

Scilab Textbook Companion for
Modern Physics
by B.L.Theraja¹

Created by
Harshit Bajpai
B.tech
Chemical Engineering
Visvesvaraya National Institute of Technology
College Teacher
Vilas K. Deshpande
Cross-Checked by
Mukul R. Kulkarni

May 23, 2016

¹Funded by a grant from the National Mission on Education through ICT, <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab codes written in it can be downloaded from the "Textbook Companion Project" section at the website <http://scilab.in>

Book Description

Title: Modern Physics

Author: B.L.Theraja

Publisher: S.Chand & Company Ltd., New Delhi

Edition: 16

Year: 2008

ISBN: 81-219-0163-4

Scilab numbering policy used in this document and the relation to the above book.

Exa Example (Solved example)

Eqn Equation (Particular equation of the above book)

AP Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

Contents

List of Scilab Codes	4
1 ELECTRIC AND MAGNETIC FIELD	5
2 THE ELECTRON	11
3 THE ATOMIC STRUCTURE	14
4 CRYSTALLOGRAPHY	19
5 QUANTUM THEORY	23
7 CLASSIFICATION OF SOLIDS	33
8 X RAYS	36
9 WAVES AND PARTICLES	45
10 THE ATOMIC NUCLEUS	51
11 NATURAL RADIOACTIVITY	55
12 ARTIFICIAL RADIOACTIVITY	59
13 NUCLEAR REACTIONS	60
14 NUCLEAR FISSION AND FUSION	63
15 NUCLEAR ENERGY SOURCES	66

List of Scilab Codes

Exa 1.1	CALCUALTION OF ELECTROSTATIC FORCE . . .	5
Exa 1.2	CALCUALTION OF DISTANCE OF SEPARATION	6
Exa 1.3	CALCULATE FIELD INTENSITY AND FORCE . . .	6
Exa 1.4	CALCULATE MASS OF OIL DROP	7
Exa 1.5	CALCULATE VELOCITY OF ELECTRON	7
Exa 1.6	CALCULATE ENERGY	8
Exa 1.7	CALCULATE TOTAL ENERGY	9
Exa 1.8	CALCULATE FORCE AND RADIUS	9
Exa 2.1	CALCULATE FORCE ACCELERATION AND KINETIC ENERGY	11
Exa 2.2	CALCULATE LINEAR VELOCITY AND RADIUS .	12
Exa 2.3	CALCULATE MASS	12
Exa 2.4	CALCULATE RADIUS AND CHARGE ON OIL DROP	13
Exa 3.1	CALCULATE DISTANCE OF CLOSEST APPROACH	14
Exa 3.2	CALCULATE VELOCITY RADIUS TIME TAKEN AND RYDBERG CONST	14
Exa 3.3	CALCULATE FREQUENCY	15
Exa 3.4	CALCULATE FREQUENCY IN FIRST ORBIT . . .	16
Exa 3.5	AT WHAT SPEED MUST ELECTRON	16
Exa 3.8	CALCULATE PRINCIPAL QUANTUM NO AND WAVE-LENGTH	17
Exa 3.9	CALCULATE WAVELENGTH FOR LYMAN SERIES	17
Exa 4.2	CALCULATE DISTANCE BETWEEN ADJACENT ATOMS	19
Exa 4.3	CALCULATE SPACING	19
Exa 4.4	CALCULATE DENSITY OF Cu	20
Exa 4.10	CALCULATION OF INTERPLANAR SPACING . .	21
Exa 4.11	CALCULATION OF ANGLE BETWEEN PAIR OF PLANES	21

Exa 4.13	CALCUALTION OF NO OF ATOMS PER SQMM . . .	22
Exa 5.1	CALCULATE ENERGY	23
Exa 5.2	HOW MANY PHOTONS	24
Exa 5.3	HOW MANY LIGHT QUANTA	24
Exa 5.4	FIND THE NO OF PHOTOELECTRONS	25
Exa 5.5	HOW MANY PHOTONS AND AT WHAT RATE . . .	25
Exa 5.6	HOW MANY PHOTONS EMITTED BY LAMP . . .	26
Exa 5.7	CALCULATE NUMBER OF PHOTONS	26
Exa 5.8	WITH WHAT VELOCITY	27
Exa 5.9	CALCULATE THRESHOLD FREQUENCY	27
Exa 5.10	CALCULATE MAXIMUM ENERGY AND WORK FUNC- TION	28
Exa 5.11	WHAT IS THRESHOLD WAVELENGTH	28
Exa 5.12	CALCULATE THRESHOLD FREQUENCY	29
Exa 5.13	WHEN VIOLET LIGHT	29
Exa 5.14	CALCULATE THRESHOLD WAVELENGTH AND PLANKS CONSTANT	30
Exa 5.15	A CERTAIN METAL	30
Exa 5.16	FIND UNKNOWN WAVELENGTH	31
Exa 5.17	LIGHT OF WAVELENGTH 2000 ANGSTROM . . .	31
Exa 5.18	CALCULATE PLANKS CONSTANT	32
Exa 7.1	CALCULATE ENERGY GAP	33
Exa 7.2	FIND MOBILITY OF ELECTRONS	33
Exa 7.3	FIND RESISTIVITY	34
Exa 7.4	CALCULATE WAVELENGTH	34
Exa 8.1	CALCULATE WAVELENGTH	36
Exa 8.2	CALCULATE NUMBER OF ELECTRONS STRIKING PER SECOND	36
Exa 8.3	CALCULATE MIN APPLIED POTENTIAL	37
Exa 8.4	CALCULATE MAX SPEED	37
Exa 8.5	CALCULATE PLANKS CONSTANT	38
Exa 8.6	CALCULATE SCREENING CONST	38
Exa 8.9	CALCULATE LINEAR ADSORPTION COEFF . . .	39
Exa 8.10	CALCULATE ANGLE OF THIRD ORDER REFLEC- TION NOTE CALCUALTION MISTAKE IN BOOK . .	39
Exa 8.11	HOW MANY ORDERS OF BRAGG RELFLECTION . .	40
Exa 8.12	CALCULATE INTERPLANAR SPACING	40
Exa 8.13	DETERMINE THE SPACING	41

Exa 8.14	DETERMINE THE TYPE OF CRYSTAL POSSESSED	41
Exa 8.15	CALCULATE SHORT WAVELENGTH AND GALNC- ING ANGLE	42
Exa 8.16	CALCULATE LATTICE SPACING OF NaCl	43
Exa 8.17	WHAT IS PRIMARY WAVELENGTH	43
Exa 8.18	WHAT IS LATTICE PARAMETER	44
Exa 9.1	WHAT IS DE BROGLIE WAVELENGTH	45
Exa 9.2	CALCULATE MOMENTUM DE BROGLIE WAVE- LENGTH AND WAVE NUMBER	45
Exa 9.3	AN ELECTRON INTIALLY AT REST	46
Exa 9.4	COMPUTE DE BROGLIE WAVELENGTH	47
Exa 9.5	WHAT IS DE BROGLIE WAVELENGTH OF NEU- TRON	47
Exa 9.6	CALCULATE THE SCATTERED WAVELENGTH	48
Exa 9.7	COMPUTE ENERGY DIFFERENCE	48
Exa 9.8	CALCULATE THE LOWEST THREE PREMISSIBLE ENERGIES	49
Exa 10.1	COMPARE DENSITIES OF WATER AND NUCLEUS	51
Exa 10.2	CALCULATE ENERGY EQUIVALENCE FOR MeV	52
Exa 10.3	CALCULATE ENERGY EQUIVALENCE FOR kWh	52
Exa 10.4	CALCULATE BINDING ENERGY	52
Exa 10.5	CALCULATE BINDING ENERGY OF $^{17}\text{Cl}35$	53
Exa 10.6	CALCULATE BINDING ENERGY FOR LITHIUM NU- CLEUS	54
Exa 11.1	CALCUALTE HOW MUCH RADIUM	55
Exa 11.2	CALCULATE RADIOACTIVE DISINTEGRATING CONST	55
Exa 11.3	COMPUTE DECAY CONST AND HALF LIFE	56
Exa 11.4	ESTIMATE HALF LIFE OF PLUTONIUM	57
Exa 11.5	CALCULATE ACTIVITY	57
Exa 11.6	HALF LIFE OF RADIUM IS 1620 YRS	58
Exa 12.1	ESTIMATE ITS AGE	59
Exa 13.1	CALCULATE THE ENERGY AVAILABLE	60
Exa 13.2	CALCULATE THE ENERGY RELEASED	60
Exa 13.3	WHAT IS THE Q VALUE OF THE REACTION	61
Exa 13.4	FIND THE MASS	61
Exa 13.6	EXPLAIN MASS DEFECT	62
Exa 14.1	COMPUTE FISSION ENERGY	63

Exa 14.2	FIND ELEMENTARY PARTICLES RELEASED IN BINARY FISSION OF $^{92}\text{U}235$	63
Exa 14.3	HOW MUCH HYDROGEN MUST BE CONVERTED INTO HELIUM	64
Exa 14.4	CALCULATE THE ENERGY LIBERATED	65
Exa 15.1	CALCULATE THE MAXIMUM FRACTION OF THE KE	66
Exa 15.2	CALCULATE THE FISSION RATE	67
Exa 15.3	HOW MANY KG OF U 235	67
Exa 15.4	HOW MUCH U 235 WOULD BE CONSUMED IN THE RUN	68
Exa 15.5	NUCLEUR REACTOR PRODUCES 200MW	69
Exa 15.6	A CITY REQUIRES 100 MW	69
Exa 15.7	BOMBAY REQUIRES 300 MWh	70
Exa 15.8	THE MOTOR OF AN ATOMIC ICE BREAKER	70
Exa 16.1	WHAT MUST BE THE FLUX DENSITY	72
Exa 16.2	WHAT IS FREQUENCY OF ALTERNATING POTENTIAL	72
Exa 16.3	A CYCLOTRON OF DEES OF RADIUS 2 METERES	73
Exa 16.4	CALCULATE THE RATIO	74
Exa 16.5	IN A CERTAIN BETATRON	74
Exa 16.6	CALCULATE MASS AND VELOCITY OF ELCTRONS	75
Exa 16.7	DETERMINE THE FREQUENCY OF GENERATOR	75
Exa 16.8	WHAT WOULD BE THE ENERGY OF ELECTRON	76
Exa 16.9	FIND THE MAX ENERGY AND CORRESPONDING WAVELENGTH	76

Chapter 1

ELECTRIC AND MAGNETIC FIELD

Scilab code Exa 1.1 CALCUALTION OF ELECTROSTATIC FORCE

```
1  clc;clear;
2  //Example 1.1
3  //Comparision of electrostatic force of replusion
       between two particles with the gravitational
       forces between them
4
5  //given values
6  q1=3.2*10^-19;
7  q2=q1;//q1 and q2 are the values of charge on alpha-
   particle in C
8  d=10^-13;//distance between two alpha-particles in m
9  m1=6.68*10^-27;
10 m2=m1;//m1 and m2 are masses of alpha-particles in
    kg
11 G=6.67*10^-11;//Gravitational constant in N-(m^2)/(
    kg^2)
12
13 //calculation
14 F1=(9*10^9)*(q1*q2)/(d^2);//calculation of
```

```

        electrostatic force
15 disp(F1, 'The electrosatic force(in N) is ');
16 F2=G*(m1*m2)/(d^2); //calculation of electrostatic
        force
17 disp(F2, 'The gravitational force (in N) is ')

```

Scilab code Exa 1.2 CALCUALTION OF DISTANCE OF SEPARATION

```

1  clc;clear;
2  //Example 1.2
3  //Calculation of the distance of separation between
        two electrons
4
5  // given values
6  m=9.1*10^-31; //mass of elctron in kg
7  q=1.6*10^-19; //charge on electron in C
8  g=9.81; //acceleration due to gravity in m/(s^2)
9
10 //calculation
11 Fg=m*g; //gravitational force in N
12 d=sqrt((9*10^9*q^2)/Fg); //equating gravitational
        force with electrosatic force for calculating
        distance
13 disp(d, 'The distance of separation(in m) is ')

```

Scilab code Exa 1.3 CALCULATE FIELD INTENSITY AND FORCE

```

1  clc;clear;
2  //Example 1.3
3  //Calculation of electric intensity between plates
        and force on proton
4
5  //given values

```

```

6 d=0.02//distance between plates in m
7 V=400;//potential difference of plates in V
8 q=1.6*10^-19;//charge on a proton in C
9
10 //calculation
11 E=V/d;//
12 disp(E,'The electric field intensity(in V/m) between
    plates is ');
13 F=q*E;//
14 disp(F,'The force(in N) on proton is ')

```

Scilab code Exa 1.4 CALCULATE MASS OF OIL DROP

```

1 clc;clear;
2 //Example 1.4
3 //calculation of mass of oil drop
4
5 //given values
6 d=0.02//distance between plates in m
7 q=1.6*10^-19;//charge on oil drop in C
8 V=6000;//potential difference of plates in V
9 g=9.81;//acceleration due to gravity in m/(s^2)
10
11 //calculation
12 E=V/d;//electric field intensity between plates in V
    /m
13 F=q*E;//electrostatic force on oil drop in N
14 m=F/g;//equating the weight of oil drop to the
    electrostatic force on it
15 disp(m,'The mass(in kg) of oil drop')

```

Scilab code Exa 1.5 CALCULATE VELOCITY OF ELECTRON

```

1  clc;clear;
2  //Example 1.5
3  //Calculation of velocity of an electron
4
5  //given values
6  V=150;//potential difference between anode and
   cathode in V
7  m=9.31*10-31;//mass of an electron in kg
8  q=1.6*10-19;//charge on an electron in C
9
10 //Calculation
11 E=q*V;//energy(in J) gained by electron during
   speeding from cathode to anode
12 vel=sqrt(E*2/m);//equating with kinetic energy of
   electron i.e m(v2)/2
13 disp(vel,'The velocity(in m/s) is ')

```

Scilab code Exa 1.6 CALCULATE ENERGY

```

1  clc;clear;
2  //Example 1.6
3  //Calculation of energy in eV,MeV,J
4
5  //given values
6  V=5*106;//potential difference through which alpha-
   particle is accelerated
7  e=1.6*10-19;//charge on electron in C
8
9  //calculation
10 E1=2*V;//electronic charge on alpha-particle is 2e
   Coulombs
11 disp(E1,'The energy in eV is ');
12 E2=E1/106;//energy in MeV
13 disp(E2,'The energy in MeV is ');
14 E3=E1*e;//energy in Joules

```

```
15 disp(E3, 'The energy in J is')
```

Scilab code Exa 1.7 CALCULATE TOTAL ENERGY

```
1 clc;clear;
2 //Example 1.7
3 //Calculation of the total energy
4
5 //given values
6 r=0.528*10^-10;//radius of the orbit in m
7 q=-1.6*10^-19;//charge on electron in C
8 Q=1.6*10^-19;//charge on Hydrogen nucleus in C
9 Eo=8.854*10^-12;//permittivity in free space in F/m
10
11 //calculation
12 E=(q*Q)/(8*3.14*Eo*r);//
13 disp(E, 'The total energy(in J) is');
14 E1=E/(1.6*10^-19);//
15 disp(E1, 'The total energy(in eV) is')
```

Scilab code Exa 1.8 CALCULATE FORCE AND RADIUS

```
1 clc;clear;
2 //Example 1.8
3
4 //given values
5 Q=3.2*10^-19;//charge on alpha-particle in C
6 m=6.68*10^-27;//mass on alpha-particle in kg
7 B=1.5;//transverse magnetic field of flux density in
  Wb/(m^2)
8 v=5*10^6;//velocity of alpha-particle in m/s
9
10 //Calculation
```

```
11 F=B*Q*v; //
12 disp(F, 'The force(in N) on particle is ');
13 R=m*v/(Q*B); //
14 disp(R, 'The radius(in m) of its circular path')
```

Chapter 2

THE ELECTRON

Scilab code Exa 2.1 CALCULATE FORCE ACCELERATION AND KINETIC ENERGY

```
1  clc;clear;
2  //Example 2.1
3
4  //given values
5  E=2400;//electric field intensity in V/m
6  V=90;//potential difference in V
7  e=1.6*10^-19;//the charge on electron in C
8  m=9.12*10^-31;//mass of electron in kg
9
10 //Calculation
11 F=e*E;
12 disp(F,'The force(in N) on electron is');
13 a=F/m;
14 disp(a,'Its acceleration (in m/s^2)');
15 KE=e*V;
16 disp(KE,'The Kinetic Energy(in J) of particle is');
17 v=sqrt(2*KE/m);
18 disp(v,'The velocity(in m/s) of the electron')
```

Scilab code Exa 2.2 CALCULATE LINEAR VELOCITY AND RADIUS

```
1  clc;clear;  
2  //Example 2.2  
3  
4  //given values  
5  V=900;//potential difference in V  
6  B=0.01;//uniform magnetic field in Wb/m^2  
7  em=1.76*10^11;//value of e/m in C/kg  
8  
9  //calculation  
10 v=sqrt(2*em*V);  
11 disp(v,'The linear velocity(in m/s) of electron is')  
    ;  
12 R=v/(em*B);  
13 disp(R,'The radius(in m) of the circular path is')
```

Scilab code Exa 2.3 CALCULATE MASS

```
1  clc;clear;  
2  //Example 2.3  
3  
4  //given values  
5  d=6*10^-3;//distance between plates in m  
6  V=900;//potential difference in V  
7  B=0.5;//uniform magnetic field in Wb/m^2  
8  Q=1.6*10^-19;//the charge on electron in C  
9  R=10.6*10^-2;//circular track radius in m  
10  
11 //calculation  
12 v=V/(B*d);  
13 m=R*Q*B/v;
```

```
14 disp(m, 'The mass(in kg) of particle')
```

Scilab code Exa 2.4 CALCULATE RADIUS AND CHARGE ON OIL DROP

```
1  clc;clear;
2  //Example 2.4
3
4  //given values
5  V=6920; //potential difference in V
6  d=1.3*10^-3; //distance between in m
7  v=1.9*10^-4; //velocity in m/s
8  p=0.9*10^3; //density of oil in kg/m^3
9  n=1.81*10^-5; //coefficient of viscosity in N-s/m^2
10 g=9.81; //accelaration due to gravity in m/s^2
11 pi=3.14; //standard constant
12
13 //calculation
14 a=sqrt((9*n*v)/(2*g*p));
15 disp(a, 'The radius(in m) of the drop is');
16 E=V/d;
17 Q=4*pi*(a^3)*p*g/(3*E);
18 disp(Q, 'The value of charge(in C) on oil drop is')
```

Chapter 3

THE ATOMIC STRUCTURE

Scilab code Exa 3.1 CALCULATE DISTANCE OF CLOSEST APPROACH

```
1  clc;clear;
2  //Example 3.1
3
4  //given values
5  Z=79;//atomic number of gold
6  e=1.6*10^-19;//electron charge in C
7  Eo=8.854*10^-12;//absolute permittivity of free space
   in F/m
8  K=7.68*1.6*10^-13;//kinectic energy in J
9  pi=3.14;//standard constant
10
11 //calculations
12 D=(2*Z*e^2)/(4*pi*Eo*K);
13 disp(D,'The closest distance(in m) of approach is')
```

Scilab code Exa 3.2 CALCULATE VELOCITY RADIUS TIME TAKEN
AND RYDBERG CONST

```

1  clc;clear;
2  //Example 3.2
3
4  //given values
5  Z=1;//atomic number of hydrogen
6  e=1.6*10^-19;//electron charge in C
7  pi=3.14;//standard constant
8  h=6.625*10^-34;//plank's constant in J-s
9  m=9.1*10^-31;//mass of an electron in kg
10 Eo=8.854*10^-12;//absolute permitivity of free space
    in F/m
11 c=3*10^8;//speed of light in m/s
12 n=1;//ground state
13
14 //calculation
15 v=9*10^9*(2*pi*Z*e^2)/(n*h);
16 disp(v,'velocity(in m/s) of ground state');
17 r=(Eo*n^2*h^2)/(pi*m*e^2);
18 disp(r,'radius(in m) of Bohr orbit in ground state')
19 t=(2*pi*r)/v;
20 disp(t,'time taken(in s) by electron to traverse the
    bohr first orbit');
21 R=(m*e^4)/(8*Eo^2*h^3*c);
22 disp(R,'Rhydberg constant (in m^-1)')

```

Scilab code Exa 3.3 CALCULATE FREQUENCY

```

1  clc;clear;
2  //Example 3.3
3
4  //given values
5  B=2.179*10^-16;//a constant in J
6  h=6.625*10^-34;//plank's constant in J-s
7
8  //calculation

```

```

9 E3=-B/3^2;
10 E2=-B/2^2;
11 f=(E3-E2)/h;
12 disp(f, 'frequency (in Hz) of radiation ')

```

Scilab code Exa 3.4 CALCULATE FREQUENCY IN FIRST ORBIT

```

1 clc;clear;
2 //Example 3.4
3
4 //given values
5 Z=1;//atomic number of hydrogen
6 e=1.6*10^-19;//electron charge in C
7 h=6.625*10^-34;//plank's constant in J-s
8 m=9.1*10^-31;//mass of an electron in kg
9 Eo=8.854*10^-12;//absolute permittivity of free space
   in F/m
10 n=1;//ground state
11
12 //Calculations
13 f=(m*Z^2*e^4)/(4*Eo^2*h^3);
14 disp(f, 'the frequency (in Hz) is ')

```

Scilab code Exa 3.5 AT WHAT SPEED MUST ELECTRON

```

1 clc;clear;
2 //Example 3.5
3
4 //given data
5 Z=1;
6 n=1;
7 e=1.6*10^-19;//the charge on electron in C
8 h=6.625*10^-34;//Plank's constant

```

```

9 Eo=8.854*10^-12;//absolute permitivity of free space
  in F/m
10 m=9.12*10^-31;//mass of electron in kg
11
12 //calculations
13 v=Z*e^2/(2*Eo*n*h);
14 disp(v,'velcocity in m/s');
15 E=-m*Z^2*e^4/(8*(Eo*n*h)^2);
16 disp(E,'energy of hydrogen atom in J');
17 f=m*Z^2*e^4/(4*Eo^2*(n*h)^3);
18 disp(f,'frequecy in Hz')

```

Scilab code Exa 3.8 CALCULATE PRINCIPAL QUANTUM NO AND WAVELENGTH

```

1  clc;clear;
2  //Example 3.8
3
4  //given data
5  h=6.625*10^-34;//Plank's constant
6  c=3*10^8;//speed of light in m/s
7  E1=10.2;//in eV energy
8  E2=12.09;//in eV energy
9  e=1.6*10^-19;//the charge on electron in C
10
11 //calcualtion
12 //principal quantum no are 2 & 3 respectively
13 W=c*h/(E1*e)*10^10;
14 disp(W,'wavelength in angstrom is for 10.2 eV');
15 W=c*h/(E2*e)*10^10;
16 disp(W,'wavelength in angstrom is for 12.09 eV')

```

Scilab code Exa 3.9 CALCULATE WAVELENGTH FOR LYMAN SERIES

```
1  clc;clear;
2  //Example 3.9
3
4  //given data
5  R=10967700; //Rydberg constant in 1/m
6
7  //calculation
8  W1=4/(3*R); //as n1=1 and n2=2
9  disp((W1*10^10), 'Long wavelength in angstrom');
10 W2=1/R; //as n1=1 and n2=infinity
11 disp((W2*10^10), 'Short wavelength in angstrom')
```

Chapter 4

CRYSTALLOGRAPHY

Scilab code Exa 4.2 CALCULATE DISTANCE BETWEEN ADJACENT ATOMS

```
1  clc; clear;
2  //Example 4.2
3
4  //given data
5  d=2180; //density of NaCl
6  M=23.5+35.5; //Molecular weight
7  Na=6.02*10^26; //Avgrado no. in 1/kg mole
8  n=4; //for f.c.c
9
10 //calculations
11 a=(n*M/(Na*d))^(1/3);
12 d=a/2;
13 disp((d*10^10), 'distance between to adajcent atoms
    in angstrom')
```

Scilab code Exa 4.3 CALCULATE SPACING


```

1  clc;clear;
2  //Example 4.3
3
4  //given data
5  d=2.163;//density in gm/cm^3
6  M=58.45;
7  Na=6.02*10^23;//Avgraodo no. in 1/gm mole
8
9  //calcualtions
10 n=Na/M;//no. of molecules/gram
11 n=n*d;//no. of molecules/cm^3
12 n=2*n;//no. of atom/cm^3;
13 n=n^(1/3);//no. of atoms in a row 1cm long
14 d1=1/n;
15 disp((d1*10^8),'Spacing in angstrom')

```

Scilab code Exa 4.4 CALCULATE DENSITY OF Cu

```

1  clc;clear;
2  //Example 4.4
3
4  //given data
5  r=1.278;//radius in A.U
6  n=4;//structure is f.c.c
7  M=63.54;
8  Na=6.02*10^23;//Avgraodo no. in 1/gm mole
9
10 //calculations
11 a=4*r/(sqrt(2));
12 V=a^3;
13 d=n*M/(Na*V);
14 disp((d*(10^8)^3),'Density in g/cm^3')

```

Scilab code Exa 4.10 CALCULATION OF INTERPLANAR SPACING

```
1  clc; clear;
2  //Example 4.10
3
4  //given data
5  r=1.746; //atomic radius in angstrom
6
7  //calculations
8  a=4*r/sqrt(2);
9  //for (i)
10 h=2;k=0;l=0;
11 d=a/sqrt(h^2+k^2+l^2);
12 disp(d, 'distace for (200) in A.U');
13 //for (ii)
14 h=2;k=2;l=0;
15 d=a/sqrt(h^2+k^2+l^2);
16 disp(d, 'distace for (220) in A.U ');
17 //for (iii)
18 h=1;k=1;l=1;
19 d=a/sqrt(h^2+k^2+l^2);
20 disp(d, 'distace for (111) in A.U')
```

Scilab code Exa 4.11 CALCULATION OF ANGLE BETWEEN PAIR OF PLANES

```
1  clc; clear;
2  //Example 4.11
3
4  //calculations
5  //for (i)
6  l=1;m=0;n=0;
7  p=0;q=1;r=0;
8  d=acosd((l*p+m*q+n*r)/(sqrt(l^2+m^2+n^2)*sqrt(p^2+q^2+r^2)));
```

```

9  disp(d, 'angle b/w pair of miller indices in (i)');
10 //for (ii)
11 l=1;m=2;n=1;
12 p=1;q=1;r=1;
13 d=acosd((l*p+m*q+n*r)/(sqrt(l^2+m^2+n^2)*sqrt(p^2+q
    ^2+r^2)));
14 disp(d, 'angle b/w pair of miller indices in (ii)')

```

Scilab code Exa 4.13 CALCUALTION OF NO OF ATOMS PER SQMM

```

1  clc;clear;
2  //Example 4.13
3
4  //given data
5  a=3.61*10^-7;//lattice constant in nm
6
7  //calcualtions
8  //for (i) plane (100)
9  SA=a*a;
10 tamc=2;//total atoms included according to sketch
11 ans=tamc/SA;
12 disp(ans, 'atoms per mm^2 for (i)');
13 //for (ii) plane (110)
14 A=a*(sqrt(2)*a);
15 tamc=2;//total atoms included according to sketch
16 ans=tamc/A;
17 disp(ans, 'atoms per mm^2 for (ii)');
18 //for (iii) plane (111)
19 A=0.866*a*a;
20 tamc=2;//total atoms included according to sketch
21 ans=tamc/A;
22 disp(ans, 'atoms per mm^2 for (iii)')

```

Chapter 5

QUANTUM THEORY

Scilab code Exa 5.1 CALCULATE ENERGY

```
1  clc;clear;
2  //Example 5.1
3
4  //given values
5  W1=4;//wavelength in Angstrom
6  W2=1;//wavelength in Angstrom
7  e=1.6*10^-19;//the charge on electron in C
8  m=9.12*10^-31;//mass of electron in kg
9
10 //calculation
11 disp("Part (i)");
12 E=12400/W1;
13 disp(E,'The energy in eV is ');
14 v=sqrt(E*e*2/m);
15 disp(v,'The velocity in m/s is ');
16 disp("Part (ii)");
17 E=12400/W2;
18 disp(E,'The energy in eV is ');
19 v=sqrt(E*e*2/m);
20 disp(v,'The velocity in m/s is ')
```

Scilab code Exa 5.2 HOW MANY PHOTONS

```
1  clc;clear;  
2  //Example 5.2  
3  
4  //given values  
5  f=880*103; //frequency in Hz  
6  P=10*103; //Power in W  
7  h=6.625*10-34; //Plank's constant  
8  
9  //calculation  
10 E=h*f;  
11 n=P/E;  
12 disp(n, 'The number of photons emitted per second are  
    ')
```

Scilab code Exa 5.3 HOW MANY LIGHT QUANTA

```
1  clc;clear;  
2  //Example 5.3  
3  
4  //given values  
5  P=200; //power in W  
6  W=6123*10-10; //wavelength in m  
7  c=3*108; //speed of light in m/s  
8  h=6.625*10-34; //Plank's constant  
9  
10 //calculation  
11 Op=0.5*P; //radiant o/p  
12 E=h*c/W;  
13 n=2/E;  
14 disp(n, 'No. of Quanta emitted/s')
```

Scilab code Exa 5.4 FIND THE NO OF PHOTOELECTRONS

```
1 clc; clear;
2 //Example 5.4
3
4 //given values
5 N=5*104; //no. of photons
6 W=3000*10-10; //wavelength in A
7 J=5*10-3; //sensitivity for W in A/W
8 h=6.625*10-34; //Plank's constant
9 c=3*108; //speed of light in m/s
10 e=1.6*10-19; //the charge on electron in C
11
12 //calculation
13 E=h*c/W; //energy content of each photon
14 TE=N*E; //total energy
15 I=J*TE; //current produced
16 n=I/e;
17 disp(n, 'no. photoelectrons emitted are')
```

Scilab code Exa 5.5 HOW MANY PHOTONS AND AT WHAT RATE

```
1 clc; clear;
2 //Example 5.5
3
4 //given values
5 W=5*10-7; //wavelength in m
6 F=10-5; //force in N
7 h=6.625*10-34; //Plank's constant
8 m=1.5*10-3; //mass in kg
9 c=3*108; //speed of light in m/s
```

```

10 S=0.1//specific heat
11
12 //calculation
13 n=F*W/h;
14 disp(n, 'no. of photons ');
15 E=F*c/4200;//in kcal/s
16 T=E/(m*S);
17 disp(T, 'the rate of temperature rise ')

```

Scilab code Exa 5.6 HOW MANY PHOTONS EMITTED BY LAMP

```

1 clc;clear;
2 //Example 5.6
3
4 //given values
5 W=4500*10^-10;//wavelength in m
6 V=150;//rated voltage in W
7 h=6.625*10^-34;//Plank's constant
8 c=3*10^8;//speed of light in m/s
9
10 //calculation
11 P=V*8/100;//lamp power emitted
12 E=h*c/W;
13 n=P/E;
14 disp(n, 'No. photons emitted/s ')

```

Scilab code Exa 5.7 CALCULATE NUMBER OF PHOTONS

```

1 clc;clear;
2 //Example 5.7
3
4 //given values
5 f=1*10^12;//frequency in Hz

```

```

6 h=6.625*10^-34; //Plank 's constant
7
8 //calculation
9 E=h*f;
10 n=E/6.625;
11 disp(n, 'the no. of photons required')

```

Scilab code Exa 5.8 WITH WHAT VELOCITY

```

1 clc;clear;
2 //Example 5.8
3
4 //given values
5 W=5200*10^-10; //wavelength in m
6 h=6.625*10^-34; //Plank 's constant
7 m=9.12*10^-31; //mass of electron in kg
8
9 //calculations
10 p=h/W;
11 v=p/m;
12 disp(v, 'the velocity in m/s')

```

Scilab code Exa 5.9 CALCULATE THRESHOLD FREQUENCY

```

1 clc;clear;
2 //Example 5.9
3
4 //given data
5 v=7*10^5; //maximum speed in m/sec
6 f=8*10^14; //frequency in Hz
7 h=6.625*10^-34; //Plank 's constant
8 c=3*10^8; //speed of light in m/s
9 m=9.12*10^-31; //mass of electron in kg

```



```

10
11 //calculations
12 E=0.5*m*v*v;
13 fo=f-(E/h);
14 disp(fo,'the threshold frequency in Hz is')

```

Scilab code Exa 5.10 CALCULATE MAXIMUM ENERGY AND WORK FUNCTION

```

1  clc;clear;
2  //Example 5.10
3
4  //given data
5  Wo=2300*10;//threshold wavelength in Angstrom
6  W=1800*10;//incident light wavelength in Angstrom
7
8  //calculations
9  w=124000/Wo;
10 disp(w,'The work function in eV is');
11 E=124000*((1/W)-(1/Wo));
12 disp(E,'The maximum energy in eV')

```

Scilab code Exa 5.11 WHAT IS THRESHOLD WAVELENGTH

```

1  clc;clear;
2  //Example 5.11
3
4  //given data
5  W=6000;//wavelegth in Angstrom
6  v=4*10^5;//velocity in m/sec
7  m=9.12*10^-31;//mass of electron in kg
8  e=1.6*10^-19;//the charge on electron in C
9

```

```

10 //calculations
11 KE=0.5*m*v^2/e;
12 disp(KE,'The Kinetic energy in eV is ');
13 WF=12400/W;
14 Wo=12400/(WF-KE);
15 disp(Wo,'The threshold wavelength in Angstrom is ')

```

Scilab code Exa 5.12 CALCULATE THRESHOLD FREQUENCY

```

1  clc;clear;
2  //Example 5.12
3
4  //given data
5  Wo=4.8;//work function in eV
6  W=2220;//wavelength in angstrom
7
8  //calculations
9  E=12400/W;
10 Emax=E-Wo;
11 disp(Emax,'The maximum Kinetic energy in eV is ')

```

Scilab code Exa 5.13 WHEN VIOLET LIGHT

```

1  clc;clear;
2  //Example 5.13
3
4  //given data
5  W=4000*10^-10;//wavelength in m
6  Vs=0.4;//retarding potential in eV
7  h=6.625*10^-34;//Plank's constant
8  c=3*10^8;//speed of light in m/s
9  e=1.6*10^-19;//the charge on electron in C
10

```

```

11 //calculations
12 f=c/W;
13 disp(f, 'The light frequency in Hz');
14 E=h*f/e;
15 disp(E, 'The photon energy in eV');
16 Wo=E-Vs;
17 disp(Wo, 'The work function in eV');
18 fo=Wo/h*e;
19 disp(fo, 'The threshold frequency in Hz');
20 NE=(E-Wo)*e;
21 disp(NE, 'The net energy in J')

```

Scilab code Exa 5.14 CALCULATE THRESHOLD WAVELENGTH AND PLANKS CONSTANT

```

1 clc;clear;
2 //Example 5.14
3
4 //given data
5 W1=3310*10^-10;//photon wavelength in m
6 W2=5000*10^-10;//photon wavelength in m
7 E1=3*10^-19;//electron energy in J
8 E2=0.972*10^-19;//electron energy in J
9 c=3*10^8;//speed of light in m/s
10
11 //given values
12 h=(E1-E2)*(W1*W2)/(c*(W2-W1));
13 disp(h, 'the plancks const in Js');
14 Wo=c*h/E1;
15 disp(Wo, 'The threshold wavelength in m')

```

Scilab code Exa 5.15 A CERTAIN METAL

```

1  clc;clear;
2  //Example 5.15
3
4  //given data
5  W=6525;//wavelength in angstrom
6
7  //calcualation
8  Vo=12400*((1/4000)-(1/W));
9  disp(Vo,'Stopping potential in (a) in volts');
10 Vo=12400*((1/2000)-(1/W));
11 disp(Vo,'Stopping potential in (b) in volts');
12 Vo=12400*((1/2000)-(2/W));
13 disp(Vo,'Stopping potential in (c) in volts')

```

Scilab code Exa 5.16 FIND UNKNOWN WAVELENGTH

```

1  clc;clear;
2  //EXample 5.16
3
4  //given data
5  Wo=5000;//wavelength in angstrom
6  V=3.1;//stopping potential in V
7
8  //calcualtion
9  W=1/((V/12400)+(1/Wo));
10 disp(W,'The unknown wavelength in Angstrom')

```

Scilab code Exa 5.17 LIGHT OF WAVELENGTH 2000 ANGSTROM

```

1  clc;clear;
2  //Example 5.17
3
4  //given values

```

```

5 W=2000; //wavelength in Angstrom
6 Vs=4.2; //Work Function in eV
7 e=1.6*10^-19; //the charge on electron in C
8
9 //calculations
10 E=12400/W;
11 Emax=(E-Vs)*e;
12 disp(Emax, 'KE of fastest photoelectron in J');
13 Emin=0;
14 disp(Emin, 'KE of slowest moving electron in J');
15 Vo=Emax/e;
16 disp(Vo, 'Stopping potential in V');
17 Wo=12400/Vs;
18 disp(Wo, 'The cutoff wavelength in Angstrom')

```

Scilab code Exa 5.18 CALCULATE PLANKS CONSTANT

```

1 clc;clear;
2 //Example 5.18
3
4 //given values
5 Vs1=4.6; //Stopping Potential in V
6 Vs2=12.9; //Stopping Potential in V
7 f1=2*10^15; //frequency in Hz
8 f2=4*10^15; //frequency in Hz
9 e=1.6*10^-19; //the charge on electron in C
10
11 //calculations
12 h=((Vs2-Vs1)*e)/(f2-f1)
13 disp(h, 'The Plancks const in Js')

```

Chapter 7

CLASSIFICATION OF SOLIDS

Scilab code Exa 7.1 CALCULATE ENERGY GAP

```
1 clc;clear;
2 //Example 7.1
3
4 //given data
5 W=11000;//wavelength in angrstrom
6
7 //calcuations
8 Eg=W/12400;
9 disp(Eg,'Energy Gap in eV')
```

Scilab code Exa 7.2 FIND MOBILITY OF ELECTRONS

```
1 clc;clear;
2 //Example 7.2
3
4 //given data
```

```

5 p=1.7*10^-6; //resistivity in ohm-cm
6 d=8.96; //density in gm/cc
7 W=63.5;
8 Na=6.02*10^23; //Avgrado no. in 1/g mole
9 e=1.6*10^-19; //the charge on electron in C
10
11 //calcualtions
12 n=8.96*Na/W;
13 ue=1/(p*e*n);
14 disp(ue, 'mobilty of electrons in cm^2/V-s');

```

Scilab code Exa 7.3 FIND RESISTIVITY

```

1 clc;clear;
2 //Example 7.3
3
4 //given data
5 d1=2.5*10^19; //density of charge carries in 1/m^3
6 d2=4.2*10^28; //density of germanium atoms
7 ue=0.36; //mobilty of electrons in m^2/V-s
8 Na=6.02*10^23; //Avgrado no. in 1/g mole
9 e=1.6*10^-19; //the charge on electron in C
10
11 //calcualtions
12 Nd=d2/10^6;
13 cn=Nd*e*ue;
14 pn=1/cn;
15 disp(pn, 'resistivity of doped germanium in ohm-m')

```

Scilab code Exa 7.4 CALCULATE WAVELENGTH

```

1 clc;clear;
2 //Example 7.4

```

```
3
4 //given data
5 Eg=0.75;//energy gap in eV
6
7 //calcualtions
8 W=12400/Eg;
9 disp(W, 'wavelength in angstrom')
```

Chapter 8

X RAYS

Scilab code Exa 8.1 CALCULATE WAVELENGTH

```
1 clc; clear;  
2 //Example 8.1  
3  
4 //given data  
5 V=60000; //working voltage in V  
6  
7 //calculation  
8 Wmin=12400/V;  
9 disp(Wmin, 'Wavelength emitted in Angstrom')
```

Scilab code Exa 8.2 CALCULATE NUMBER OF ELECTRONS STRIKING PER SECOND

```
1 clc; clear;  
2 //Example 8.2  
3  
4 //given data  
5 V=12400; //Volatage applied in V
```

```

6 I=0.002;//current drop in A
7 e=1.6*10^-19;//the charge on electron in C
8
9 //calculations
10 n=I/e;
11 disp(n,'No. of electrons');
12 v=(5.93*10^5)*(sqrt(V));
13 disp(v,'the speed with which they strike in m/s');
14 Wmin=12400/V;
15 disp(Wmin,'shortest wavelength in Angstrom')

```

Scilab code Exa 8.3 CALCULATE MIN APPLIED POTENTIAL

```

1 clc;clear;
2 //Example 8.3
3
4 //given values
5 Wmin=1;//shortest wavelength in Angstrom
6
7 //calculations
8 V=(12400/Wmin)/1000;
9 disp(V,'The minimum applied voltage in kV')

```

Scilab code Exa 8.4 CALCULATE MAX SPEED

```

1 clc;clear;
2 //Exmaple 8.4
3
4 //given data
5 I=0.005;//current in A
6 V=100000;//potential difference in V
7
8 //calcualtions

```

```

9 v=(5.98*10^5)*(sqrt(V));
10 disp(v, 'Maximum speed in m/s');
11 IP=V*I; //incident power in W
12 P=.999*IP; //power converted into heat in W
13 H=P/4.18;
14 disp(H, 'The heat produced/second in cal/s');

```

Scilab code Exa 8.5 CALCULATE PLANKS CONSTANT

```

1 clc;clear;
2 //Example 8.5
3
4 //given data
5 V=30000; //potential difference in V
6 Wmin=0.414*10^-10; //short wavelength limit in m
7 e=1.602*10^-19; //the charge on electron in C
8 c=3*10^8; //speed of light in m/s
9
10 //calcualtions
11 h=(e*V*Wmin)/c;
12 disp(h, 'The Plancks const in Js');

```

Scilab code Exa 8.6 CALCULATE SCREENING CONST

```

1 clc;clear;
2 //Example 8.6
3
4 //given data
5 W=1.43*10^-10; //wavelength in m
6 Z=74; //atomic no
7 R=10.97*10^6; //Rydberg constant in 1/m
8
9 //calcualation

```

```
10 b=74-sqrt(36/(5*R*W)); //Transition from to M to L
11 disp(b, 'the screening const.')
```

Scilab code Exa 8.9 CALCULATE LINEAR ADSORPTION COEFF

```
1  clc;clear;
2  //Example 8.9
3
4  //given data
5  um=0.6; //mass adsoption coeffcient in cm^2/g
6  p=2.7; //density of aluminium in g/cm^3
7
8  //calculations
9  u=p*um;
10 disp(u, 'linear adsorption coeffcient of aluminium in
    1/cm');
11 T=0.693/u
12 disp(T, 'the hvl in cm');
13 x=(log(20))*(1/u);
14 disp(x, 'the thickness in cm')
```

Scilab code Exa 8.10 CALCULATE ANGLE OF THIRD ORDER REFLECTION NOTE CALCUALTION MISTAKE IN BOOK

```
1      clc;clear;
2  //Example 8.10
3
4  //given values
5  D=12; //glancing angle in Degree
6  n=1;
7  d=3.04*10^-10; //grating space in m
8
9  //calculation
```

```

10 W=(2*d*sind(D));
11 disp((W/(10^-10)), 'the wavelength in Angstrom');
12 D3=asind((3*W)/(2*d));
13 disp(D3, 'the angle for third order reflection in
    degrees')

```

Scilab code Exa 8.11 HOW MANY ORDERS OF BRAGG RELFLECTION

```

1  clc;clear;
2  //Example 8.11
3
4  //given data
5  d=1.181; //distance of seperation in Angstrom
6  W=1.540; //wavelength in Angstrom
7
8  //calculations
9  n=2*d/W; //sin(D) = 1 for max value
10 disp(n, 'the orders of bragg reflection')

```

Scilab code Exa 8.12 CALCULATE INTERPLANAR SPACING

```

1  clc;clear;
2  //Example 8.12
3
4  //given data
5  W=0.6; //wavelength in angstrom
6  D1=6.45;
7  D2=9.15;
8  D3=13; //angles in degree
9
10 //calculation
11 d=W/(2*sind(D1));

```

```

12 disp(d, 'interplanar spacing for (a) in angstrom');
13 d=W/(2*sind(D2));
14 disp(d, 'interplanar spacing for (b) in angstrom');
15 d=W/(2*sind(D3))*2; //n=2 for (c)
16 disp(d, 'interplanar spacing for (c) in angstrom')

```

Scilab code Exa 8.13 DETERMINE THE SPACING

```

1 clc;clear;
2 //Example 8.13
3
4 //given data
5 W=3*10^-10; //wavelength in m
6 D=40; //angle in degree
7 n=1;
8
9 //calculation
10 d=n*W/(2*sind(D));
11 disp((d/10^-10), 'spacing in AU')
12 a=2*d;
13 v=a^3;
14 disp(v, 'the volumne in m^3 is')

```

Scilab code Exa 8.14 DETERMINE THE TYPE OF CRYSTAL POSSESED

```

1 clc;clear;
2 //Example 8.14
3
4 //given data
5 D1=5.4;
6 D2=7.6;
7 D3=9.4; //angles in degree

```

```

8
9 //calcualtion
10 d1=1/(2*sind(D1));
11 d2=1/(2*sind(D2));
12 d3=1/(2*sind(D3));
13 m=min(d1,d2,d3);
14 d1=d1/m;
15 d2=d2/m;
16 d3=d3/m;
17 disp(d1,d2,d3,'d1:d2:d3 =')

```

Scilab code Exa 8.15 CALCULATE SHORT WAVELENGTH AND GAL-
NCING ANGLE

```

1 clc;clear;
2 //Example 8.15
3
4 //given data
5 V=50000;//applied voltage in V
6 p=1.99*10^3;//density in kg/m^3
7 n=4;
8 Na=6.02*10^26;//Avgraodo no. in 1/kg mole
9 M=74.6;//molecular mass
10 W=0.248*10^-10;//wavelength in m
11
12 //calculation
13 Wmin=12400/V;
14 disp(Wmin,'short wavelength limit');
15 a=(n*M/(Na*p))^(1/3);
16 d=a/2;
17 D=asind(W/(2*d));
18 disp(D,'glancing angle in degree')

```

Scilab code Exa 8.16 CALCULATE LATTICE SPACING OF NaCl

```
1  clc; clear;
2  //Example 8.16
3
4  //given data
5  W=1.54; //wavelength in angstrom
6  D=15.9; //angle in degree
7  M=58.45 //molecular weight
8  p=2164*103; //density in kg/m3
9  n=2; //for NaCl molecule
10
11 //calculation
12 d=W/(2*sind(D));
13 disp(d, 'lattice spacing in angstrom');
14 d=d*10-10;
15 Na=M/(2*d3*p);
16 disp(Na, 'Avogrado number in 1/gm mole')
```

Scilab code Exa 8.17 WHAT IS PRIMARY WAVELENGTH

```
1  clc; clear;
2  //Example 8.17
3
4  //given data
5  D=60; //angle in degree
6  W=0.254; //wavelength in angstrom
7
8  //calcualtion
9  dW=0.024*(1-cosd(D));
10 W1=W-dW;
11 disp(W1, 'primary X-ray wavelength in angstrom')
```

Scilab code Exa 8.18 WHAT IS LATTICE PARAMETER

```
1  clc; clear;
2  //Example 8.18
3
4  //given data
5  D=32; //angle in degree
6  W=1.54*10^-10; //wavelength in angstrom
7  h=2; k=2; l=0; //lattice consts
8
9  //calculations
10 d=W/(2*sind(D));
11 a=d*sqrt(h^2+k^2+l^2);
12 disp(a, 'lattice parameter in m');
13 r=sqrt(2)*a/4;
14 disp(r, 'radius of atom in m')
```

Chapter 9

WAVES AND PARTICLES

Scilab code Exa 9.1 WHAT IS DE BROGLIE WAVELENGTH

```
1 clc; clear;  
2 //Example 9.1  
3  
4 //given data  
5 V=20000; //applied voltage in V  
6  
7 //calculation  
8 W=12.25/(sqrt(V));  
9 disp(W, 'de broglie wavelength in angstrom')
```

Scilab code Exa 9.2 CALCULATE MOMENTUM DE BROGLIE WAVELENGTH AND WAVE NUMBER

```
1 clc; clear;  
2 //Example 9.2  
3  
4 //given data  
5 V=5000; //applied voltage in V
```

```

6 e=1.602*10^-19;//the charge on electron in C
7 m=9.12*10^-31;//mass of electron in kg
8 d=2.04*10^-10;//distance in m
9
10 //calculations
11 p=sqrt(2*m*e*V);
12 disp(p, 'momentum in kg-m/s^2');
13 W=12.25/sqrt(V);
14 disp(W, 'de broglie wavelength in angstrom');
15 v=1/(W*10^-10);
16 disp(v, 'the wave number in m');
17 D=asind((W*10^-10)/(2*d));
18 disp(D, 'the Bragg angle in degrees')

```

Scilab code Exa 9.3 AN ELECTRON INTIALLY AT REST

```

1 clc;clear;
2 //Example 9.3
3
4 //given data
5 V=54;//applied voltage in V
6 e=1.602*10^-19;//the charge on electron in C
7 m=9.12*10^-31;//mass of electron in kg
8 h=6.625*10^-34;//Plank's constant
9
10 //calcualtions
11 v=sqrt(2*e*V/m);
12 disp(v, 'velocity of electron in m/s');
13 W=12.25/sqrt(V);
14 disp(W, 'de broglie wavelength in angstrom');
15 u=h/(2*m*W*10^-10);
16 disp(u, 'phase velocity in m/s')

```

Scilab code Exa 9.4 COMPUTE DE BROGLIE WAVELENGTH

```
1 clc; clear;
2 //Example 9.4
3
4 //given data
5 e=1.6*10^-19; //the charge on electron in C
6 m=9.12*10^-31; //mass of electron in kg
7 c=3*10^8; //speed of light in m/s
8 h=6.625*10^-34; //Plank's constant
9
10 //calculations
11 E=m*c^2;
12 mp=1836*m;
13 // (0.5*m*v^2)=E
14 mv=sqrt(E*2*mp);
15 W=h/mv;
16 disp((W/10^-10), 'de broglie wavelength in Angstrom')
```

Scilab code Exa 9.5 WHAT IS DE BROGLIE WAVELENGTH OF NEUTRON

```
1 clc; clear;
2 //Example 9.5
3
4 //given data
5 e=1.6*10^-19; //the charge on electron in C
6 m=1.676*10^-27; //mass of neutron in kg
7 c=3*10^8; //speed of light in m/s
8 h=6.625*10^-34; //Plank's constant
9
10 //calculations
11 E=1; //in eV
12 E=1*e; //in V
13 mv=sqrt(2*E*m);
```

```
14 W=h/mv;  
15 disp((W/10^-10), 'de broglie wavelength in Angstrom')
```

Scilab code Exa 9.6 CALCULATE THE SCATTERED WAVELENGTH

```
1 clc;clear;  
2 //Example 9.6  
3  
4 //calculations  
5 W=0.09; //wavelength in Angstrom  
6 D=54; //scattering angle in degree  
7 h=6.625*10^-34; //Plank's constant  
8 c=3*10^8; //speed of light in m/s  
9 e=1.6*10^-19; //the charge on electron in C  
10  
11 //calculations  
12 dW=0.0243*(1-cosd(D));  
13 W1=W+dW;  
14 disp(W1, 'wavelegth of scattered X-ray in Angstrom');  
15 E=h*c/(W*10^-10);  
16 disp((E/(e*10^6)), 'Energy of incident photon in MeV'  
17 );  
17 E=h*c/(W1*10^-10);  
18 disp((E/(e*10^6)), 'Energy of scattered photon in MeV'  
19 )
```

Scilab code Exa 9.7 COMPUTE ENERGY DIFFERENCE

```
1 clc;clear;  
2 //Example 9.7  
3  
4 //given data  
5 h=6.625*10^-34; //Plank's constant
```

```

6 m=9.12*10^-31; //mass of electron in kg
7
8 //calculations
9 //for (a)
10 nx=1;
11 ny=1;
12 nz=1;
13 L=1;
14 E=h^2*(nx^2+ny^2+nz^2)/(8*m*L^2);
15 disp(E, 'energy in first quantum state in J');
16 //for (b) (nx^2+ny^2+nz^2)=6
17 L=1;
18 E=h^2*6/(8*m*L^2);
19 disp(E, 'energy in second quantum state in J')

```

Scilab code Exa 9.8 CALCULATE THE LOWEST THREE PERMISSIBLE ENERGIES

```

1 clc;clear;
2 //Example 9.8
3
4 //given data
5 h=6.625*10^-34; //Plank's constant
6 m=9.12*10^-31; //mass of electron in kg
7 L=2.5*10^-10;
8 e=1.6*10^-19; //the charge on electron in C
9
10 //calcualtions
11 n=1;
12 E1=n^2*h^2/(8*m*L^2*e);
13 disp(E1, 'E1 in eV');
14 n=2;
15 E2=4*E1;
16 disp(E2, 'E2 in eV');
17 n=3;

```

```
18 E3=9*E1;  
19 disp(E3, 'E3 in eV');
```

Chapter 10

THE ATOMIC NUCLEUS

Scilab code Exa 10.1 COMPARE DENSITIES OF WATER AND NUCLEUS

```
1  clc;clear;
2  //Example 10.1
3
4  //given data
5  R=1.2*10^-15;//radius in A^(1/3)-m *A is mass number
6  mp=1.008;
7  mn=mp;//mass of proton and neutron in a.m.u
8  pi=3.14;//const
9  Dw=1000;//density of water in kg/m^3
10
11 //calculation
12 Vn=4/3*pi*R^3;
13 mp=mp*1.66*10^-27;//conversion in kg
14 mn=mp;
15 m=mn;//m is combined mass in A-kg
16 Dn=m/Vn;
17 R=Dn/Dw;
18 disp(R,'the ratio is')
```

Scilab code Exa 10.2 CALCULATE ENERGY EQUIVALENCE FOR MeV

```
1 clc; clear;  
2 //Example 10.2  
3  
4 //calculations  
5 amu=1.66*10^-27; //1 amu in kg  
6 c=3*10^8; //speed of light in m/s  
7 m=amu;  
8 E=m*c^2;  
9 kWh=1.6*10^-13; //conversion of kWh in J  
10 E=E/kWh;  
11 disp(E, 'energy equivalence in MeV')
```

Scilab code Exa 10.3 CALCULATE ENERGY EQUIVALENCE FOR kWh

```
1 clc; clear;  
2 //Example 10.3  
3  
4 //calculations  
5 gm=10^-3; //1 gram in kg  
6 m=gm;  
7 c=3*10^8; //speed of light in m/s  
8 E=m*c^2;  
9 kWh=36*10^5; //1 kWh in J  
10 EE=E/kWh;  
11 disp(EE, 'energy equivalence in kWh')
```

Scilab code Exa 10.4 CALCULATE BINDING ENERGY

```

1  clc;clear;
2  //Example 10.4
3
4  //given data
5  mn=1.00893; //mass of neutron in a.m.u
6  mp=1.00813; //mass of proton in a.m.u
7  md=2.01473; //mass of deuteron in a.m.u
8  ma=4.00389; //mass of alpha-particle in a.m.u
9
10 //calculations
11 dm=md-(mn+mp);
12 disp((-dm*931), 'binding energy in MeV');
13 dm=ma-2*(mn+mp);
14 disp((-dm*931), 'binding energy in MeV')

```

Scilab code Exa 10.5 CALCULATE BINDING ENERGY OF $^{17}\text{Cl}^{35}$

```

1  clc;clear;
2  //Example 10.5
3
4  //given data
5  m1=1.008665; //mass of  $^0\text{n}^1$  in a.m.u
6  m2=1.007825; //mass of  $^1\text{H}^1$  in a.m.u
7  m3=34.9800; //mass  $^{17}\text{Cl}^{35}$  in a.m.u
8  n=17+18;
9
10 //calculations
11 dm=(17*m2)+(18*m1)-m3;
12 Q=dm*931;
13 disp(Q, 'Binding energy in MeV');
14 BEn=Q/n;
15 disp(BEn, 'Binding energy per nucleon in MeV')

```

Scilab code Exa 10.6 CALCULATE BINDING ENERGY FOR LITHIUM NUCLEUS

```
1  clc; clear;
2  //Example in 10.6
3
4  //given data
5  m1=1.00814; //mass of proton in a.m.u
6  m2=1.00893; //mass of neutron in a.m.u
7  m3=7.01822; //mass of lithium in a.m.u
8
9  //calculations
10 dm=(3*m1)+(4*m2)-m3;
11 Q=dm*931;
12 disp(Q, 'Binding energy in MeV')
```

Chapter 11

NATURAL RADIOACTIVITY

Scilab code Exa 11.1 CALCULATE HOW MUCH RADIUM

```
1  clc;clear;
2  //Example 11.1
3
4  //given data
5  ttg=8378-1898;//total time gap in yrs
6  hf=1620;//half life in yrs
7  n=ttg/hf;//no of half-periods
8  Mo=200;//amt of radium in mg
9
10 //calculations
11 M=Mo*(0.5)^n;
12 disp(M,'radium left in mg')
```

Scilab code Exa 11.2 CALCULATE RADIOACTIVE DISINTEGRATING
CONST

```
1  clc;clear;
2  //Example 11.2
```

```

3
4 //given data
5 hf=30;//half life in days
6 //M is intial conc.
7
8 //calcuations
9 k=0.693/hf;
10 disp(k, 'radioactive disintegration constant in 1/day
    ');
11 //M/4 is left
12 t=-log(1/4)/k;
13 disp(t, 'time taken for (ii) in days');
14 //M/8 is left
15 t=-log(1/8)/k;
16 disp(t, 'time taken for (iii) in days')

```

Scilab code Exa 11.3 COMPUTE DECAY CONST AND HALF LIFE

```

1 clc;clear;
2 //Example 11.3
3
4 //given data
5 No=4750;
6 N=2700;//rate in counts/minute
7 t=5;//time in minutes
8
9 //calculation
10 k=log(No/N)/t;
11 disp(k, 'radioactive disintegration constant in 1/min
    ');
12 hf=0.693/k;
13 disp(hf, 'half life in minutes')

```

Scilab code Exa 11.4 ESTIMATE HALF LIFE OF PLUTONIUM

```
1  clc;clear;
2  //Example 13.4
3
4  //given data
5  m=4.00387;//mass of alpha particle in a.m.u
6  M=10^-6;//mass of Pu-239 in kg
7
8  //calculations
9  m=m*1.66*10^-24;//conversion in gm
10 Mo=2300*m;
11 k=(Mo/1)/M;
12 hf=0.693/k;
13 hf=hf/(365*24*3600);//conversion in yrs
14 disp(hf,'half life in yrs')
```

Scilab code Exa 11.5 CALCULATE ACTIVITY

```
1  clc;clear;
2  //Example 11.5
3
4  //given data
5  hf=2.48*10^5;//half life in yrs
6  k=8.88*10^-14//decay const in 1/s
7  Mo=4;//intial mass in mg
8  Na=6.02*10^23;//Avgraodo no. in 1/gm mole
9
10 //calculations
11 kt=0.693/hf*62000;
12 M=Mo*(exp(-kt));
13 disp(M,'mass remain unchanged in mg');
14 N=M*10^-3*Na/234;
15 A=k*N;
16 disp(A,'Activity in disintegrations/second')
```

Scilab code Exa 11.6 HALF LIFE OF RADIUM IS 1620 YRS

```
1  clc; clear;
2  //Example 11.6
3
4  //given data
5  hf=1620; //half life in yrs
6  Mo=1/100; //mass in gm
7
8  //calculations
9  k=0.693/hf;
10 M=(1-Mo);
11 t=log(1/M)/k;
12 disp(t, 'time reqd for (i) in yrs');
13 M=Mo;
14 t=log(1/M)/k;
15 disp(t, 'time reqd for (ii) in yrs')
```

Chapter 12

ARTIFICIAL RADIOACTIVITY

Scilab code Exa 12.1 ESTIMATE ITS AGE

```
1  clc; clear;  
2  //Example 12.1  
3  
4  //given data  
5  r=0.5; //ratio of mass of Pb206 and mass of U238  
6  t=4.5*10^9; //half life in years  
7  
8  //calculation  
9  T=(log(1+r))*(t/0.693);  
10 disp(T, 'age in years')
```

Chapter 13

NUCLEAR REACTIONS

Scilab code Exa 13.1 CALCULATE THE ENERGY AVAILABLE

```
1  clc; clear;
2  //Example 13.1
3
4  //given data
5  m1=7.0183; //mass of 3Li7 in a.m.u
6  m2=4.0040; //mass of 2He4 in a.m.u
7  m3=1.0082; //mass of 1H1 in a.m.u
8  Na=6.02*10^26; //Avgrado no. in 1/kg mole
9  //rxn = 3Li7 + 1H1 = 2He4 + 2He4
10
11 //calculations
12 dm=m1+m3-(2*m2);
13 E=dm*931;
14 n=0.1*Na/7; //no of atoms in 100 gm of lithium
15 TE=n*E;
16 disp(TE, 'Total energy available in MeV')
```

Scilab code Exa 13.2 CALCULATE THE ENERGY RELEASED

```

1  clc;clear;
2  //Example 13.2
3
4  //given data
5  m1=6.015126;//mass of  ${}^3\text{Li}_7$  in a.m.u
6  m2=4.002604;//mass of  ${}^2\text{He}_4$  in a.m.u
7  m3=1.00865;//mass of  ${}^1_0\text{n}$  in a.m.u
8  m4=3.016049;//mass of  ${}^1_1\text{H}$  in a.m.u
9  //rxn =  ${}^3\text{Li}_7 + {}^1_0\text{n} = {}^2\text{He}_4 + {}^1_1\text{H} + \text{Q}$ 
10
11 //calculations
12 dm=m1+m3-(m2+m4);
13 Q=dm*931;
14 disp(Q,'energy released in MeV')

```

Scilab code Exa 13.3 WHAT IS THE Q VALUE OF THE REACTION

```

1  clc;clear;
2  //Example 13.3
3
4  //given data
5  m1=14.007515;//mass of  ${}^7\text{N}_{14}$  in a.m.u
6  m2=4.003837;//mass of  ${}^2\text{He}_4$  in a.m.u
7  m3=17.004533;//mass of  ${}^8\text{O}_{17}$  in a.m.u
8  m4=1.008142;//mass of  ${}^1_1\text{H}$  in a.m.u
9  //rxn =  ${}^7\text{N}_{14} + {}^2\text{He}_4 = {}^8\text{O}_{17} + {}^1_1\text{H}$ 
10
11 //calculations
12 dm=m3+m4-(m1+m2);
13 Q=dm*931;
14 disp(Q,'energy released in MeV')

```

Scilab code Exa 13.4 FIND THE MASS

```

1  clc;clear;
2  //Example 13.3
3
4  //given data
5  m1=14.007520;//mass of  ${}^7\text{N}14$  in a.m.u
6  m2=1.008986;//mass of  ${}^1_0\text{n}1$  in a.m.u
7  //m3=mass of  ${}^6\text{C}14$  in a.m.u
8  m4=1.008145;//mass of  ${}^1_1\text{H}1$  in a.m.u
9  //rxn =  ${}^7\text{N}14 + {}^1_0\text{n}1 = {}^6\text{C}14 + {}^1_1\text{H}1 + 0.55 \text{ MeV}$ 
10
11 //calculations
12 Q=0.55;
13 dm=Q/931;
14 m3=dm+m1+m2-m4;
15 disp(m3,'mass of  ${}^6\text{C}14$  in a.m.u')

```

Scilab code Exa 13.6 EXPLAIN MASS DEFECT

```

1  clc;clear;
2  //Example 13.6
3
4  //given data
5  m0=11.01280;//mass  ${}^5\text{B}11$  in a.m.u
6  m1=4.00387;//mass of alpha particle in a.m.u
7  m2=14.00752;//mass of  ${}^7\text{N}14$  in a.m.u
8  //m3=mass of neutron
9  E1=5.250;//energy of alpha particle in MeV
10 E2=2.139;//energy of  ${}^7\text{N}14$  in MeV
11 E3=3.260;//energy of  ${}^1_0\text{n}1$  in MeV
12
13 //calculations
14 m3=(m0*931)+((m1*931)+E1)-((m2*931)+E2)-E3;
15 disp((m3/931),'mass of neutron in a.m.u')

```

Chapter 14

NUCLEUR FISSION AND FUSION

Scilab code Exa 14.1 COMPUTE FISSION ENERGY

```
1  clc; clear;  
2  //Example 14.1  
3  
4  //given data  
5  E1=7.8; //avg. B.E per nucleon in MeV  
6  E2=8.6; //for fission fragments in MeV  
7  
8  //calculations  
9  FER=(234*E2) - (236*E1);  
10 disp(FER, 'Fission energy released in MeV')
```

Scilab code Exa 14.2 FIND ELEMENTARY PARTICLES RELEASED
IN BINARY FISSION OF $^{92}\text{U}^{235}$

```
1  clc; clear;  
2  //Example 14.2
```

```

3
4 //given data
5 m1=235.044; //mass of 92U235 in a.m.u
6 m2=97.905; //mass of 42Mo98 in a.m.u
7 m3=135.917; //mass of 54Xe136 in a.m.u
8 //rxn = 0n1 + 92U235 = 42Mo98 + 54Xe136 + 4 -1e0 + 2
      0n1
9
10 //calculation
11 LHSm=1.009+m1;
12 RHSm=m2+m3+(4*0.00055)+(2*1.009);
13 dm=LHSm-RHSm;
14 disp(dm, 'mass defect in a.m.u');
15 E=dm*931;
16 disp(E, 'energy released in MeV')

```

Scilab code Exa 14.3 HOW MUCH HYDROGEN MUST BE CONVERTED INTO HELIUM

```

1 clc;clear;
2 //Example 14.3
3
4 //given data
5 m1=1.00813; //mass of 1H1 in a.m.u
6 m2=4.00386; //mass of 2He4 in a.m.u
7 SC=1.35; //solar constant in kW/m^2
8 d=1.5*10^11; //dist b/w earth and sun in m
9 e=1.6*10^-19; //the charge on electron in C
10 Na=6.02*10^26; //Avgrado no. in 1/kg mole
11 pi=3.14; //const
12 //rxn = 4 1H1 = 2He4 + 2 1e0
13
14 //calculations
15 dm=(4*m1)-m2
16 E=dm*931; //energy produced in MeV

```

```

17 EP=E/4; //energy produced per atom
18 EP=EP*10^6*e; //conversion in J
19 EPkg=EP*Na; //energy produced by 1 kg of hydrogen
20 SC=SC*1000; //conversion in J/s-m^2
21 SA=4*pi*d^2; //surface area of sphere
22 ER=SC*SA; //energy recieved per second
23 m=ER/EPkg;
24 disp((m/10^3), 'mass of hydrogen consumed in tonnes/
      second ')

```

Scilab code Exa 14.4 CALCULATE THE ENERGY LIBERATED

```

1  clc;clear;
2  //Example 14.4
3
4  //given data
5  m1=2.01478; //mass of 1H2 in a.m.u
6  m2=4.00388; //mass of 2He4 in a.m.u
7  //rxn 1H2 + 1H2 = 2He4 + Q
8
9  //calculations
10 Q=2*m1-m2;
11 Q=Q*931; //conversion in MeV
12 disp(Q, 'energy liberated in MeV')

```

Chapter 15

NUCLEUR ENERGY SOURCES

Scilab code Exa 15.1 CALCULATE THE MAXIMUM FRACTION OF
THE KE

```
1  clc; clear;
2  //Example 15.1
3
4  //given data
5  ma=1;
6  Ma=2;
7  mb=1;
8  Mb=12;
9  mc=1;
10 Mc=238; //m is mass of neutron and M is mass of other
           nucleus
11
12 //calculation
13 n=(4*ma*Ma/(ma+Ma)^2)*100;
14 disp(n, 'Maximum fraction of KE lost by a neutron for
           (a) ');
15 n=(4*mb*Mb/(mb+Mb)^2)*100;
16 disp(n, 'Maximum fraction of KE lost by a neutron for
```

```

    (a) ');
17 n=(4*mc*Mc/(mc+Mc)^2)*100;
18 disp(n, 'Maximum fraction of KE lost by a neutron for
    (a) ')

```

Scilab code Exa 15.2 CALCULATE THE FISSION RATE

```

1  clc;clear;
2  //Example 15.2
3
4  //given data
5  E=200; //energy released per fission in MeV
6  e=1.6*10^-19; //the charge on electron in C
7  Na=6.02*10^26; //Avgrado no. in 1/kg mole
8
9  //calculations
10 CE=E*e*10^6; //conversion in J
11 RF=1/CE;
12 disp(RF, 'fission rate of one watt in fissions/second
    ');
13 Ekg=CE*Na/235;
14 disp(Ekg, 'Energy realeased in complete fission of 1
    kg in J')

```

Scilab code Exa 15.3 HOW MANY KG OF U 235

```

1  clc;clear;
2  //Example 15.3
3
4  //given data
5  R=3*10^7; //rate of energy development in J s
6  E=200; //energy released per fission in MeV
7  e=1.6*10^-19; //the charge on electron in C

```



```

8 t=1000; //time is hours
9 Ekg=8.2*10^13; //energy released per kg of U-235
10
11 //calculation
12 CE=E*e*10^6; //conversion in J
13 n=R/CE;
14 disp(n, 'no of atoms undergo fission/second ');
15 TE=R*t*3600; //energy produced in 1000 hours
16 MC=TE/Ekg;
17 disp(MC, 'mass consumed in kg')

```

Scilab code Exa 15.4 HOW MUCH U 235 WOULD BE CONSUMED IN THE RUN

```

1 clc;clear;
2 //Example 15.4
3
4 //given data
5 EPF=180; //Energy consumed per disintegration in MeV
6 E=1200; //average power in kW
7 t=10; //time in hours
8 Na=6.02*10^26; //Avgrado no. in 1/kg mole
9 e=1.6*10^-19; //the charge on electron in C
10
11 //calculation
12 TE=E*t; //energy consumed in kWh
13 TE=TE*36*10^5; //conversion in J
14 EE=TE/0.2; //efficient energy
15 CE=EPF*e*10^6; //conversion in J
16 n=EE/CE;
17 m=235*n/Na*1000;
18 disp(m, 'mass consumed in gram')

```

Scilab code Exa 15.5 NUCLEUR REACTOR PRODUCES 200MW

```
1  clc;clear;
2  //Example 15.5
3
4  //given data
5  OE=200;//o/p power in MW
6  E=200;//energy released per fission in MeV
7  WF=3.1*10^10;//fission rate in fissions/second
8  Na=6.02*10^26;//Avgraodo no. in 1/kg mole
9
10 //calculations
11 IE=OE/0.3*10^6;//reactor input in W
12 TFR=WF*IE;
13 n=TFR*24*3600;//no. of U-235 for one day
14 m=235*n/Na;
15 disp((m*100/0.7),'amt of natural uranium conumed/day
      in kg')
```

Scilab code Exa 15.6 A CITY REQUIRES 100 MW

```
1  clc;clear;
2  //Example 15.6
3
4  //given data
5  AE=100;//electrical power in MW
6  E=200;//energy released per fission in MeV
7  e=1.6*10^-19;//the charge on electron in C
8  Na=6.02*10^26;//Avgraodo no. in 1/kg mole
9
10 //calculations
11 TE=AE*10^6*24*3600;//energy consumed in city in one
      day in J
12 EE=TE/0.2;
13 CE=E*e*10^6;//conversion in J
```

```

14 n=EE/CE;
15 m=235*n/Na;
16 disp(m, 'amt of fuel required in kg')

```

Scilab code Exa 15.7 BOMBAY REQUIRES 300 MWh

```

1  clc;clear;
2  //Example 15.7
3
4  //given data
5  OE=3000;//output power in MWh
6  E=200;//energy released per fission in MeV
7  e=1.6*10^-19;//the charge on electron in C
8  Na=6.02*10^26;//Avgrado no. in 1/kg mole
9
10 //calculations
11 IE=OE/0.2;
12 TE=IE*36*10^8;//conversion in J
13 CE=E*e*10^6;//conversion in J
14 n=TE/CE;
15 m=235*n/Na;
16 disp(m, 'daily fuel requirement in kg')

```

Scilab code Exa 15.8 THE MOTOR OF AN ATOMIC ICE BREAKER

```

1  clc;clear;
2  //Example 15.8
3
4  //given data
5  OP=32824;//o/p power in kW
6  E=200;//energy released per fission in MeV
7  Ekg=8.2*10^13;//energy released per kg of U-235
8

```

```
9 //calculations
10 DOP=OP*1000*24*3600;//daily o/p power in J
11 IP=DOP/0.2;
12 DFC=IP/Ekg;//daily fuel consumption
13 disp(DFC,'daily fuel consumption in kg');
14 DI=DOP/(0.8*4186);//daily input at 80% efficiency
15 Crqd=DI/(7*103);
16 disp(Crqd,'Coal reqd/day in tonnes')
```

Chapter 16

PARTICLE ACCELERATORS

Scilab code Exa 16.1 WHAT MUST BE THE FLUX DENSITY

```
1  clc;clear;
2  //Example 16.1
3
4  //given data
5  fo=9*10^6;//frequency in Hz
6  m=6.643*10^-27;//mass in kg
7  pi=3.14;//constant
8  e=1.6*10^-19;//the charge on electron in C
9
10 //calculations
11 Q=2*e;
12 B=fo*2*pi*m/Q;
13 disp(B,'magnetic flux density in Wb/m^2')
```

Scilab code Exa 16.2 WHAT IS FREQUENCY OF ALTERNATING POTENTIAL

```
1  clc;clear;
```

```

2 //Example 16.2
3
4 //given data
5 B=0.7;//magnetic flux intensity in Wb/m^2
6 m=3.34*10^-27;//mass in Kg
7 e=1.6*10^-19;//the charge on electron in C
8 pi=3.14;//const
9
10 //calculations
11 Q=e;
12 fo=B*Q/(2*pi*m*10^6);
13 disp(fo,'The cyclotron frequency in MHz ')

```

Scilab code Exa 16.3 A CYCLOTRON OF DEES OF RADIUS 2 METRES

```

1 clc;clear;
2 //Example 16.3
3
4 //given data
5 B=0.75;//magnetic flux intensity in Wb/m^2
6 m1=1.67*10^-27;//mass in Kg
7 m2=3.31*10^-27;//mass in Kg
8 e=1.6*10^-19;//the charge on electron in C
9 Rm=2;//radius in m
10
11 //calculations
12 Q=e;
13 Emax=3.12*10^12*B^2*Q^2*Rm^2/m1;
14 disp(Emax,'Maximum energies in Mev for proton');
15 Emax=3.12*10^12*B^2*Q^2*Rm^2/m2;
16 disp(Emax,'Maximum energies in Mev for deuteron')

```

Scilab code Exa 16.4 CALCULATE THE RATIO

```
1  clc; clear;
2  //Example 16.4
3
4  //given data
5  mo=9.1*10^-31; //mass of electron in kg
6  m=1.67*10^-27; //mass of proton in kg
7  c=3*10^8; //speed of light in m/s
8  E=1; //given energy in MeV
9
10 //calculations
11 Eo=mo*c^2/(1.6*10^-13);
12 mbymo=1+(E/Eo);
13 disp(mbymo, 'Ratio for electron');
14 Eo=m*c^2/(1.6*10^-13);
15 mbymo=1+(E/Eo);
16 disp(mbymo, 'Ratio for proton')
```

Scilab code Exa 16.5 IN A CERTAIN BETATRON

```
1  clc; clear;
2  //Example 16.5
3
4  //given data
5  B=0.5; //magnetic field in Wb/m^2
6  d=1.5; //diameter in m
7  f=59; //frequency in Hz
8  e=1.6*10^-19; //the charge on electron in C
9  c=3*10^8; //speed of light in m/s
10 pi=3.14; //const
11
12 //calculations
13 R=d/2;
14 N=c/(4*(2*pi*50)*R);
```

```

15 E=B*e*R*c/(1.6*10^-13);
16 disp(E,'final energy in MeV');
17 AE=E/N*10^6;
18 disp(AE,'average energy in eV')

```

Scilab code Exa 16.6 CALCULATE MASS AND VELOCITY OF ELCTRONS

```

1  clc;clear;
2  //Example 16.6
3
4  //given data
5  E=0.51;//kinetic energy in MeV
6  R=0.15;//radius in m
7  e=1.6*10^-19;//the charge on electron in C
8  mo=9.12*10^-31;//mass of electron in kg
9  c=3*10^8;//speed of light in m/s
10
11 //calculation
12 Eo=E;
13 m=mo*(1+(E/Eo));
14 b=sqrt(1-(mo/m)^2);
15 v=b*c;
16 B=mo*v/(e*R);
17 disp(B,'magnetic field intensity')

```

Scilab code Exa 16.7 DETERMINE THE FREQUENCY OF GENERATOR

```

1  clc;clear;
2  //Example 16.7
3
4  //given data
5  E=4;//applied voltage in MeV

```



```

6 m=3.334*10^-27; //mass of deuteron in kg
7 R=0.75; //radius in m
8 pi=3.14; //const
9 e=1.6*10^-19; //the charge on electron in C
10
11 //calcuations
12 E=4*10^6*e;
13 fo=sqrt(E/(2*m))/(pi*R);
14 disp(fo, 'frequency of generator in Hz')

```

Scilab code Exa 16.8 WHAT WOULD BE THE ENERGY OF ELECTRON

```

1 clc;clear;
2 //Example 16.8
3
4 //given data
5 roi=15; //rate of increase in Wb/s
6 tr=10^6; //total revolutions
7
8 //calcualtion
9 IE=roi*10^-6; //increased energy in MeV
10 FE=IE*tr;
11 disp(FE, 'Fianl Energy in MeV')

```

Scilab code Exa 16.9 FIND THE MAX ENERGY AND CORRESPONDING WAVELENGTH

```

1 clc;clear;
2 //Example 16.9
3
4 //given data
5 R=0.1; //radius in m

```

```

6 pi=3.14; //const
7 h=6.625*10^-34; //Plank's constant
8 c=3*10^8; //speed of light in m/s
9 roi=15; //rate of increase in Wb/s
10 t=4*10^-4; //period of accerleartion in s
11 e=1.6*10^-19; //the charge on electron in C
12
13 //calculations
14 N=c*t/(2*pi*R);
15 IE=roi; //incresed energy in eV
16 ME=N*IE*10^-6;
17 disp(ME, 'Maximum energy in MeV');
18 ME=ME*10^6*e; //conversion in V
19 p=ME/c;
20 Y=h/p;
21 disp(Y, 'corresponding wavelength of X-rays in m')

```
