

Scilab Textbook Companion for
Material Science
by S. L. Kakani and A. Kakani¹

Created by
Sumit Joshi
B. tech
Electrical Engineering
Uttarakhand Technical University
College Teacher
Laxmi Datt Pant
Cross-Checked by
Mukul Kulkarni and Lavitha Pereira

February 16, 2014

¹Funded by a grant from the National Mission on Education through ICT,
<http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab
codes written in it can be downloaded from the "Textbook Companion Project"
section at the website <http://scilab.in>

Book Description

Title: Material Science

Author: S. L. Kakani and A. Kakani

Publisher: New Age International Publishers, New Delhi

Edition: 1

Year: 2005

ISBN: 81-224-1528-8

Scilab numbering policy used in this document and the relation to the above book.

Exa Example (Solved example)

Eqn Equation (Particular equation of the above book)

AP Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

Contents

List of Scilab Codes	5
2 Atomic structure and electronic configuration	9
3 Crystal Geometry Structure and Defects	17
4 Bonds in solid	29
5 Electron Theory of Metals	33
6 Photoelectric Effect	39
7 Diffusion in Solids	44
8 Mechanical Properties of Materials and Mechanical Tests	48
9 Alloys Systems Phase Diagrams and Phase Transformations	51
10 Heat Treatment	53
11 Deformation of Materials	54
12 Oxidation and Corrosion	57
13 Thermal and Optical Properties of Materials	59
14 Electrical and Magnetic Properties of Materials	60
15 Semiconductors	66

16 Superconductivity and Superconducting Materials	69
18 Composites	70

List of Scilab Codes

Exa 2.1	Calculate the distance of the closest approach alpha particles from the copper nucleus	9
Exa 2.2	Calculate the radius and frequency of an electron in the bohr first orbit in hydrogen atom	10
Exa 2.3	Determine the radius of the second bohr orbit in a singly ionized helium atom	10
Exa 2.4	Calculate the unit cell dimensions and atomic diameter	11
Exa 2.5	Calculate the revolutions per second of an electron in the bohr orbit of hydrogen atom	11
Exa 2.6	Determine the orbital frequency of an electron in the first bohr orbit in a hydrogen atom	12
Exa 2.7	Calculate the values of kinetic energy potential energy and total energy of an electron	13
Exa 2.8	Calculate the velocity of an electron in hydrogen atom in bohr first orbit	13
Exa 2.9	Calculate the principal quantum number and wavelengths of radiation in both energies	14
Exa 2.13	Assuming that the weight of one proton is equal to that of one neutron find the weight of copper atom and weight of one proton	15
Exa 2.15	find the percentage of Si in Copper silicide	15
Exa 3.10	Find the angle Between normals to the planes	17
Exa 3.11	Determine the packing efficiency and density of sodium chloride	17
Exa 3.12	Calculate the unit cell dimensions and atomic diameter	18
Exa 3.13	Calculate the interplaner distance	19
Exa 3.14	Find the number of atoms per mm ²	20
Exa 3.15	Determine the planer density of Ni	20

Exa 3.16	Calculate the planar atomic densities of planes	21
Exa 3.17	Calculate the linear atomic densities of planes	21
Exa 3.18	Find lattice constant	22
Exa 3.19	Find the numbers of atoms per unit cell	23
Exa 3.20	Calculate the line energy of disslocation in bcc iron . .	23
Exa 3.22	Calculate the number of vacancies	23
Exa 3.23	Calculate the surface energy of copper	24
Exa 3.24	Calculate the equilibrium concentration of vacancies in aluminium	25
Exa 3.25	Determine the interplanar spacing	25
Exa 3.26	Calculate the size of unit cell	26
Exa 3.27	Calculate the Bragg angle	26
Exa 3.28	Calculate the bragg reflection index	27
Exa 3.29	Calculate the angle for 3rd order reflection	27
Exa 3.30	Obtain the interplanar spacing and miller indices of the reflection plane	28
Exa 3.31	Determine interatomic spacing	28
Exa 4.1	Find A and B and the force needed to dissociate this molecule into atoms and the interatomic distance at which the dissociation occurs	29
Exa 4.2	Find the repulsive exponent n	30
Exa 4.3	Find the radius of Cl ion	30
Exa 4.4	Calculate the force of attraction between ions	31
Exa 4.5	How much net energy is spent in the process	31
Exa 4.6	Estimate the fraction of hydrogen bonds that are broken when ice melts	32
Exa 5.1	Evaluate the temperature	33
Exa 5.2	Find the drift velocity of carriers	34
Exa 5.3	Find the conductivity of copper at 300K	34
Exa 5.4	Find the mobility of condution electron	35
Exa 5.5	Find the mobility of condution electron and drift velocity	35
Exa 5.6	Find the number of free electrons and also calculate mobility of electrons	36
Exa 5.7	Calculate the maximum velocity of an electron in a metal and mobility of electrons	36
Exa 5.8	Calculate the velocity of an electrons at fermi level . .	37
Exa 5.9	Estimate the mean path of free electrons in pure copper	37
Exa 5.10	Find the conductivity of copper	38

Exa 6.1	Find the kinetic energy of electrons ejected from the surface	39
Exa 6.2	Calculate the threshold frequency and the corresponding wavelength	40
Exa 6.3	Calculate the threshold frequency and the work function of metal	40
Exa 6.4	Calculate the photons emitted by lamp per second	41
Exa 6.5	Determine the region of electrons spectrum	41
Exa 6.6	Calculate the photons emitted by radio receiver	42
Exa 6.7	Calculate the wavelength of light which can just eject electron from tungsten and from barium	42
Exa 7.1	find the time required for carburization	44
Exa 7.2	Calculate time required to get a boron content of 1023 atoms per m ³ at a depth of 2 micro meter	44
Exa 7.3	find the constant of the equation and activation energy and determine the diffusion coefficient	45
Exa 7.4	Determine the approximate time that will produce same diffussion at 500 C	46
Exa 8.1	Determine the fracture strength	48
Exa 8.2	Compute the true stress and strain with engineering stress and strain	48
Exa 8.3	Determine the fracture strength	49
Exa 8.4	Find the resultant elongation	50
Exa 8.5	Compute the strain hardening exponent	50
Exa 9.1	Find the percentage of proeutectoid ferrite	51
Exa 9.2	Determine the degrees of freedom of a system of two components	51
Exa 9.3	Find the minimum number of components in the system	52
Exa 10.1	Determine the grain diameter of an ASTM Number 8	53
Exa 11.1	Determine the value of critical resolved shear stress	54
Exa 11.3	Find the yield stress	55
Exa 11.4	Find the yield stress	55
Exa 12.1	Find the distance at which magnesium anode capable of giving 2MA	57
Exa 12.2	Calculate the quality of magnesium required per square meter of the hull surface for a design life of 10 years	57

Exa 13.1	Determine the maximum temperature to which the rod may be heated without exceeding a compressive stress of 172 MPa	59
Exa 14.1	Calculate the resistance of an aluminium wire	60
Exa 14.2	Find the resistivity of a copper alloy containing 1 atomic percent nickel and 3 atomic percent silver	60
Exa 14.3	Find the resistivity due to impurity scattering per percent of Ni in the Cu lattice	61
Exa 14.4	Calculate the relative dielectric constant of a barium titanate crystal	62
Exa 14.5	Calculate the polarization	62
Exa 14.6	Find net magnetic moment per iron atom in the crystal	63
Exa 14.7	Calculate the saturation magnetization and the saturation flux density	64
Exa 14.8	Calculate the saturation magnetization	64
Exa 14.9	Calculate eddy current loss at the normal voltage and frequency	65
Exa 15.1	Find the conductivity and resistivity of a pure silicon crystal	66
Exa 15.2	Find the resistivity at room temperature	67
Exa 15.3	Find the resistance of an intrinsic germanium rod . . .	67
Exa 15.4	Obtain density relation in P type material	68
Exa 16.1	Calculate the energy gap in electron volts and Calculate the wavelength of a photon	69
Exa 18.1	Calculate the modulus of elasticity	70
Exa 18.2	Calculate the elastic modulus when the stress is applied perpendicular to the direction of the fibre alignment .	70

Chapter 2

Atomic structure and electronic configuration

Scilab code Exa 2.1 Calculate the distance of the closest approach alpha particles from the copper nucleus

```
1 //Exam : 2 . 1
2 clc;
3 clear;
4 close;
5 Eg_k=5; //kinetic energy of alpha particles (in MeV)
6 Eg_K=5*(10^6)*1.6*(10^-19); //kinetic energy of alpha
    particles (in J)
7 mv2=2*Eg_K;
8 pi=22/7;
9 phi=180; //firing angle
10 Z=29; //Atomic number
11 e=1.6*(10^-19); //electron charge (in C)
12 Eo=8.85*10^-12; //permittivity of free space
13 d=(Z*e^2/(2*pi*Eo*mv2))*(1+cscd(90))/;
14 disp(d, 'distance of the closest approach alpha
    particles from the copper nucleus (in meter)=');
```

Scilab code Exa 2.2 Calculate the radius and frequency of an electron in the bohr first orbit in hydrogen atom

```
1 //Exam:2.2
2 clc;
3 clear;
4 close;
5 e=1.6*10^(-19); //electron charge (in C)
6 m=9.1*10^(-31); //mass of electron (in Kg)
7 E_o=8.854*10^(-12); //permittivity of free space
8 h=6.625*10^(-34); //Planck constant
9 n=1; //Orbit number
10 Z=1; //atomic number
11 pi=22/7;
12 r_1=(E_o*h^2)/(pi*m*e^2); //first orbit radius of
    hydrogen atom
13 disp(r_1,'first orbit radius of hydrogen atom (in m)=
    ');
14 Freq=m*(Z^2)*(e^4)/(4*(E_o^2)*(n^3)*h^3); //
15 disp(Freq,'Orbital frequency of electron (in Hz)=');
```

Scilab code Exa 2.3 Determine the radius of the second bohr orbit in a singly ionized helium atom

```
1 //Exam:2.3
2 clc;
3 clear;
4 close;
5 Z_1=1; //atomic number for hydrogen
6 n_1=1; //first orbit
7 r_1=0.529; //radius of first orbit of electron for
    hydrogen
```

```
8 Z_2=2; //atomic number for helium
9 n_2=2; //second orbit
10 k=r_1*Z_1/n_1;
11 r_2=k*((n_2)^2)/Z_2; //radius of first orbit of
    electron for helium
12 disp(r_2, 'radius of the second bohr orbit in a
    singly ionized helium atom(in A)=');
```

Scilab code Exa 2.4 Calculate the unit cell dimensions and atomic diameter

```
1 //Exam:2.4
2 clc;
3 clear;
4 close;
5 n_1=1; //first orbit
6 n_2=2; //second orbit
7 n_3=3; //third orbit
8 //E_1=-13.6*(Z^2)/(1^2);
9 //E_2=-13.6*(Z^2)/(2^2);
10 //E_3=-13.6*(Z^2)/(3^2);
11 //E_3-E_1=-13.6*(Z^2)*(-8/9);
12 //E_2-E_1=-13.6*(Z^2)*(-3/4);
13 E_1=-13.6/(1^2); //energy of electron in the first
    bohr orbit of an atom
14 E_2=-13.6/(2^2); //energy of electron in the second
    bohr orbit of an atom
15 E_3=-13.6/(3^2); //energy of electron in the third
    bohr orbit of an atom
16 disp((E_3-E_1)/(E_2-E_1), 'ratio of energy released =
    ');
```

Scilab code Exa 2.5 Calculate the revolutions per second of an electron in the bohr orbit of hydrogen atom

```
1 //Exam:2.5
2 clc;
3 clear;
4 close;
5 m=9.1*10^(-31); //electron mass (in Kg)
6 Z=1; //atomic number
7 e=1.6*10^(-19); //electron charge(in C)
8 E_o=8.25*10^(-12); //permittivity of free space
9 n=1; //first bohr orbit
10 h=6.63*10^(-34); //planck constant
11 R_ps=m*(e^4)/(4*(E_o^2)*(h^3)); //number of
    revolutions per second
12 disp(R_ps, 'revolutions per second of an electron in
    the bohr orbit of hydrogen atom=');
```

Scilab code Exa 2.6 Determine the orbital frequency of an electron in the first bohr orbit in a hydrogen atom

```
1 //Exam:2.6
2 clc;
3 clear;
4 close;
5 n=1; //first bohr orbit
6 Z=1; //atomic number
7 m=9.1*10^(-31); //electron mass in Kg.
8 e=1.6*10^(-19); //electron charge(in C)
9 E_o=8.85*10^(-12); //permittivity of free space
10 h=6.63*10^(-34); //planck constant
11 v_n=m*(Z^2)*(e^4)/(4*(E_o^2)*(h^3)*(n^3)); //orbital
    frequency of an electron in the first bohr orbit
    in a hydrogen atom
12 disp(v_n, 'orbital frequency of an electron in the
```

```
first bohr orbit in a hydrogen atom( in Hz)=') ;
```

Scilab code Exa 2.7 Calculate the values of kinetic energy potential energy and total energy of an electron

```
1 //Exam:2.7
2 clc;
3 clear;
4 close;
5 m=9.11*10^-31; //mass of electron (in Kg)
6 Z=1; //atomic number
7 n=1; //first bohr orbit
8 E_o=8.854*10^-12; //permittivity of free space
9 h=6.62*10^-34; //planck constant
10 e=1.6*10^-19; //electron charge(in C)
11 E_k=(m*(Z^2)*(e^4))/(8*(E_o^2)*(n^2)*(h^2)); //
    Kinetic energy(in joule)
12 E=E_k/e; //Kinetic energy(in eV)
13 E_t=-13.6*(Z^2/n^2); //Total Energy(in eV)
14 E_p=E_t-E; //Potential energy(in eV)
15 disp(E_t, 'Total energy (in eV)=');
16 disp(E, 'kinetic energy (in eV)=');
17 disp(E_p, 'potential energy (in eV)=');
```

Scilab code Exa 2.8 Calculate the velocity of an electron in hydrogen atom in bohr first orbit

```
1 //Exam:2.8
2 clc;
3 clear;
4 close;
5 h=6.626*10^-34; //planck constant
6 E_o=8.825*10^-12; //permittivity of free space
```

```

7 e=1.6*10^-19; //electron charge(in C)
8 n=1; //first bohr orbit
9 Z=1; //atomic number
10 v=Z*(e^2)/(2*E_o*n*h); //velocity of electron in
    hydrogen atom in bohr first orbit
11 disp(v, 'velocity of electron in hydrogen atom in
    bohr first orbit(in meter/sec)=');

```

Scilab code Exa 2.9 Calculate the principal quantum number and wavelengths of radiation in both energies

```

1 //Exam:2.9
2 clc;
3 clear;
4 close;
5 n_1=1; //electron excited from ground state
6 h=6.62*10^-34; //Planck constant
7 c=3*10^8; //speed of light
8 E_o=8.825*10^-12; //permittivity of free space
9 e=1.6*10^-19; //electron charge(in C)
10 m=9.11*10^-31; //mass of electron(in Kg)
11 E_1=10.2; //energy excites the hydrogen from ground
    level(in eV)
12 K=m*e^4/(8*(E_o^2)*(h^2))//in joule
13 K_e=K/e;//in eV
14 //E_1=K_e*((1/n_1^2)-(1/n^2))
15 //1/(n^2)=1/(n_1^2)-E_1/K_e
16 //n^2=1/(1/(n_1^2)-E_1/K_e)
17 n=(1/(1/(n_1^2)-E_1/K_e))^(1/2); //principal quntum
    number when 10.2 eV energy excites electron
18 disp(ceil(n), 'principal quntum number when 10.2 eV
    energy excites electron=');
19 W_1=h*c/(E_1*e)*10^10; //wavelength of radiation when
    10.2 eV energy excites electron
20 disp(W_1, 'wavelength of radiation when 10.2 eV

```

```

        energy excites electron (in A)=')
21 E_2=12.09; //energy excites the hydrogen from ground
    level (in eV)
22 n_2=(1/(1/(n_1^2)-E_2/K_e))^(1/2); //principal quantum
    number when 12.09 eV energy excites electron
23 W_2=h*c/(E_2*e)*10^10; //wavelength of radiation when
    12.09 eV energy excites electron
24 disp(ceil(n_2), 'principal quantum number when 12.09
    eV energy excites electron=')
25 disp(W_2, 'wavelength of radiation when 12.09 eV
    energy excites electron (in A)=')

```

Scilab code Exa 2.13 Assuming that the weight of one proton is equal to that of one neutron find the weight of copper atom and weight of one proton

```

1 //Exam:2.13
2 clc;
3 clear;
4 close;
5 At_w=63.54; //atomic weight of copper
6 N=6.023*10^23; //avogadro's number
7 W_a=At_w/N; //weight of one atom(in gm)
8 W_p=W_a/63; //weight of one proton(in gm)
9 disp(W_a, 'weight of one atom(in gm)=');
10 disp(W_p, 'weight of one proton(in gm)=');

```

Scilab code Exa 2.15 find the percentage of Si in Copper silicide

```

1 //Exam:2.15
2 clc;
3 clear;
4 close;
5 Atw_Cu=63.54; //atomic weight of copper

```

```
6 Atw_Si=28.09; //atomic weight of silicon
7 // 5 atoms of copper working in Cu_5_Si
8 Tw_Cu=5*Atw_Cu; //total weight of copper used in
copper silicide
9 Tw_Si=Atw_Si; //total weight of silicon used in
copper silicide
10 Percentage=(Tw_Si/(Tw_Cu+Tw_Si))*100; //percentage of
Si in Copper silicide
11 disp(Percentage , 'percentage of Si in Copper silicide
(Cu_5_Si)=')
```

Chapter 3

Crystal Geometry Structure and Defects

Scilab code Exa 3.10 Find the angle Between normals to the planes

```
1 //Exam:3.10
2 clc;
3 clear;
4 close;
5 // Miller indices of plane
6 h_1=1;
7 k_1=1;
8 l_1=1;
9 h_2=1;
10 k_2=2;
11 l_2=1;
12 angle=acosd((h_1*h_2+k_1*k_2+l_1*l_2)/(((h_1^2+k_1
    ^2+l_1^2)^(1/2))*( (h_2^2+k_2^2+l_2^2)^(1/2))) );
13 disp(angle,'angle Between normals to the planes
    (111) and (121)(in degrees)=');
```

Scilab code Exa 3.11 Determine the packing efficiency and density of sodium chloride

```
1 //Exam:3.11
2 clc;
3 clear;
4 close;
5 r_Na=0.98; //Radius of Na+(in A)
6 r_Cl=1.81; //Radius of Cl-(in A)
7 a=2*(r_Na+r_Cl); //Lattice parameter (in A)
8 pi=22/7;
9 V_i=4*(4/3)*pi*((r_Na^3)+(r_Cl^3)); //Volume of ions
    present in unit cell
10 V_u=a^3; //Volume of unit cell
11 Apf=V_i/V_u; //Atomic packing fraction
12 Ef_p=(Apf)*100; //Packing efficiency (in %)
13 AM_sodium=22.99; //Atomic mass of sodium (in amu)
14 AM_chlorine=35.45; //Atomic mass of chlorine (in amu)
15 M_1=4*(AM_sodium+AM_chlorine)*1.66*10^(-27); //Mass
    of the unit cell
16 a_1=a*10^(-10); //Lattice parameter (in meter)
17 V_u1=(a_1)^3;
18 Density=M_1/V_u1;
19 disp(Ef_p, 'Packing efficiency of sodium chloride (in
    %)=');
20 disp(Density, 'density of sodium chloride (in Kg/m3)='
    );
```

Scilab code Exa 3.12 Calculate the unit cell dimensions and atomic diameter

```
1 //Exam:3.12
2 clc;
3 clear;
4 close;
```

```

5 Density=2.7; // (in g/cm^3)
6 n=4;
7 m=26.98; //atomic weight of Al
8 N_a=6.023*10^(23); //avogadro number
9 a=((n*m/(Density*N_a))^(1/3)); //Lattice parameter (in
Cm)
10 A=a*10^(8); //Lattice parameter (in A)
11 disp(A, 'radius (in A)=');
12 r=A/(2*1.414); //radius for fcc structure
13 disp(2*r, 'Diameter (in A)=');

```

Scilab code Exa 3.13 Calculate the interplaner distance

```

1 //Exam:3.13
2 clc;
3 clear;
4 close;
5 r=1.245; //radius of nickel (in A)
6 a=4*r/(2)^(1/2); //Lattice constant (in A)
7 //Miller indices of plane 200
8 h_1=2;
9 k_1=0;
10 l_1=0;
11 //Miller indices of plane 111
12 h_2=1;
13 k_2=1;
14 l_2=1;
15 d_200=a/((h_1^2)+(k_1^2)+(l_1^2))^(1/2);
16 d_111=a/((h_2^2)+(k_2^2)+(l_2^2))^(1/2);
17 disp(d_200, 'interplaner distance of (200) plane of
nickel crystal (in A)=');
18 disp(d_111, 'interplaner distance of (111) plane of
nickel crystal (in A)=');

```

Scilab code Exa 3.14 Find the number of atoms per mm²

```
1 //Exam:3.14
2 clc;
3 clear;
4 close;
5 a=3.03*10^(-7); //lattice constant(in mm)
6 N_100=1/(a^2); //Number of atoms in the (100) plane
                  of a simple cubic structure
7 N_110=0.707/(a^2); //Number of atoms in the (110)
                  plane of a simple cubic structure
8 N_111=0.58/(a^2); //Number of atoms in the (111)
                  plane of a simple cubic structure
9 disp(N_100,'Number of atoms in the (100) plane of a
      simple cubic structure(in per mm^2)=');
10 disp(N_110,'Number of atoms in the (110) plane of a
      simple cubic structure(in per mm^2)=');
11 disp(N_111,'Number of atoms in the (111) plane of a
      simple cubic structure(in per mm^2)=');
```

Scilab code Exa 3.15 Determine the planer density of Ni

```
1 //Exam:3.15
2 clc;
3 clear;
4 close;
5 r=1.245*10^(-7); //Radius of the Ni atom(in mm)
6 NA_100=1+(1/4)*4; //Numbers of atom in (100) plane
7 a=4*r/(2)^(1/2); //Lattice constant(in mm)
8 Area=a^2;
9 P_density=NA_100/Area;
```

```
10 disp(P_density, 'the planer density of Ni (in atoms  
per mm^2)=');
```

Scilab code Exa 3.16 Calculate the planar atomic densities of planes

```
1 //Exam:3.16  
2 clc;  
3 clear;  
4 close;  
5 N_a1=4*(1/4)+1; //Number of atoms contained in (100)  
plane  
6 r=1.75*10^(-7); //radius of lead atom (in mm)  
7 a_1=2*2^(1/2)*r; //edge of unit cell in case of  
(100) plane  
8 PD_100=N_a1/(a_1^2); //Planar density of plane (100)  
9 N_a2=4*(1/4)+2*(1/2); //Number of atoms contained in  
(110) plane  
10 a_21=4*r; //top edge of the plane (110)  
11 a_22=2*2^(1/2)*r; //vertical edge of the plane (110)  
12 PD_110=N_a2/(a_21*a_22); //Planar density of plane  
(110)  
13 N_a3=3*(1/6)+3/2; //Number of atom contained in (111)  
plane  
14 Ar_111=4*(3^(1/2))*r^2; //area of (111) plane  
15 PD_111=N_a3/Ar_111; //Planar density of plane (111)  
16 disp(PD_100, 'Planar density of plane 100(in atoms/mm  
^2)=');  
17 disp(PD_110, 'Planar density of plane 110(in atoms/mm  
^2)=');  
18 disp(PD_111, 'Planar density of plane 111(in atoms/mm  
^2)=');
```

Scilab code Exa 3.17 Calculate the linear atomic densities of planes

```

1 //Exam:3.17
2 clc;
3 clear;
4 close;
5 N_a1=(1/2)+1+(1/2); //Number of diameters of atom
    along (110) direction
6 a=3.61*10^(-7); //lattice constant of copper in mm
7 L_d1=2^(1/2)*a; //length of the face diagonal in case
    of (110) direction
8 p_110=N_a1/L_d1; //linear atomic density along (110)
    of copper crystal lattice (in atoms/mm)
9 N_a2=(1/2)+(1/2); //Number of diameters of atom along
    (111) direction
10 L_d2=3^(1/2)*a; //length of the face diagonal in case
    of (111) direction
11 p_111=N_a2/L_d2; //linear atomic density along (110)
    of copper crystal lattice (in atoms/mm)
12 disp(p_110,'linear atomic density along (110) of
    copper crystal lattice (in atoms/mm)=');
13 disp(p_111,'linear atomic density along (111) of
    copper crystal lattice (in atoms/mm)=');

```

Scilab code Exa 3.18 Find lattice constant

```

1 //Exam:3.18
2 clc;
3 clear;
4 close;
5 A=55.8; //atomic weight of Fe
6 n=2; //number of atoms per unit cell
7 N=6.02*10^(26); //Avogadro's number
8 p=7.87*10^3; //density of Fe (in kg/m^3)
9 a=((A*n/(N*p))^(1/3))*10^10; //Value of lattice
    constant
10 disp(a,'Value of lattice constant (in A)=');

```

Scilab code Exa 3.19 Find the numbers of atoms per unit cell

```
1 //Exam:3.19
2 clc;
3 clear;
4 close;
5 a=2.9*10^(-10); //lattice parameter (in m)
6 A=55.8; //atomic weight of Fe
7 N=6.02*10^(26); //Avogadro's number
8 p=7.87*10^3; //density of Fe (in kg/m^3)
9 n=(a^3)*N*p/A; //Numbers of atoms per unit cell
10 disp(floor(n), 'Numbers of atoms per unit cell=');
```

Scilab code Exa 3.20 Calculate the line energy of disslocation in bcc iron

```
1 //Exam:3.20
2 clc;
3 clear;
4 close;
5 a=2.87*10^(-10); //lattice parameter for bcc iron
6 b=a*(3^(1/2))/2; //Magnitude of burgers vector
7 u=80*10^9; //shear modulus
8 E=(1/2)*u*b^2; //line energy of disslocation
9 disp(E, 'line energy of disslocation (in J/m)=');
```

Scilab code Exa 3.22 Calculate the number of vacancies

```
1 //Exam:3.22
2 clc;
```

```

3 clear;
4 close;
5 N=6.023*10^23; // avogadro number
6 T=1000; // absolute temperature
7 R=8.314; // constant
8 H_f=100*1000; // enthalpy of formation of vacancies (in
J/mol)
9 n=N*exp(-(H_f)/(R*T)); //number of vacancies created
during heating (in per mol)
10 V=5.5*10^(-6); //volume of 1 mole of the crystal in m
^3
11 n_1=n/V; //number of vacancies created during heating
(in per m^3)
12 disp(n_1, 'number of vacancies created during heating
(in per m^3)=');

```

Scilab code Exa 3.23 Calculate the surface energy of copper

```

1 //Exam:3.23
2 clc;
3 clear;
4 close;
5 //bond energy per atom of copper=bond energy per
bond*numbers of bond per atom*(1/2)
6 A=56.4*1000; //
7 N=6.023*10^23; // avogadro number
8 n_1=12; // numbers of bond per atom
9 n_2=3; // bonds broken at the surface
10 E=A*n_1/(2*N); // Energy of total bonds
11 E_b=E*(n_2/n_1); // Energy of broken bonds on surface
12 disp(E_b, 'E_b');
13 n_a=1.77*10^19; // no. of atoms on {111} planes in
copper (in m^-2)
14 E_c=n_a*E_b; // Surface energy (enthalpy) of copper
15 disp(E_c, 'Surface energy (enthalpy) of copper (in J/m

```

$\wedge 2) = ') ;$

Scilab code Exa 3.24 Calculate the equilibrium concentration of vacancies in aluminium

```
1 //Exam:3.24
2 clc;
3 clear;
4 close;
5 H_f=68*1000; //enthalpy of formation of vacancies (in
J/mol)
6 T_1=0; //temp (in K)
7 T_2=300; //temp (in K)
8 R=8.314; //constant
9 n=exp(-H_f/(R*T_2)); //equilibrium concentration of
vacancies in aluminium at 300 K
10 disp(n, 'equilibrium concentration of vacancies in
aluminium at 300 K=');
```

Scilab code Exa 3.25 Determine the interplanar spacing

```
1 //Exam:3.25
2 clc;
3 clear;
4 close;
5 Wavelength=1.54*10^(-10); //in meter
6 Angle=20.3; //in degree
7 n=1; //First order
8 d=Wavelength*n/(2*sind(Angle)); //the interplanar
spacing (in Meter)
9 disp(d/(10^-10), 'the interplanar spacing between
atomic plane (in A)=');
```

Scilab code Exa 3.26 Calculate the size of unit cell

```
1 //Exam:3.26
2 clc;
3 clear;
4 close;
5 wavelength=0.58; //in Angstrom
6 angle=9.5; //in degree
7 n=1; //First order
8 d_200=wavelength*n/(2*sind(angle)); //interplanar
      spacing(in Angstrom)
9 //Miller indices of plane
10 h=2;
11 k=0;
12 l=0;
13 a=d_200*(h^2+k^2+l^2)^(1/2); //Size of unit cell(in
      Angstrom)
14 disp(a, 'Size of unit cell(in Angstrom)=');
```

Scilab code Exa 3.27 Calculate the Bragg angle

```
1 //Exam:3.27
2 clc;
3 clear;
4 close;
5 //Miller indices of plane
6 h=1;
7 k=1;
8 l=1;
9 wavelength=0.54; //in angstrom
10 a=3.57; //size of a cube
11 n=1;
```

```
12 d_111=a/(h^2+k^2+l^2)^(1/2); //interplanar spacing (in  
Angstrom)  
13 angle=asind(n*wavelength/(2*d_111));  
14 disp(angle,'Bragg angle (in degree)=');
```

Scilab code Exa 3.28 Calculate the bragg reflection index

```
1 //Exam:3.28  
2 clc;  
3 clear;  
4 close;  
5 d=1.181; //  
6 wavelength=1.540; //in angstrom  
7 angle=90; //in degree  
8 n=2*d*sind(angle)/(wavelength); //the bragg  
reflection index  
9 disp(n,'bragg reflection index for BCC crystal=');
```

Scilab code Exa 3.29 Calculate the angle for 3rd order reflection

```
1 //Exam:3.29  
2 clc;  
3 clear;  
4 close;  
5 n_1=1; //1st order reflection index  
6 angle_1=10; //1st order reflection angle  
7 n_3=3; //3rd order reflection index  
8 //sind(angle_1)/sind(angle_3)=n_1/n_3  
9 angle_3=asind(n_3*sind(angle_1)/n_1); //  
10 disp(angle_3,'3rd order reflection angle=')
```

Scilab code Exa 3.30 Obtain the interplanar spacing and miller indices of the reflection plane

```
1 //Exam:3.30
2 clc;
3 clear;
4 close;
5 angle=20.3; //in degree
6 wavelength=1.54; //in angstrom
7 n=1;
8 a=3.16; //lattice parameter in angstrom
9 d=n*wavelength/(2*sind(angle)); //interplanar spacing
10 M_indices=a^2/(d^2);
11 disp(d, 'interplanar spacing of reflection plane');
12 disp(floor(M_indices), 'miller indices of the
    reflection plane');
13 disp((101),(110),(011));
```

Scilab code Exa 3.31 Determine interatomic spacing

```
1 //Exam:3.31
2 clc;
3 clear;
4 close;
5 //Miller indices of plane
6 n=1;
7 h=1;
8 k=1;
9 l=1;
10 angle=30; //in degree
11 wavelength=2; //in angstrom
12 d=n*wavelength/(2*sind(angle)); //interplanar spacing
13 a=d*(h^2+k^2+l^2)^(1/2); //interatomic spacing
14 disp(a, 'interatomic spacing (in angstrom)=');
```

Chapter 4

Bonds in solid

Scilab code Exa 4.1 Find A and B and the force needed to dissociate this molecule into atoms and the interatomic distance at which the dissociation occurs

```
1 //Exam:4.1
2 clc;
3 clear;
4 close;
5 r_o=2.8 //interatomic distance in
6 R_o=2.8*10^(-10); //interatomic distance in m
7 u_o=8; //released energy in eV
8 e=1.6*10^(-19); //charge of electron in C
9 U_o=8*e //released energy in Joule
10 A=(5/4)*U_o*(R_o^2); //proportionality constant for
    attraction in J-m2
11 B=A*(R_o^8)/5; //proportionality constant for
    repulsion in J-m2
12 r_c=(110*B/(6*A))^(1/8); //interatomic distance at
    which the dissociation occurs in m
13 F=-(2/r_c^3)*(A-5*B/(r_c^8)); //the force required to
    dissociate the molecule in N
14 disp(A,'proportionality constant for attraction (in
    J-m2)=');
```

```
15 disp(B,'proportionality constant for repulsion (in J  
-m2)=');  
16 disp(r_c,'interatomic distance at which the  
dissociation occurs (in m)=');  
17 disp(F,'the force required to dissociate the  
molecule (in N)=');
```

Scilab code Exa 4.2 Find the repulsive exponent n

```
1 //Exam:4.2  
2 clc;  
3 clear;  
4 close;  
5 r_o=3.14; //nearest neighbour equilibrium distance in  
6 R_o=3.14*10^(-10); //nearest neighbour equilibrium  
distance in m  
7 K=5.747*10^(-11); //compressibility of KCl in m2/N  
8 M=1.748; //Madelung constant  
9 pi=22/7;  
10 E_o=8.854*10^(-12);  
11 q=1.6*10^(-19); //electron charge  
12 n=1+18*(R_o^4)*4*pi*E_o/(K*M*q^2);  
13 disp(n,'repulsive exponent n=');
```

Scilab code Exa 4.3 Find the radius of Cl ion

```
1 //Exam:4.3  
2 clc;  
3 clear;  
4 close;  
5 F_1=3.02*10^(-9); //force of attraction b/w ions of  
Na+ and Cl-
```

```

6 Z_1=+1;
7 Z_2=-1;
8 e=1.6*10^(-19);
9 E_o=8.854*10^-12;
10 pi=22/7;
11 r_Na=0.95; //ionic radius of Na+ ion
12 r=(-Z_1*Z_2*e^2/(4*pi*E_o*F_1))^(1/2); //Radius of
    ion in meter
13 R=r/10^(-10); //Radius of ion in Angstrom
14 r_Cl=(R-r_Na); //Radius of Cl- ion in Angstrom
15 disp(r_Cl, 'Ionic Radius of Cl- ion (in Angstrom)=');

```

Scilab code Exa 4.4 Calculate the force of attraction between ions

```

1 //Exam:4.4
2 clc;
3 clear;
4 close;
5 Z_1=+2;
6 Z_2=-2;
7 r_Mg=0.65; //radius of Mg++ ion
8 r_S=1.84; //radius of S— ion
9 r=r_Mg+r_S; //net radius(in Angstrom)
10 R=r*10^(-10); //net radius(in meter)
11 e=1.6*10^(-19);
12 E_o=8.854*10^-12;
13 pi=22/7;
14 F=-Z_1*Z_2*e^2/(4*pi*E_o*R^2); //force of attraction
    between ions(in Newton)
15 disp(F, 'force of attraction between ions(in Newton)=
    ');

```

Scilab code Exa 4.5 How much net energy is spent in the process

```
1 //Exam:4.5
2 clc;
3 clear;
4 close;
5 //Na atom requires +5.14 eV of energy. When this
   electron is transferred to a vacant position , it
   gives back 4 .02 eV of energy
6 E_1=+5.14; //in eV
7 E_2=-4.02; //in eV
8 NET_energy=E_1+E_2; //in eV
9 disp(NET_energy , 'Net spent energy in whole process(
   in eV)=')
```

Scilab code Exa 4.6 Estimate the fraction of hydrogen bonds that are broken when ice melts

```
1 //Exam:4.6
2 clc;
3 clear;
4 close;
5 Enthalpy=6.02; //enthalpy of fusion of ice is 6.02 kJ
   /mol
6 E_h=20.5; //Hydrogen bond energy (in kJ/mol)
7 //There are two moles of hydrogen bonds per mole of
   H2O in ice .
8 H_b=Enthalpy/(2*E_h); //the fraction of hydrogen
   bonds that are broken when ice melts
9 disp(H_b , 'fraction of hydrogen bonds that are broken
   when ice melts=')
```

Chapter 5

Electron Theory of Metals

Scilab code Exa 5.1 Evaluate the temperature

```
1 //Exam:5.1
2 clc;
3 clear;
4 close;
5 //The probability that a particular quantum state at
   energy E is filled , is given by
6 //f(E) =1/(1+exp(E-E_f)/kT)
7 e=1.6*10^(-19); //charge on the electron
8 dE=0.5*e; //E-E_f in joule
9 //0.01=1/(1+exp(x))
10 //1+exp(x)=100
11 x=log(99);
12 k=1.38*10^(-23); //constant
13 T=dE/(x*k); //temperature
14 disp(ceil(T), 'temperature at which there is one per
   cent probability that a state with an energy 0.5
   eV above the Fermi energy will be occupied by an
   electron (in K)=');
```

Scilab code Exa 5.2 Find the drift velocity of carriers

```
1 //Exam:5.2
2 clc;
3 clear;
4 close;
5 n=10^19; //electrons per m^3
6 V=0.017; //applied voltage
7 d=0.27*10^-2; //distance with material
8 e=1.602*10^-19; //in coulomb
9 m=9.1*10^-31; //mass of an electron (in kg)
10 conductivity=0.01; //in mho.m^-1
11 E=V/d; //Electric field (in V/m)
12 v=(conductivity*E/(n*e))*10^2; //drift velocity of
   carriers (in meter/sec)
13 disp(v, 'drift velocity of carriers (in meter/sec)=');
```

Scilab code Exa 5.3 Find the conductivity of copper at 300K

```
1 //Exam:5.3
2 clc;
3 clear;
4 close;
5 T=300; //Temperature (in Kelvin)
6 t=2*10^-14; //time (in sec)
7 V_c=8.9; //volume of 63.54gm of copper (in cc)
8 Aw_c=63.54; //Atomic weight of copper (in a.m.u)
9 e=1.6*10^(-19);
10 m=9.1*10^-31;
11 N_a=6.023*10^23; //avogadro's number
12 n=(N_a/(Aw_c/V_c))*10^6; //Number of electrons per m
   ^3
13 conductivity=(e^2)*n*t/m; //conductivity of copper at
   300K (in mho/m)
14 disp(conductivity, 'conductivity of copper at 300K (in
```

mho/m)=') ;

Scilab code Exa 5.4 Find the mobility of conduction electron

```
1 //Exam:5.4
2 clc;
3 clear;
4 close;
5 t=10^(-14); //mean free time between the collisions (
    in second)
6 e=1.6*10^-19;
7 m=9.1*10^-31;
8 Mobility=e*t/m; //in m^2/V-s
9 disp(Mobility , 'mobility of conduction electron (in m
    ^2/V-s)=') ;
```

Scilab code Exa 5.5 Find the mobility of conduction electron and drift velocity

```
1 //Exam:5.5
2 clc;
3 clear;
4 close;
5 n=6*10^23; //conduction electron per m^3
6 conductivity=6.5*10^7; //in mho/m
7 E=1; //electric field intensity (in V/m)
8 e=1.6*10^-19;
9 m=9.1*10^-31;
10 Mobility=conductivity/(n*e); //in m^2/V-s
11 v=Mobility*E; //drift velocity (in m/sec)
12 disp(Mobility , 'mobility of conduction electron (in m
    ^2/V-s)=') ;
13 disp(v , 'drift velocity (in m/sec)=') ;
```

Scilab code Exa 5.6 Find the number of free electrons and also calculate mobility of electrons

```
1 //Exam:5.6
2 clc;
3 clear;
4 close;
5 d=10.5; // density of silver (in gm/cc)
6 At_w=107.9;
7 e=1.6*10^-19;
8 conductivity=6.8*10^5; //in mho/centimeter
9 N=6.023*10^23;
10 n=N*d/At_w; //number of free electrons
11 Mobility=conductivity/(n*e); //mobility of electrons(
    in cm^2/V-s);
12 disp(n,'number of free electrons=');
13 disp(Mobility,'mobility of electrons (in cm^2/V-s)' );
;
```

Scilab code Exa 5.7 Calculate the maximum velocity of an electron in a metal and mobility of electrons

```
1 //Exam:5.7
2 clc;
3 clear;
4 close;
5 E_f=3.75; //Fermi energy (in eV)
6 e=1.602*10^-19;
7 W_f=e*E_f; //fermi energy in joules
8 t=10^-14; //mean free time between the collisions (in
    second)
```

```
9 m=9.1*10^-31; //mass of electron
10 v_f=(2*W_f/m)^(1/2); //maximum velocity of an
    electron in a metal(in m/s)
11 mobility=e*t/m; //mobility of electrons(in m^2/V-s)
12 disp(v_f,'maximum velocity of an electron in a metal
    (in m/s)=');
13 disp(mobility,'mobility of electrons (in m^2/V-s)=')
```

Scilab code Exa 5.8 Calculate the velocity of an electrons at fermi level

```
1 //Exam:5.8
2 clc;
3 clear;
4 close;
5 E_f=2.1; //fermi energy (in eV)
6 e=1.602*10^-19;
7 m=9.1*10^-31;
8 W_f=e*E_f; //fermi energy in joules
9 v_f=(2*W_f/m)^(1/2); //velocity of an electrons at
    fermi level(in m/sec)
10 disp(v_f,'velocity of an electrons at fermi level(in
    m/sec)')
```

Scilab code Exa 5.9 Estimate the mean path of free electrons in pure copper

```
1 //Exam:5.9
2 clc;
3 clear;
4 close;
5 t=10^-9; //collision time(in seconds)
6 E_f=7; //fermi energy(in eV)
7 e=1.6*10^-19;
```

```
8 m=9.1*10^-31;
9 W_f=E_f*e; //fermi energy(in joules)
10 v_f=(2*W_f/m)^(1/2); //velocity of an electrons at
    fermi level(in m/sec)
11 P=v_f*t; //Mean free path(in meter)
12 disp(P, 'Mean free path(in meter)=')
```

Scilab code Exa 5.10 Find the conductivity of copper

```
1 //Exam:5.10
2 clc;
3 clear;
4 close;
5 N_a=6.023*10^23;
6 V_c=8.9; //volume of 63.54gm of copper(in cc)
7 Aw_c=63.54; //Atomic weight of copper(in a.m.u)
8 n=(N_a/(Aw_c/V_c))*10^6; //Number of electrons per m
    ^3
9 e=1.6*10^-19;
10 m=9.1*10^-31;
11 t=2*10^-14; //collision time
12 conductivity=n*(e^2)*t/m; //conductivity of copper
13 disp(conductivity, 'conductivity of copper(in ohm^-1/
    m)=');
```

Chapter 6

Photoelectric Effect

Scilab code Exa 6.1 Find the kinetic energy of electrons ejected from the surface

```
1 //Exam :6 . 1
2 clc;
3 clear;
4 close;
5 h=6.62*10^-34;
6 c=3*10^8;
7 e=1.6*10^-19;
8 Wavelength_1=2300*10^-10;
9 W=h*c/Wavelength_1; //Work function
10 Wavelength_2=1800*10^-10;
11 E_in=h*c/Wavelength_2;
12 E=E_in-W; //kinetic energy of the ejected electron (in
   Joules)
13 E_1=E/e; //kinetic energy of the ejected electron (in
   eV)
14 disp(E_1, 'kinetic energy of the ejected electron (in
   eV)=');
```

Scilab code Exa 6.2 Calculate the threshold frequency and the corresponding wavelength

```
1 //Exam:6.2
2 clc;
3 clear;
4 close;
5 h=6.625*(10^(-34)); //Planck's constant (in m2*kg/s)
6 c=3*10^8; //speed of light (in m/s)
7 e=1.602*10^-19; //electron charge (in coulomb)
8 W=2.3; //work (in eV)
9 W_1=W*e; //work (in joules)
10 v_o=W_1/h; //threshold frequency (in Hz)
11 Wavelength=(h*c/W_1)/10^(-10); //Wavelength in
    Angstrom
12 disp(v_o,'threshold frequency (Hz)=');
13 disp(Wavelength,'Wavelength (in Angstrom)=');
```

Scilab code Exa 6.3 Calculate the threshold frequency and the work function of metal

```
1 //Exam:6.3
2 clc;
3 clear;
4 close;
5 h=6.625*(10^(-34)); //Planck's constant (in m2*kg/s)
6 c=3*10^8; //speed of light (in m/s)
7 e=1.602*10^-19; //electron charge (in coulomb)
8 wavelength=6800*10^-10; //wavelength of radiation
9 v_o=c/wavelength; //frequency
10 W=h*v_o; //Work function
11 disp(v_o,'threshold frequency (in Hz)=')
12 disp(W,'work function of metal (in joule)=')
```

Scilab code Exa 6.4 Calculate the photons emitted by lamp per second

```
1 //Exam:6.4
2 clc;
3 clear;
4 close;
5 h=6.625*(10^(-34)); //Planck's constant (in m2*kg/s)
6 c=3*10^8; //speed of light (in m/s)
7 L_r =150*8/100; //Lamp rating (in joule)
8 wavelength=4500*10^-10; //in meter
9 W=h*c/wavelength; //work function
10 N=L_r/W; //number of photons emitted by lamp per
    second
11 disp(N, 'number of photons emitted by lamp per second
    =' )
```

Scilab code Exa 6.5 Determine the region of electrons spectrum

```
1 //Exam:6.5
2 clc;
3 clear;
4 close;
5 h=6.6*(10^(-34)); //Planck's constant (in m2*kg/s)
6 c=3*10^8; //speed of light (in m/s)
7 e=1.6*10^-19; //electron charge (in coulomb)
8 W=2.24; //work function (in eV)
9 W_1=W*e; //work function (in joule)
10 v=(W_1/h)*10^-10; //frequency
11 wavelength=c/v; //region of electrons spectrum is
    less than (in angstrom)
12 disp(wavelength, 'region of electrons spectrum is
    less than (in angstrom) ')
```

Scilab code Exa 6.6 Calculate the photons emitted by radio receiver

```
1 //Exam:6.6
2 clc;
3 clear;
4 close;
5 h=6.625*(10^(-34)); //Planck's constant (in m2*kg/s)
6 c=3*10^8; //speed of light (in m/s)
7 P_o=10*10^3; //Power of radio receiver (in Watt)
8 v=440*10^3; //Operating frequency
9 E=h*v; //Energy of each electron
10 N=P_o/E; //Number of photons emitted/sec
11 disp(N, 'Number of photons emitted/sec by radio
    receiver=')
```

Scilab code Exa 6.7 Calculate the wavelength of light which can just eject electron from tungsten and from barium

```
1 //Exam:6.7
2 clc;
3 clear;
4 close;
5 W_t=4.52; //Work function for tungsten (in eV)
6 W_b=2.5; //Work function for barium (in eV)
7 h=6.62*(10^(-34)); //Planck's constant (in m2*kg/s)
8 c=3*10^8; //speed of light (in m/s)
9 e=1.6*10^-19; //electron charge (in coulomb)
10 W_T=W_t*e; //Work function for tungsten (in Joule)
11 W_B=W_b*e; //Work function for barium (in Joule)
12 Wavelength_T=(h*c/W_T)*10^10; //wavelength of light
    which can just eject electron from tungsten
```

```
13 Wavelength_B=(h*c/W_B)*10^10; //wavelength of light  
    which can just eject electron from barrium  
14 disp(Wavelength_T , 'wavelength of light which can  
    just eject electron from tungsten(in Angstrom)=')  
15 disp(Wavelength_B , 'wavelength of light which can  
    just eject electron from barrium(in Angstrom)=')
```

Chapter 7

Diffusion in Solids

Scilab code Exa 7.1 find the time required for carburization

```
1 //Exam:7.1
2 clc;
3 clear;
4 close;
5 D=1.28*10^(-11); //diffusion coefficient of carbon in
                     given steel in m2/s
6 c_s=0.9; //Surface concentration of diffusion element
            in the surface
7 c_o=0.2; //Initial uniform concentration of the
            element in the solid
8 c_x=0.4; //Concentration of the diffusing element at a
            distance x from the surface
9 x=0.5*10^(-3); //depth from the surface in m
10 //((c_s-c_x)/(c_s-c_o))=erf(x/(2*(D*t)(1/2)))
11 t=(x/(2*erfinv((c_s-c_x)/(c_s-c_o))*D^(1/2)))^2; //
            time required for carburization(in sec)
12 disp(t,'time required for carburization(in sec)=');
```

Scilab code Exa 7.2 Calculate time required to get a boron content of 1023 atoms per m³ at a depth of 2 micro meter

```

1 //Exam:7.2
2 clc;
3 clear;
4 close;
5 D=4*10^(-17); //diffusion coefficient of carbon in
     given steel in m2/s
6 c_s=3*10^26; //Surface concentration of boron atoms
     in the surface
7 c_1=0; //Initial uniform concentration of the element
     in the solid
8 c_x=10^23; //Concentration of the diffusing element
     at a distance x from the surface
9 x=2*10^(-6); //depth from the surface in m
10 //((c_s-c_x)/(c_s-c_1))=erf(x/(2*(D*t)^(1/2)))
11 a=erfinv((c_s-c_x)/(c_s-c_1));
12 disp(a,'==')
13 t=(x^2/(D*4*(2.55)^2)); //time required to get a
     boron content of 1023 atoms per m3 at a depth of
     2 micro meter
14 disp(t,'time required to get a boron content of 1023
     atoms per m3 at a depth of 2 micro meter(in sec)
     ==');
15 disp((c_s-c_x)/(c_s-c_1));
16 T=(x/(2*(2.55)*D^(1/2)))^2;
17 disp(T,'==')
```

Scilab code Exa 7.3 find the constant of the equation and activation energy and determine the diffusion coefficient

```

1 //Exam:7.3
2 clc;
3 clear;
```

```

4 close;
5 t_1=736; //Temperature in C
6 t_2=782; //Temperature in C
7 T_1=t_1+273; //Temperature in K
8 T_2=t_2+273; //Temperature in K
9 D_1=2*10^(-13); //Coefficient of diffusion at T_1 (in
m2/s)
10 D_2=5*10^(-13); //Coefficient of diffusion at T_2 (in
m2/s)
11 k=1.38*10^(-23); //in J/K
12 //log(d_1)=log(d_o)-E/(k*T_1)
13 //log(d_2)=log(d_o)-E/(k*T_2)
14 E=(log(D_1)-log(D_2))/((1/(k*T_1))-(1/(k*T_2))); //
15 disp(E, 'activation energy (in J)=');
16 D_o=2*10^(-13)/exp(E/(k*T_1));
17 disp(D_o, 'constant of the equation (in m2/s)' );
18 t_4=500; //Temperature in C
19 T_4=t_4+273; //Temperature in K
20 D_4=D_o*exp(E/(k*T_4)); //diffusion coefficient at
500 C
21 disp(D_4, 'diffusion coefficient at 500 C (in m2/s)' )

```

Scilab code Exa 7.4 Determine the approximate time that will produce same diffussion at 500 C

```

1 //Exam:7.4
2 clc;
3 clear;
4 close;
5 D_500=4.8*10^(-14); //Diffusion coefficient for
copper in aluminimium at 500*C(in m2/s)
6 D_600=5.3*10^(-13); //Diffusion coefficient for
copper in aluminimium at 600*C(in m2/s)
7 t_600=10; //time of diffussion at 600*C(in Hours)

```

```
8 //D_500*t_500=D_600*t_600
9 t_500=D_600*t_600/D_500; //time of diffussion at 500*
C
10 disp(t_500,'Time at 500*C that will produce the same
diffusion as in 600*C(in Hours)=');
```

Chapter 8

Mechanical Properties of Materials and Mechanical Tests

Scilab code Exa 8.1 Determine the fracture strength

```
1 //Exam:8.1
2 clc;
3 clear;
4 close;
5 Y=180*10^9; //Young's modulus of a certain material(
    in N/m^2)
6 E=1.8; //true surface energy (in J/m^2)
7 c=(5/2)*10^-6; //Crack (in meter)
8 pi=3.14;
9 F_strength=(2*Y*E/(pi*c))^(1/2);
10 disp(F_strength*10^-6, 'fracture strength (in MN/m^2)=
    ');
```

Scilab code Exa 8.2 Compute the true stress and strain with engineering stress and strain

```

1 //Exam:8.2
2 clc;
3 clear;
4 close;
5 d_o=12.7; // tensile test specimen diameter (in mm)
6 d=12; // tensile test specimen diameter after load (in
      mm)
7 P=76*10^3; //load (in N)
8 pi=22/7;
9 A_o=(pi/4)*(d_o^2); //Initial area of cross section (
      in mm^2)
10 A=(pi/4)*(d^2); //area of cross section after load of
      76 kN
11 E_stress=P/A_o; //engineering stress
12 T_stress=P/A; //true stress
13 T_strain=log(A_o/A); //true strain
14 E_strain=exp(T_strain)-1; //engineering strain
15 disp(E_stress,'engineering stress (in N/mm^2)=');
16 disp(T_stress,'true stress (in N/mm^2)=');
17 disp(E_strain,'engineering strain=');
18 disp(T_strain,'true strain=');

```

Scilab code Exa 8.3 Determine the fracture strength

```

1 //Exam:8.3
2 clc;
3 clear;
4 close;
5 Y=210*10^9; //Young's modulus of a certain material(
      in N/m^2)
6 E=10; //true surface energy (in J/m^2)
7 c=(100/2)*10^-6; //Crack (in meter)
8 pi=3.14;
9 F_strength=(2*Y*E/(pi*c))^(1/2);
10 disp(F_strength,'fracture strength (in Newton/m^2)=')

```

;

Scilab code Exa 8.4 Find the resultant elongation

```
1 //Exam:8.4
2 clc;
3 clear;
4 close;
5 l_o=305*10^-3; //length of copper piece(in meter)
6 E=110*10^9; //surface energy
7 stress=276*10^6; //in Pa
8 dl=stress*l_o/E; //resultant elongation(in meter)
9 disp(dl*10^3, 'resultant elongation(in mm)=');
```

Scilab code Exa 8.5 Compute the strain hardening exponent

```
1 //Exam:8.5
2 clc;
3 clear;
4 close;
5 T_stress=415; //True stress (in Megapascal)
6 T_strain=0.10; //True strain
7 K=1035; //(in Megapascal)
8 n=(log(T_stress)-log(K))/log(T_strain); //
9 disp(n, 'Strain hardening exponent for an alloy=')
```

Chapter 9

Alloys Systems Phase Diagrams and Phase Transformations

Scilab code Exa 9.1 Find the percentage of proeutectoid ferrite

```
1 //Exam:9.1
2 clc;
3 clear;
4 close;
5 //Fulcrum is at 0.5% carbon
6 //from lever rule
7 Pro_f=((0.80-0.5)/(0.80-0.0))*100; // % Proeutectoid
    ferrite
8 Pea_f=100-Pro_f; // % Pearlite ferrite
9 disp(Pro_f ,'% Proeutectoid ferrite=');
10 disp(Pea_f ,'% Pearlite ferrite=');
```

Scilab code Exa 9.2 Determine the degrees of freedom of a system of two components

```
1 //Exam:9.2
```

```

2 clc;
3 clear;
4 close;
5 N=2;
6 C=2;
7 //F=C-P+N
8 P_1=1;
9 P_2=2;
10 P_3=3;
11 P_4=4;
12 F_1=C-P_1+N;
13 F_2=C-P_2+N;
14 F_3=C-P_3+N;
15 F_4=C-P_4+N;
16 disp(F_1, 'Degrees of freedom for 1 phase=');
17 disp(F_2, 'Degrees of freedom for 2 phases=');
18 disp(F_3, 'Degrees of freedom for 3 phases=');
19 disp(F_4, 'Degrees of freedom for 4 phases=');

```

Scilab code Exa 9.3 Find the minimum number of components in the system

```

1 //Exam:9.3
2 clc;
3 clear;
4 close;
5 P=4; //Number of phases exhibit by a material
6 F=0; //Minimum degrees of freedom
7 //modified form of the phase rule F=C-P+1
8 C=F+P-1; //minimum number of components in the system
9 disp(C, 'the minimum number of components in the
system=')

```

Chapter 10

Heat Treatment

Scilab code Exa 10.1 Determine the grain diameter of an ASTM Number 8

```
1 //Exam:10.1
2 clc;
3 clear;
4 close;
5 N=8; //ASTM grain size number
6 n=2^(N-1); //Number of grains per inch square at a
               magnification
7 N_1=n*100*100; //Number of grains per inch square
                   without magnification
8 N_2=N_1/(25.4)^2; //Number of grains per mm square
                   without magnification
9 A_a=1/(N_2); //Average area of each grain(in mm^2)
10 D=(A_a)^(1/2); //Average grain diameter(in mm)
11 disp(D, 'Average grain diameter(in mm)=')
```

Chapter 11

Deformation of Materials

Scilab code Exa 11.1 Determine the value of critical resolved shear stress

```
1 //Exam:11.1
2 clc;
3 clear;
4 close;
5 h_1=1;
6 k_1=1;
7 l_1=1;
8 //Miller indices of slip plane
9 h_2=1;
10 k_2=-1;
11 l_2=1;
12 //Miller indices of stress plane
13 h_3=1;
14 k_3=1;
15 l_3=0;
16 //Miller indices of slip direction
17 A=(h_1*h_2+k_1*k_2+l_1*l_2)/(((h_1^2+k_1^2+l_1^2)
    ^^(1/2))*((h_2^2+k_2^2+l_2^2)^(1/2))); //Value of
    cos(x) where x =angle between slip plane and
    stress plane
18 B=(h_1*h_3+k_1*k_3+l_1*l_3)/(((h_1^2+k_1^2+l_1^2)
```

```


$$^{(1/2)} * ((h_3^2 + k_3^2 + l_3^2)^{(1/2)}) ; // Value of$$

cos(y) where y = angle between slip direction and
stress direction
19 C = (1 - A^2)^{(1/2)} ; // Value of sin(x)
20 stress = 3.5 ; // Applied Stress in MPa
21 T_cr = stress * A * B * C ; // Critical resolved shear stress (
    in MPa)
22 disp(T_cr, 'Critical resolved shear stress (in MPa)=')
;
```

Scilab code Exa 11.3 Find the yield stress

```

1 //Exam:11.3
2 clc;
3 clear;
4 close;
5 D=0.002; //Grain diameter (in mm)
6 d=D*10^(-3); //Grain diameter (in m)
7 K=0.63; //Constant (in MNm^(-3/2))
8 sigma_i=80; //in MNm^-2
9 sigma_y=sigma_i+K*d^(-1/2); //Yield stress for a
    polycrystalline alloy
10 disp(sigma_y, 'Yield stress for a polycrystalline
    alloy (in MN/m^2)' );
;
```

Scilab code Exa 11.4 Find the yield stress

```

1 //Exam:11.4
2 clc;
3 clear;
4 close;
5 sigma_y1=120; //primary yield strength of
    polycrystalline material (in MN*m^-2)
```

```

6 sigma_y2=220; //increased yield strength of
    polycrystalline material(in MN*m^-2)
7 d_1=0.04*10^(-3); //primary grain diameter(in meter)
8 d_2=0.01*10^(-3); //grain diameter after decreasing(
    in meter)
9 //sigma_y1=sigma_i+K*(d_1)^(-1/2)
10 //sigma_y2=sigma_i+K*(d_2)^(-1/2)
11 //putting the values and solving the equation
12 K=(220-120)/((d_2^(-1/2))-((d_1^(-1/2)))); //constant
    (in MN*m(-3/2))
13 sigma_i=sigma_y1-K*(d_1)^(-1/2); //in MN*m^-2
14 d=1/((10^4)*(256/645))^(1/2); //grain diameter for
    grain size ASTM 9(in mm)
15 D=d*10^(-3); //grain diameter for grain size ASTM 9(
    in meter)
16 sigma_y=sigma_i+K*(D)^(-1/2); //Yield stress for a
    polycrystalline alloy for grain size ASTM 9(in MN
    *m^-2)
17 disp(ceil(sigma_y), 'Yield stress for a
    polycrystalline alloy for grain size ASTM 9(in MN
    *m^-2)= ');

```

Chapter 12

Oxidation and Corrosion

Scilab code Exa 12.1 Find the distance at which magnesium anode capable of giving 2MA

```
1 //Exam:12.1
2 clc;
3 clear;
4 close;
5 D=320*10^-3; //in meter
6 L=1; //in meter
7 A=%pi*D*L; //Surface area in meter^2
8 l=ceil (200/A);
9 disp(l,'the distance at which magnisium anode
    capable of giving 2MA (in meters)=');
```

Scilab code Exa 12.2 Calculate the quality of magnesium required per square meter of the hull surface for a design life of 10 years

```
1 //Exam:12.2
2 clc;
3 clear;
```

```
4 close;
5 W=0.0243; //1 mole of magnesium weight (in Kg)
6 C=2*96490; //used charge (in A-s)
7 A=15*10^(-3); //current density (in A/metre2)
8 t=10; //time (in years)
9 T=10*365*24*3600; //time (in sec)
10 //amount of magnesium required =charge required per
    m2 of hull surface for a design life of 10 years
    /(used charge for anode)
11 Mg_required=W*A*T/C; //magnesium required per square
    meter of the hull surface for a design life of 10
    years
12 disp(Mg_required, 'magnesium required per square
    meter of the hull surface for a design life of 10
    years (in Kg/m2)=');
```

Chapter 13

Thermal and Optical Properties of Materials

Scilab code Exa 13.1 Determine the maximum temperature to which the rod may be heated without exceeding a compressive stress of 172 MPa

```
1 //Exam:13.1
2 clc;
3 clear;
4 close;
5 alpha=20*10^(-6); //linear coefficient of thermal
                     expansion per C
6 Sigma=-(172); //compressive stress MPa
7 T=20; //Temprature at which rod is stress free(in C
         )
8 E=100*10^3; //modulus of elasticity (in MPa)
9 T_f=T-(Sigma/(alpha*E)); //maximum temperature the
                           rod may be heated without exceeding a compressive
                           stress of 172 MPa
10 disp(T_f, 'maximum temperature(in C) the rod may be
          heated without exceeding a compressive stress of
          172 MPa=');
```

Chapter 14

Electrical and Magnetic Properties of Materials

Scilab code Exa 14.1 Calculate the resistance of an aluminium wire

```
1 //Exam:14.1
2 clc;
3 clear;
4 close;
5 l=100; //length of wire
6 p=2.66*10^(-8); //resistivity
7 A=3*10^(-6); //cross sectional area
8 R=p*l/A; //resistance of an aluminium wire
9 disp(R,'resistance of an aluminium wire(in Ohm)=');
```

Scilab code Exa 14.2 Find the resistivity of a copper alloy containing 1 atomic percent nickel and 3 atomic percent silver

```
1 //Exam:14.2
2 clc;
3 clear;
```

```

4 close;
5 R_Cu=1.56; // Resistivity of pure copper (in micro-ohm-
cm)
6 R_CuNi = 4.06; // Resistivity of Cu containing two
atomic percent (in micro-ohm-cm)
7 R_Ni=(R_CuNi-R_Cu)/2; // Increase in resistivity due
to one atomic % Ni
8 R_CuAg= 1.7; // resistivity of copper , containing one
atomic percent silver (in micro-ohm-cm)
9 R_Ag=R_CuAg-R_Cu; // Increase in resistivity due to
one atomic % Ag
10 R_CuNiAg=R_Cu+R_Ni+3*R_Ag; // Resistivity of copper
alloy containing one atomic percent Ni and 3
atomic percent Ag
11 disp(R_CuNiAg,' Resistivity of copper alloy
containing one atomic percent Ni and 3 atomic
percent Ag(in micro-ohm-cm)=')

```

Scilab code Exa 14.3 Find the resistivity due to impurity scattering per percent of Ni in the Cu lattice

```

1 //Exam:14.3
2 clc;
3 clear;
4 close;
5 R_Cu=1.8*10^(-8); // resistivity of pure copper at
room temperature
6 R_CuNi=7*10^(-8); // resistivity of Cu 4% Ni alloy at
room temperature
7 R_Ni=(R_CuNi-R_Cu)/4; // resistivity due to Impurity
scattering per % of Ni
8 disp(R_Ni,' resistivity due to impurity scattering
per percent of Ni in the Cu lattice (in ohm-meter)
= ')

```

Scilab code Exa 14.4 Calculate the relative dielectric constant of a barium titanate crystal

```
1 //Exam:14.4
2 clc;
3 clear;
4 close;
5 C=10^(-9); //capacitance (in F)
6 d=2*10^(-3); //distance of separation in a parallel
    plate condenser
7 E_o=8.854*10^(-12); //dielectric constant
8 A=(10*10^(-3))*(10*10^(-3)); //area of parallel plate
    condenser
9 //C=E_o*E_r*A/d
10 E_r=C*d/(E_o*A); //Relative dielectric constant
11 disp(ceil(E_r), 'Relative dielectric constant of a
    barium titanate crystal')
```

Scilab code Exa 14.5 Calculate the polarization

```
1 //Exam:14.5
2 clc;
3 clear;
4 close;
5 q=1.6*10^(-19); //charge (in C)
6 d_1=0.06 //shift of the titanium ion from the body
    centre (in )
7 d_2=0.08 //shift of the oxygen anions of the side
    faces (in )
8 d_3=0.06 //shift of the oxygen anions of the top and
    bottom face (in )
```

```

9 D_1=d_1*10^(-10); // shift of the titanium ion from
the body centre (in m)
10 D_2=d_2*10^(-10); // shift of the oxygen anions of the
side faces (in m)
11 D_3=d_3*10^(-10); // shift of the oxygen anions of the
top and bottom face (in m)
12 U_1=4*q*D_1; // dipole moment due to two O2 ions on
the four side faces (in C-m)
13 U_2=2*q*D_2; // dipole moment due to one O2 on top
and bottom (in C-m)
14 U_3=4*q*D_3; // dipole moment due to one Ti4+ ion at
body centre (in C-m)
15 U=U_1+U_2+U_3; // Total dipole moment (in C-m)
16 V=4.03*((3.98)2)*10^(-30); // volume (in m3)
17 P=U/V; // polarization the total dipole moments per
unit volume
18 disp(P, 'polarization (in C/m2)=');
19 disp(U, '==')

```

Scilab code Exa 14.6 Find net magnetic moment per iron atom in the crystal

```

1 //Exam:14.6
2 clc;
3 clear;
4 close;
5 V=((2.87)3)*10^(-30) //Volume of unit cell of BCC
iron (in m3)
6 N=2 //Number of atoms in the unit cell
7 M=1750*103; //saturation magnetization of BCC Iron A
/m
8 M_Net=V*M*(1/N) //net magnetic moment per atom
9 Bohr_magneton=9.273*10^(-24); //Bohr_magneton (
magnetic moment) in A/m2
10 M_moment=M_Net/Bohr_magneton; //The magnetic moment (

```

```
    in units of U_B)
11 disp(M_moment,'The magnetic moment (in units of U_B)
=');
```

Scilab code Exa 14.7 Calculate the saturation magnetization and the saturation flux density

```
1 //Exam:14.7
2 clc;
3 clear;
4 close;
5 p=8.90*10^6; //density of nickel in gm/m3.
6 N_A=6.023*10^23; //Avogadro's number atoms/mol
7 At_w=58.71; //Atomic weight of Ni in gm/mol
8 N=p*N_A/At_w; //number of atoms/m3
9 U_B=9.273*10^(-24); //Bohr_magneton
10 M_s=0.60*U_B*N; //saturation magnetization
11 pi=22/7;
12 U_o=4*pi*10^(-7); //magnetic constant
13 B_s=U_o*M_s; //Saturation flux density
14 disp(M_s,'the saturation magnetization=');
15 disp(B_s,'Saturation flux density=');
```

Scilab code Exa 14.8 Calculate the saturation magnetization

```
1 //Exam:14.8
2 clc;
3 clear;
4 close;
5 //Each cubic unit cell of ferrous ferric oxide
//contains 8 Fe2+ and 16 Fe3+ ions and
6 n_b=32; //
7 U_B=9.273*10^(-24); //Bohr_magneton
```

```
8 a=0.839*10^(-9); //the unit cell edge length in m
9 V=a^3; //volume(in m3)
10 M_s=n_b*U_B/V; //the saturation magnetization
11 disp(M_s,'the saturation magnetization=');
```

Scilab code Exa 14.9 Calculate eddy current loss at the normal voltage and frequency

```
1 //Exam:14.9
2 clc;
3 clear;
4 close;
5 //hysteresis loss (Ph) and the induced emf loss (Pe)
//are proportional to the frequency
6 //Pe is proportional to the square of the induced
//emf (Pe)
7 //Pe + Ph = 750 W (at 25 Hz)
8 //4Pe + 2Ph = 2300 W(at 50Hz)
9 //solving equation
10 P_e=800/2; //induced emf loss
11 I_d=4*P_e; //The eddy current loss at the normal
//voltage and frequency
12 disp(I_d,'The eddy current loss at the normal
//voltage and frequency (in W)=');
```

Chapter 15

Semiconductors

Scilab code Exa 15.1 Find the conductivity and resistivity of a pure silicon crystal

```
1 //Exam:15.1
2 clc;
3 clear;
4 close;
5 U_n=1350 // mobility of electron in cm2/volt-sec
6 U_h=480 // hole mobility in cm2/volt-sec
7 Sigma=1.072*1010 //density of electron hole pair per
cc at 300 K for a pure silicon crystal
8 e=1.6*10-19; //charge on the electron in C
9 Sigma_i=Sigma*e*(U_n+U_h); //Conductivity of pure
silicon crystal
10 p_i=1/(Sigma_i); //Resistivity of silicon crystal in
Ohm-cm
11 P_i=p_i*10-2; //Resistivity of silicon crystal in
Ohm-m
12 disp(Sigma_i,'Conductivity of pure silicon crystal(
in mho/cm)=');
13 disp(P_i,'Resistivity of silicon crystal (in Ohm-m)=
');
```

Scilab code Exa 15.2 Find the resistivity at room temperature

```
1 //Exam:15.2
2 clc;
3 clear;
4 close;
5 U=1200; //electron mobility in cm2/Volt-sec
6 e=1.6*10^(-19); //charge on the electron in C
7 n=10^13; //concentration of phosphorus
8 sigma=U*e*n; //conductivity of crystal in mho/cm
9 p_i=1/sigma; //resistivity of silicon wafer if all
    donor atom are active
10 disp(p_i,'resistivity of silicon wafer if all donor
    atom are active(in ohm-cm)=');
```

Scilab code Exa 15.3 Find the resistance of an intrinsic germanium rod

```
1 //Exam:15.3
2 clc;
3 clear;
4 close;
5 U_n=3900 //mobility of electron in cm2/volt-sec
6 U_h=1900 //hole mobility in cm2/volt-sec
7 n_i=2.5*10^13; //concentration of electron
8 u_n=U_n*10^(-4); //mobility of electron in m2/volt-
    sec
9 u_h=U_h*10^(-4); //hole mobility in m2/volt-sec
10 e=1.6*10^(-19); //charge on the electron in C
11 Sigma_i=n_i*e*(u_n+u_h)*10^6; //Conductivity
12 p_i=1/(Sigma_i); //resistivity of intrinsic germanium
    rod
13 l=1*10^(-2); //length of germanium rod in m
```

```
14 w=1*10^(-3); //width of germanium rod in m
15 t=1*10^(-3); //thick of germanium rod in m
16 A=w*t; //Area of cross section in m2
17 R=p_i*1/A; //Resistance of an intrinsic germanium rod
    in Ohm
18 disp(R/10^3, 'Resistance of an intrinsic germanium
    rod (in K-Ohm)=');
```

Scilab code Exa 15.4 Obtain density relation in P type material

```
1 //Exam:15.4
2 clc;
3 clear;
4 close;
5 N_a=1.1*10^20; //acceptor density in atoms/m3
6 n_i=2.5*10^19; //concentration of majority carrier
    per m3
7 n_p=(n_i^2)/N_a; //intrinsic density
8 R=n_p/n_i; //Ratio of n_p and n_i
9 disp(R, 'n_p/n_i=');
```

Chapter 16

Superconductivity and Superconducting Materials

Scilab code Exa 16.1 Calculate the energy gap in electron volts and Calculate the wavelength of a photon

```
1 //Exam:16.1
2 clc;
3 clear;
4 close;
5 T_c=4.2; // critical temperature of mercury
6 k=1.4*10^(-23); //
7 E_g=3*k*T_c; //energy gap (in Joule)
8 e=1.6*10^(-19); //charge on the electron
9 E=E_g/e; //energy gap (in electron volt)
10 h=6.6*10^(-34) // in J-s
11 c=3*10^8; //in m/s
12 wavelength=h*c/E_g; //wavelength of a photon (in m)
13 disp(E,'energy gap (in electron volt)=');
14 disp(wavelength,'wavelength of a photon (in m)=');
```

Chapter 18

Composites

Scilab code Exa 18.1 Calculate the modulus of elasticity

```
1 //Exam:18.1
2 clc;
3 clear;
4 close;
5 E_f=69; //modulus of elasticity in GPa
6 V_f=40/100; //Volume of glass fibres %
7 E_m=3.4; //modulus (in GPa)
8 V_m=60/100; //Volume of polyester resin %
9 E_cl=E_m*V_m+E_f*V_f; //modulus of elasticity (in Gpa
    )
10 disp(ceil(E_cl), 'modulus of elasticity (in Gpa)=') ;
```

Scilab code Exa 18.2 Calculate the elastic modulus when the stress is applied perpendicular to the direction of the fibre alignment

```
1 //Exam:18.2
2 clc;
3 clear;
```

```
4 close;
5 E_f=69; //modulus of elasticity in GPa
6 V_f=40/100; //Volume of glass fibres %
7 E_m=3.4; //modulus (in GPa)
8 V_m=60/100; //Volume of polyester resin %
9 E_cl=E_m*E_f/(E_m*V_f+E_f*V_m); //modulus of
    elasticity when the stress is applied
    perpendicular to the direction of the fibre
    alignment(in Gpa)
10 disp(E_cl,'modulus of elasticity when the stress is
    applied perpendicular to the direction of the
    fibre alignment(in Gpa)=');
```
