

Scilab Textbook Companion for  
Applied Physics For Engineers  
by N. Mehta<sup>1</sup>

Created by  
Divya Sharma  
Modern Physics  
Physics  
Shri Mata Vaishno Devi University  
College Teacher  
Mr. Pankaj Biswas  
Cross-Checked by  
Chaya Ravindra

March 14, 2014

<sup>1</sup>Funded by a grant from the National Mission on Education through ICT, <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab codes written in it can be downloaded from the "Textbook Companion Project" section at the website <http://scilab.in>

# Book Description

**Title:** Applied Physics For Engineers

**Author:** N. Mehta

**Publisher:** Phi Learning Pvt. Ltd., New Delhi

**Edition:** 1

**Year:** 2011

**ISBN:** 978-81-203-4242-2

Scilab numbering policy used in this document and the relation to the above book.

**Exa** Example (Solved example)

**Eqn** Equation (Particular equation of the above book)

**AP** Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

# Contents

List of Scilab Codes	5
1 Relativistic Mechanics	14
2 Quantum Mechanics	24
3 Statistical Mechanics	31
4 Geometrical Optics	41
5 Physical Optics	57
6 X Rays	82
7 Lasers and Holography	88
8 Ultrasonics	91
9 Fibre Optics	97
10 Electrostatics	106
11 Electromagnetic Theory	113
12 Magnetic Properties of Materials	120
13 Dielectric Properties of Materials	128
14 Solid State Electronics	137

<b>15 Digital Electronics</b>	<b>149</b>
<b>16 Crystal Physics</b>	<b>162</b>
<b>17 Nuclear Physics</b>	<b>170</b>
<b>19 Superconductivity</b>	<b>188</b>

# List of Scilab Codes

Exa 1.2	Lorentz transformations for space and time . . . . .	14
Exa 1.4	Relative speed of one photon with respect to another .	15
Exa 1.6	Areal contraction of moving circular lamina . . . . .	15
Exa 1.7	Length of a one metre stick moving parallel to its length	16
Exa 1.8	Mean lifetime of meson in motion . . . . .	17
Exa 1.9	Speed at which a moving clock ticks slow . . . . .	17
Exa 1.10	Distance that the meson beam can travel before reduction in its flux . . . . .	18
Exa 1.11	Velocity of the particle when its total energy is thrice its rest energy . . . . .	19
Exa 1.12	Kinetic energy and momentum of moving electron . .	19
Exa 1.13	Amount of work to be done to increase the speed of an electron . . . . .	20
Exa 1.14	Particle moving with relativistic speed . . . . .	21
Exa 1.15	Speed of the electron in order to have its mass equal to mass of a proton . . . . .	22
Exa 1.17	Classical and relativistic speed of an electron of given kinetic energy . . . . .	22
Exa 2.1	de broglie wavelength of an electron . . . . .	24
Exa 2.2	de broglie wavelength associated with a proton . . . . .	24
Exa 2.3	Wavelength of the matter wave associated with a proton	25
Exa 2.5	Uncertainty in determining the position of the electron	25
Exa 2.6	Minimum error in measurement of lifetime of excited state of hydrogen atom . . . . .	26
Exa 2.7	Uncertainty in the velocity of an electron . . . . .	26
Exa 2.8	Smallest possible uncertainty in position of an electron	27
Exa 2.9	Energy of an electron moving in 1D infinitely high potential box . . . . .	28

Exa 2.10	Lowest two permitted energy values of an electron . . .	28
Exa 2.11	Lowest energy of the neutron confined to the nucleus .	29
Exa 2.12	Energy difference between the ground state and the first excited state for an electron in 1D box . . . . .	29
Exa 3.1	Probability of existence of oxygen molecules within the given velocity range . . . . .	31
Exa 3.2	Probability that the speed of oxygen molecules . . . . .	31
Exa 3.3	Temperature to produce invariant average speed of hydrogen molecule . . . . .	32
Exa 3.4	Fraction of oxygen gas molecules within one percent of most probable speed . . . . .	33
Exa 3.5	Most probable distribution of 5 distinguishable particles among 3 cells . . . . .	34
Exa 3.6	Thermodynamic probability of the macrostate of distributing 8 distinguishable particles in 2 compartments	35
Exa 3.7	Three particles obeying Bose Einstein statistics distributed in three cells . . . . .	36
Exa 3.8	Probability of macrostate . . . . .	37
Exa 3.11	Fermi energy of the Na at absolute zero . . . . .	38
Exa 3.12	Fermi energy of silver in metallic state . . . . .	38
Exa 3.13	Fermi energy of free electrons in cesium . . . . .	39
Exa 3.15	Temperature at which the level above the fermi level is occupied by the electron . . . . .	40
Exa 4.1	Actual path of light using Fermat principle . . . . .	41
Exa 4.2	Light reflected from the inner surface of spherical shell	42
Exa 4.3	Equivalent focal length of the combinations of lenses .	42
Exa 4.4	Focal length of the combination of lenses of given powers	43
Exa 4.5	Focal length of the combination of coaxially placed thin convex lenses . . . . .	44
Exa 4.7	Locations of principal points and focal points . . . . .	44
Exa 4.8	Position of principal points and focal points for two coaxially placed lenses . . . . .	45
Exa 4.9	Focal lengths from dispersive powers of achromatic combination of lenses . . . . .	46
Exa 4.10	Focal length of the two component lens of an achromatic doublet . . . . .	47
Exa 4.11	Radii of curvature of the second surface of each of the lens of achromatic doublet . . . . .	48

Exa 4.12	Focal length of the convergent lens for C line . . . . .	50
Exa 4.13	Focal length of two lenses with no aberration . . . . .	51
Exa 4.14	Longitudinal chromatic aberration for an object at infinity . . . . .	51
Exa 4.15	Focal length of component lenses of a convergent doublet	52
Exa 4.16	Radii of Aplanatic surfaces and lateral magnification of the image . . . . .	53
Exa 4.17	Aplanatic surface . . . . .	53
Exa 4.18	Focal length of the field lens . . . . .	54
Exa 4.19	Equivalent focal length of a Ramsden eyepiece . . . . .	55
Exa 4.20	Focal lengths of the lenses and the eyepiece . . . . .	55
Exa 4.21	Focal length of the component lenses and the separation between them . . . . .	56
Exa 5.1	Order of interference maximum with different wavelength	57
Exa 5.2	Angle of the biprism . . . . .	58
Exa 5.3	Thickness of the mica sheet . . . . .	58
Exa 5.4	Distance between the two coherent sources . . . . .	59
Exa 5.5	Wavelength of light used in a biprism experiment . . . . .	59
Exa 5.6	Distance between the slits . . . . .	60
Exa 5.7	Lateral shift of central maximum . . . . .	60
Exa 5.8	Thickness of a soap bubble film . . . . .	61
Exa 5.9	Wavelength of the light used in Newton rings experiment	62
Exa 5.10	Radius of the curvature of the lens and the thickness of the air film . . . . .	62
Exa 5.11	Thickness of the soap film . . . . .	63
Exa 5.12	Least thickness of the soap film that will appear bright dark . . . . .	63
Exa 5.13	Thickness of the wire separating edges of two plane glass surfaces . . . . .	64
Exa 5.14	Radius of curvature of the lens and the thickness of the corresponding air film . . . . .	65
Exa 5.16	Angles at which first and second order maxima can be observed . . . . .	65
Exa 5.17	Relation between two wavelengths illuminating a single slit due to Fraunhofer diffraction . . . . .	66
Exa 5.18	Angular position of the first two minima on either side of a central maxima . . . . .	67



Exa 5.19	Wavelength of light and the missing order of Fraunhofer diffraction . . . . .	68
Exa 5.20	Deduction of wavelength of the light from given data . . . . .	69
Exa 5.21	Minimum number of lines in a grating . . . . .	69
Exa 5.22	Maximum number of visible orders . . . . .	70
Exa 5.23	Grating element of diffraction grating . . . . .	70
Exa 5.26	Coinciding spectral lines . . . . .	71
Exa 5.27	Resolution of D1 and D2 lines of sodium . . . . .	72
Exa 5.28	Distance between two stars which are just resolved . . . . .	73
Exa 5.29	Numerical aperture of the objective of microscope . . . . .	73
Exa 5.30	Aperture of the objective of a telescope . . . . .	74
Exa 5.31	Minimum distance from the telescope at which the the pinhole can be resolved . . . . .	75
Exa 5.32	Numerical aperture of the objective of microscope for given wavelength of light . . . . .	75
Exa 5.33	Angle of minimum deviation for green light for its passage through a prism . . . . .	76
Exa 5.34	Thickness of quarter wave plate . . . . .	76
Exa 5.35	Percentage purity of the sugar sample . . . . .	77
Exa 5.36	Specific rotation of sugar solution for given plane of polarization . . . . .	78
Exa 5.37	Angle of rotation produced by quartz plate . . . . .	78
Exa 5.38	Optical rotation produced by new length of sugar solution . . . . .	79
Exa 5.39	Strength of the solution . . . . .	80
Exa 5.40	Length of sugar solution for given concentration and optical rotation . . . . .	80
Exa 6.1	Electrons striking the target in X ray coolidge tube . . . . .	82
Exa 6.2	Maximum speed of the electron striking the target . . . . .	83
Exa 6.3	Longest wavelength that can be analysed by a rock salt crystal . . . . .	83
Exa 6.4	Angles at which the second and the third Bragg diffraction maxima are observed . . . . .	84
Exa 6.5	Interplanar separation of atomic planes in the crystal . . . . .	84
Exa 6.6	Wavelength of K alpha radiation in copper for its given value in Mo . . . . .	85
Exa 6.7	Wavelength of gamma radiation at 90 degree . . . . .	86
Exa 6.8	Compton shift from a carbon block . . . . .	86

Exa 6.9	Wavelength of incident photon . . . . .	87
Exa 7.1	Energy of the laser pulse . . . . .	88
Exa 7.2	Coherence length resultant bandwidth and line width of laser beam . . . . .	88
Exa 7.3	Angular spread and areal spread of laser beam . . . . .	89
Exa 8.1	Frequency of the fundamental mode of ultrasonic wave	91
Exa 8.2	Fundamental frequency of quartz crystal . . . . .	91
Exa 8.3	Thickness of steel plate using ultrasonic beam . . . . .	92
Exa 8.4	Inductance of an inductor to produce ultrasonic waves	93
Exa 8.5	Position of imperfection and the velocity of pulse inside the rod . . . . .	93
Exa 8.6	Maximum acceleration and displacement of a quartz ultrasonic transducer . . . . .	94
Exa 8.7	Fundamental frequency of a magnetostrictive hydrophone	95
Exa 8.8	Length of the copper wire used to introduce ultrasonic delay . . . . .	95
Exa 9.1	NA and the acceptance angle of optical fibre . . . . .	97
Exa 9.2	NA acceptance angle and the critical angle of optical fibre . . . . .	97
Exa 9.3	Acceptance angle for the optical fibre in water . . . . .	98
Exa 9.4	The characteristics of glass clad fibre . . . . .	99
Exa 9.5	Refractive index of core and cladding of an optical fibre	100
Exa 9.6	NA and the core radius of an optical fibre . . . . .	100
Exa 9.7	v number and the number of modes supported by the optical fibre . . . . .	101
Exa 9.8	Maximum values of refractive index of cladding and the fractional change in refractive index . . . . .	102
Exa 9.9	Normalized frequency for the optical fibre . . . . .	102
Exa 9.10	Cut off parameter or v number of modes supported by the fibre . . . . .	103
Exa 9.11	Power output through optical fibre . . . . .	104
Exa 9.12	Attenuation of power through optical fibre . . . . .	104
Exa 9.13	Minimum optical power input to an optical fibre . . . . .	105
Exa 10.1	Potential difference between the two charged horizontal plates . . . . .	106
Exa 10.2	Electric potential at a point equidistant from the three corners of a triangle . . . . .	107
Exa 10.3	Electric potential at the centre of a square . . . . .	107

Exa 10.6	New potential when the two charged drops coalesce to form a bigger drop . . . . .	108
Exa 10.7	Magnitude and the direction of electric field which would balance the weight of an electron placed in it . . . . .	109
Exa 10.8	Magnitude and the direction of electric field at a point midway between two charges . . . . .	109
Exa 10.9	Electric field strength at a point . . . . .	110
Exa 10.11	Electric field strength due to spherical charge distribution	111
Exa 10.12	Maximum electric field strength at an internal point .	112
Exa 11.1	Amplitude of field vector $E$ in free space . . . . .	113
Exa 11.2	Maximum value of magnetic induction vector . . . . .	113
Exa 11.3	Conduction and displacement current densities in the conducting medium . . . . .	114
Exa 11.8	Average value of the intensity of electric field of radiation	115
Exa 11.9	Amplitude of electric and magnetic fields of radiation .	116
Exa 11.10	Phase difference between electric and magnetic field vectors of an EM wave . . . . .	116
Exa 11.12	Skin depth of an EM wave in Al . . . . .	117
Exa 11.14	Skin depth and attenuation constant of sea water . . .	118
Exa 12.1	Current through the solenoid . . . . .	120
Exa 12.2	Magnetic moment of the iron rod . . . . .	120
Exa 12.3	Magnetic moment of the rod placed inside the solenoid	121
Exa 12.4	Magnetizing force and relative permeability of the material . . . . .	122
Exa 12.5	Magnetic flux density and magnetic intensity . . . . .	123
Exa 12.6	Total dipole moment of the sample . . . . .	124
Exa 12.7	Magnetization and magnetic moment in the bar . . . .	124
Exa 12.8	Hysteresis loss of energy per hour in the iron core of the transformer . . . . .	126
Exa 12.9	Hysteresis loss from BH loop . . . . .	126
Exa 12.10	Change in the magnetic dipole moment of the electron	127
Exa 13.1	Electric dipole placed in a uniform electric field . . . .	128
Exa 13.2	Force acting on an electric dipole in different orientations relative to the electric field . . . . .	129
Exa 13.3	Dielectric constant and the electric permittivity of the material . . . . .	130
Exa 13.4	Dielectric constant and the electric susceptibility of diamond . . . . .	130

Exa 13.5	Calculate the values of E D and P . . . . .	131
Exa 13.6	Dipole moment induced in He atom . . . . .	132
Exa 13.7	Induced dipole moment and atomic polarizability of neon gas . . . . .	132
Exa 13.8	Electronic polarizability of argon atom . . . . .	133
Exa 13.9	Individual dipole moment of carbon tetrachloride . . .	134
Exa 13.10	Atomic radius of He . . . . .	134
Exa 13.11	Percentage of ionic polarizability in NaCl crystal . . .	135
Exa 13.12	Determine the dipole moment . . . . .	135
Exa 14.1	Density of impurity atoms to N type and P type silicon	137
Exa 14.2	Electrical conductivity and resistivity of intrinsic germanium sample . . . . .	138
Exa 14.3	Electrical conductivity of undoped and doped silicon .	139
Exa 14.4	Shift in Fermi level due to change in density of donor atoms . . . . .	140
Exa 14.5	Voltage required to cause a forward current density in pn junction diode . . . . .	140
Exa 14.6	Applied voltage for forward current density . . . . .	141
Exa 14.7	Forward voltage to increase the current density of Si diode . . . . .	142
Exa 14.8	Static and dynamic values of diode resistance . . . . .	143
Exa 14.9	Half wave rectifier parameters . . . . .	143
Exa 14.10	Full wave rectifier parameters . . . . .	144
Exa 14.11	Current gains in BJT . . . . .	145
Exa 14.12	Base current of BJT in CB mode . . . . .	145
Exa 14.13	Current gain and base current of BJT in CB mode . .	146
Exa 14.14	Current gain and leakage current of BJT in CE mode	147
Exa 14.15	Voltage and power gain of PNP transistor in CB mode	147
Exa 15.1	Binary equivalent of decimal number . . . . .	149
Exa 15.2	Decimal equivalent of 6 bit binary number . . . . .	150
Exa 15.3	Decimal equivalent of octal number . . . . .	151
Exa 15.4	Octal equivalent of decimal number . . . . .	151
Exa 15.5	Hexadecimal equivalent of binary numbers . . . . .	152
Exa 15.6	Hexadecimal equivalent of decimal numbers . . . . .	154
Exa 15.7	Addition of two binary numbers . . . . .	155
Exa 15.8	Addition of two binary numbers with fractions . . . .	156
Exa 15.9	Subtraction of two binary numbers . . . . .	158
Exa 15.10	Multiplication of two binary numbers . . . . .	159

Exa 15.11	Binary division of two numbers . . . . .	160
Exa 16.1	Miller indices of the crystal plane . . . . .	162
Exa 16.2	Miller indices of the lattice plane . . . . .	163
Exa 16.3	Miller indices of the set of parallel planes . . . . .	163
Exa 16.4	Length of the intercepts on Y and Z axes . . . . .	164
Exa 16.5	Lattice constant for NaCl crystal . . . . .	165
Exa 16.6	Lattice constant for KBr crystal . . . . .	165
Exa 16.7	Lattice constant for Cu and distance between the two nearest Cu atoms . . . . .	166
Exa 16.8	Inter planar spacing for lattice planes . . . . .	167
Exa 16.9	Interplanar spacing in cubic crystal . . . . .	168
Exa 16.10	Interplanar spacing in tetragonal crystal lattice . . . . .	168
Exa 17.1	Binding energy per nucleon for the deuteron . . . . .	170
Exa 17.2	Binding energy for an alpha particle . . . . .	171
Exa 17.3	Weizsacker formula for stability of nuclei . . . . .	171
Exa 17.4	Nuclei stability . . . . .	172
Exa 17.5	Exothermicity and endothermicity of nuclear reactions	173
Exa 17.6	Q value of the formation of P30 in ground state . . . . .	175
Exa 17.7	Threshold energy required to initiate the reaction . . . . .	176
Exa 17.8	Kinetic energy of emitted protons and threshold energy	178
Exa 17.9	Energy released by the fission of 1 kg of U235 . . . . .	179
Exa 17.10	Rate of fission of U235 . . . . .	180
Exa 17.11	Rate of fission and energy released in the complete fis- sioning of U235 . . . . .	180
Exa 17.12	Energy and oscillator frequency of some positively charged particles accelerating in a cyclotron . . . . .	181
Exa 17.13	Energy of the protons issuing out of the cyclotron . . . . .	182
Exa 17.14	Energy gained per turn and maximum energy of the electron in a betatron . . . . .	183
Exa 17.15	Maximum frequency of Dee voltage and gain in energy of the deuteron in a synchrocyclotron . . . . .	184
Exa 17.16	Maximum radial field and the life of a GM counter . . . . .	185
Exa 17.17	Avalanche voltage in a GM tube . . . . .	185
Exa 17.18	Maximum permissible voltage fluctuation in a GM counter	186
Exa 19.1	Critical field for lead at 4 K . . . . .	188
Exa 19.2	Isotopic effect in mercury . . . . .	188
Exa 19.3	Critical current through superconducting aluminium wire	189

Exa 19.4	Critical current density for superconducting wire of lead at different temperatures . . . . .	190
Exa 19.5	Critical temperature of lead from London penetration depth . . . . .	191
Exa 19.6	Field strength required by lead to lose its superconducting state at 0 K . . . . .	191
Exa 19.7	Critical temperature for niobium . . . . .	192

# Chapter 1

## Relativistic Mechanics

Scilab code Exa 1.2 Lorentz transformations for space and time

```
1 // Scilab Code Ex1.2: Page:26 (2011)
2 clc; clear;
3 c = 3e+008; // Speed of light in vacuum, m/s
4 delta_x = 2.45e+03; // Space difference, m
5 delta_t = 5.35e-06; // Time difference, s
6 v = 0.855*c; // Speed of frame S_prime, m/s
7 delta_x_prime = 1/sqrt(1-v^2/c^2)*(delta_x - v*(
    delta_t))*1e-03; // Distance between two
    flashes as measured in S_prime frame, km
8 delta_t_prime = 1/sqrt(1-v^2/c^2)*(delta_t - v/c^2*
    delta_x)*1e+006; // Time between two flashes
    as measured in S_prime
9 printf("\\nThe distance between two flashes as
    measured in S_prime frame = %4.2f km",
    delta_x_prime);
10 printf("\\nThe time between two flashes as measured
    in S_prime frame = %4.2f micro-second",
    delta_t_prime);
11
12 // Result
13 // The distance between two flashes as measured in
```

```

S_prime frame = 2.08 km
14 // The time between two flashes as measured in
    S_prime frame = -3.15 micro-second

```

---

**Scilab code Exa 1.4** Relative speed of one photon with respect to another

```

1 // Scilab Code Ex1.4: Page:27 (2011)
2 clc;clear;
3 c = 1;....// Speed of light in vacuum, m/s
4 u_x_prime = c; // Velocity of photon as measured
    in S_prime frame, m/s
5 v = c; // Velocity of frame S_prime relative to S
    frame, m/s
6 u_x = (u_x_prime + v)/(1+v*u_x_prime/c^2);
7 if u_x == 1 then
8     ux = 'c';
9 else
10    ux = string(u_x)+'c';
11 end
12 printf("\nThe speed of one photon as observed by the
    other is %c", ux);
13
14 // Result
15 // The speed of one photon as observed by the other
    is c

```

---

**Scilab code Exa 1.6** Areal contraction of moving circular lamina

```

1 // Scilab Code Ex1.6 : Page:28 (2011)
2 clc;clear;
3 a = 1; // For simplicity assume length of semi
    minor axis to be unity, m
4 c = 3e+08; // Speed of light, m/s

```



```

5 v = poly(0, 'v'); // Declare velocity variable , m
  /s
6 // As b = a*sqrt(1-v^2/c^2), length of semi-major
  axis
7 // Also A_c = %pi*a^2, area of the lamina in its own
  frame and
8 // A_e = %pi*a*b, area of the lamina in stationary
  frame S, so with A_c = A_e
9 v = roots(1-v^2/c^2 - 1/4); // Velocity at which
  surface area of lamina reduces to half in S-frame
  , m/s
10 printf("\nThe velocity at which surface area of
  lamina reduces to half in S-frame = %4.2e", v(1))
  ;
11
12 // Result
13 // The velocity at which surface area of lamina
  reduces to half in S-frame = 2.60e+008

```

---

**Scilab code Exa 1.7** Length of a one metre stick moving parallel to its length

```

1 // Scilab Code Ex1.7 : Page:29 (2011)
2 clc;clear;
3 m0 = 1; // For simplicity assume the rest mass of
  stick to be unity, kg
4 m = 1.5*m0; // Mass of the moving stick, kg
5 L0 = 1; // Assume resting length of the stick to
  be unity, m
6 // As m = m0/sqrt(1-v^2/c^2) = m0*gama, solving for
  gama
7 gama = m/m0; // Relativistic factor
8 L = L0/gama; // Contracted length of the metre
  stick, m
9 printf("\nThe contracted length of the metre stick =

```

```

    %4.2 f m", L);
10
11 // Result
12 // The contracted length of the metre stick = 0.67 m

```

---

**Scilab code Exa 1.8** Mean lifetime of meson in motion

```

1 // Scilab Code Ex1.8: Page:29 (2011)
2 clc;clear;
3 c = 3e+008; // Speed of light in vacuum, m/s
4 tau0 = 2e-008; // Mean lifetime of meson at rest,
    m/s
5 v = 0.8*c; // Velocity of moving meson, m/s
6 tau = tau0/sqrt(1-v^2/c^2); // Mean lifetime of
    meson in motion, m/s
7 printf("\nThe mean lifetime of meson in motion = %4
    .2e s", tau);
8
9 // Result
10 // The mean lifetime of meson in motion = 3.33e-008
    s

```

---

**Scilab code Exa 1.9** Speed at which a moving clock ticks slow

```

1 // Scilab Code Ex1.9: Page:30 (2011)
2 clc;clear;
3 c = 3e+008; // Speed of light in vacuum, m/s
4 delta_t0 = 59; // Reading of the moving clock for
    each hour, min
5 delta_t = 60; // Reading of the stationary clock
    for each hour, min
6 // As from Time Dilation, delta_t = delta_t0/sqrt(1-
    v^2/c^2), solving for v

```

```

7 v = c*sqrt(1-(delta_t0/delta_t)^2);
8 printf("\nThe speed at which the moving clock ticks
    slow = %4.2e m/s", v);
9
10 // Result
11 // The speed at which the moving clock ticks slow =
    5.45e+007 m/s

```

---

**Scilab code Exa 1.10** Distance that the meson beam can travel before reduction in its flux

```

1 // Scilab Code Ex1.10: Page:30 (2011)
2 clc;clear;
3 c = 3e+008; // Speed of light in vacuum, m/s
4 tau0 = 2.5e-008; // Mean lifetime of meson at
    rest, m/s
5 v = 0.8*c; // Velocity of moving meson, m/s
6 tau = tau0/sqrt(1-v^2/c^2); // Mean lifetime of
    meson in motion, m/s
7 N0 = 1; // Assume initial flux of meson beam to
    be unity, watt/Sq.m
8 N = N0*%e^(-2); // Meson flux after time t, watt/
    Sq.m
9 // As N = N0*e^(-t/tau), which on comparing gives
10 t = 2*tau; // Time during which the meson beam
    flux reduces, s
11 d = 0.8*c*t; // The distance that the meson beam
    can travel before reduction in its flux, m
12 printf("\nThe distance that the meson beam can
    travel before reduction in its flux = %2d m", d);
13
14 // Result
15 // The distance that the meson beam can travel
    before reduction in its flux = 20 m

```

---

**Scilab code Exa 1.11** Velocity of the particle when its total energy is thrice its rest energy

```
1 // Scilab Code Ex1.11: Page:31 (2011)
2 clc;clear;
3 c = 3e+008; // Speed of light in vacuum, m/s
4 E0 = 1; // Rest energy of particle, unit
5 E = 3*E0; // Energy of relativistically moving
   particle, unit
6 // E = m*c^2 and E0 = m0*c^2
7 // With m = m0/sqrt(1-v^2/c^2), we have
8 v = c*sqrt(1-(E0/E)^2); // Velocity of the moving
   particle, m/s
9 printf("\nThe velocity of the moving particle = %4.2
   e m/s", v);
10
11 // Result
12 // The velocity of the moving particle = 2.83e+008 m
   /s
```

---

**Scilab code Exa 1.12** Kinetic energy and momentum of moving electron

```
1 // Scilab Code Ex1.12: Page:32 (2011)
2 clc;clear;
3 c = 3e+008; // Speed of light in vacuum, m/s
4 m0 = 9.1e-031; // Rest mass of electron, kg
5 m = 11*m0; // Mass of relativistically moving
   electron, kg
6 E_k = (m-m0)*c^2/(1.6e-019*1e+06); // Kinetic
   energy of moving electron, MeV
7 // As m = m0/sqrt(1-v^2/c^2), solving for v
```

```

8 v = c*sqrt(1-(m0/m)^2);    // The velocity of the
    moving electron , m/s
9 p = m*v;    // Momentum of moving electron , kg-m/s
10 printf("\nThe kinetic energy of moving electron = %4
    .2 f MeV", E_k);
11 printf("\nThe momentum of moving electron = %4.2e kg
    -m/s", p);
12
13 // Result
14 // The kinetic energy of moving electron = 5.12 MeV
15 // The momentum of moving electron = 2.99e-021 kg-m/
    s

```

---

**Scilab code Exa 1.13** Amount of work to be done to increase the speed of an electron

```

1 // Scilab Code Ex1.13: Page:32 (2011)
2 clc;clear;
3 c = 3e+008;    // Speed of light in vacuum, m/s
4 E0 = 0.5;    // Rest energy of the electron, MeV
5 v1 = 0.6*c;    // Initial velocity of the electron ,
    m/s
6 v2 = 0.8*c;    // Final velocity of the electron , m/
    s
7 W = (1/sqrt(1-v2^2/c^2)-1/sqrt(1-v1^2/c^2))*E0;
    // The amount of work to be done to increase the
    speed of the electron , MeV
8 printf("\nThe amount of work to be done to increase
    the speed of an electron = %4.2e J", W*1e+06*1.6e
    -019);
9
10 // Result
11 // The amount of work to be done to increase the
    speed of an electron = 3.33e-014 J

```

---

### Scilab code Exa 1.14 Particle moving with relativistic speed

```
1 // Scilab Code Ex1.14: Page:33 (2011)
2 clc; clear;
3 c = 1; // Assume speed of light in vacuum to be
    unity, unit
4 m0 = 1; // For simplicity assume rest mass of the
    particle to be unity, unit
5 v = c/sqrt(2); // Given speed of the particle, m/
    s
6 gama = 1/sqrt(1-v^2/c^2); // Relativistic factor
7 m = gama*m0; // The relativistic mass of the
    particle, unit
8 p = m*v; // The relativistic momentum of the
    particle, unit
9 E = m*c^2; // The relativistic total energy of
    the particle, unit
10 E_k = (m-m0)*c^2; // The relativistic kinetic
    energy of the particle, unit
11 printf("\\nThe relativistic mass of the particle = %5
    .3fm0", m);
12 printf("\\nThe relativistic momentum of the particle
    = %1.0gm0c", p);
13 printf("\\nThe relativistic total energy of the
    particle = %5.3fm0c^2", E);
14 printf("\\nThe relativistic kinetic energy of the
    particle = %5.3fm0c^2", E_k);
15
16 // Result
17 // The relativistic mass of the particle = 1.414m0
18 // The relativistic momentum of the particle = 1m0c
19 // The relativistic total energy of the particle =
    1.414m0c^2
20 // The relativistic kinetic energy of the particle =
```

$$0.414m_0c^2$$

---

**Scilab code Exa 1.15** Speed of the electron in order to have its mass equal to mass of a proton

```
1 // Scilab Code Ex1.15: Page:34 (2011)
2 clc;clear;
3 c = 3e+008; // Speed of light in vacuum, unit
4 m0 = 9.1e-031; // Rest mass of the electron, kg
5 m = 1.67e-027; // Rest mass of the proton, kg
6 // As  $m = m_0/\sqrt{1-v^2/c^2}$ , solving for v
7 v = c*sqrt(1-(m0/m)^2); // Velocity of the
   electron, m/s
8 printf("\nThe velocity of the electron to have its
   mass equal to mass of the proton = %5.3e m/s", v)
9
10 // Result
11 // The velocity of the electron to have its mass
   equal to mass of the proton = 3.000e+008 m/s
```

---

**Scilab code Exa 1.17** Classical and relativistic speed of an electron of given kinetic energy

```
1 // Scilab Code Ex1.17: Page:35 (2011)
2 clc;clear;
3 c = 3e+008; // Speed of light in vacuum, unit
4 m0 = 9.1e-031; // Rest mass of the electron, kg
5 E_k = 0.1*1e+006*1.6e-019; // Kinetic energy of
   the electron, J
6 v = sqrt(2*E_k/m0); // Classical speed of the
   electron, m/s
```

```

7 printf("\nThe classical speed of the electron = %5.3
  e m/s", v);
8 // As  $E_k = (m-m_0)*c^2 = (1/\sqrt{1-v^2/c^2}-1)*m_0*c^2$ , solving for v
9 v = c*sqrt(1-(m0*c^2/(E_k+m0*c^2))^2); //
  Relativistic speed of the electron, m/s
10 printf("\nThe relativistic speed of the electron =
  %5.3e m/s", v);
11
12 // Result
13 // The classical speed of the electron = 1.875e+008
  m/s
14 // The relativistic speed of the electron = 1.644e
  +008 m/s

```

---



## Chapter 2

# Quantum Mechanics

Scilab code Exa 2.1 de broglie wavelength of an electron

```
1 // Scilab Code Ex2.1: Page:79 (2011)
2 clc;clear;
3 V = 50;....// Given potential difference , V
4 lambda = 12.24/sqrt(V);....// Wavelength of the
   light , angstrom
5 printf("\nThe de-broglie wavelength of electron = %4
   .2f angstrom", lambda);
6
7 // Result
8 // The de-broglie wavelength of electron = 1.73
   angstrom
```

---

Scilab code Exa 2.2 de broglie wavelength associated with a proton

```
1 // Scilab Code Ex2.2: Page:79 (2011)
2 clc;clear;
3 h = 6.62e-34; // Planck's constant , J-s
4 m0 = 1.6e-27; // Rest mass of proton , kg
```

```

5 c = 3e+8;    // Speed of light , in m/s
6 v = c/20;    // Velocity of the proton , in m/s
7 lambda = (h*sqrt(1-v^2/c^2))/(m0*v);
8 printf("\nThe de broglie wavelength associated with
    the proton = %4.2e m",lambda);
9
10 // Result
11 // The de broglie wavelength associated with the
    proton = 2.75e-14 m

```

---

**Scilab code Exa 2.3** Wavelength of the matter wave associated with a proton

```

1 // Scilab Code Ex2.3: Page:79 (2011)
2 clc;clear;
3 c = 3e+8;....// Speed of light , m/s
4 v = 2e+8;....// Velocity of the proton , m/s
5 m0 = 1.6e-27;....// Rest mass of proton , kg
6 h = 6.62e-34;....// Plancks constant ,J-s
7 lambda = (h*sqrt(1-v^2/c^2))/(m0*v);
8 printf("\nThe wavelength of matter wave associated
    with the proton = %5.3e m", lambda);
9
10 // Result
11 // The wavelength of matter wave associated with
    the proton = 1.542e-15 m

```

---

**Scilab code Exa 2.5** Uncertainty in determining the position of the electron

```

1 // Scilab Code Ex2.5: Page:80 (2011)
2 clc;clear;
3 a = 0.003;....// Accuracy of the electron ,in percent

```

```

4 s = 5e+03;....// Speed of the electron ,in m/s
5 del_v = (a/100)*s;....// Change in velocity ,in m/s
6 m0 = 9.1e-31;....// Rest mass of the electron ,in kg
7 hcut = 1.054e-34;....// Plancks constant ,J-s
8 del_x = hcut/(2*del_v*m0);
9 printf("\nThe uncertainty in the position of the
    electron = %4.2e m", del_x);
10
11 // Result
12 // The uncertainty in the position of the electron
    = 3.86e-004 m

```

---

**Scilab code Exa 2.6** Minimum error in measurement of lifetime of excited state of hydrogen atom

```

1 // Scilab Code Ex2.6 : Page:81 (2011)
2 clc;clear;
3 del_t = 2.5e-14;....// Lifetime of the hydrogen atom
    in excited state
4 hcut = 1.054e-34;....// Planck's constant ,in J-s
5 e = 1.6e-19;....// Charge on electron ,in C
6 del_E = hcut/(2*del_t*e);....// Energy of the state ,
    in eV
7 printf("\nThe minimum error in measurement of
    lifetime of excited state of hydrogen atom = %6.4
    f eV",del_E);
8
9 // Result
10 // The minimum error in measurement of lifetime of
    excited state of hydrogen atom = 0.0132 eV

```

---

**Scilab code Exa 2.7** Uncertainty in the velocity of an electron

```

1 // Scilab Code Ex2.7 : Page:81 (2011)
2 clc;clear;
3 del_x = 1e-09; // Uncertainty in position of the
    electron , m
4 m0 = 9.1e-031;....// Rest mass of an electron , kg
5 hcut = 1.054e-034;....// Planck 's constant ,in J-s
6 del_v = hcut/(2*del_x*m0);....// Uncertainty in
    velocity of the electron
7 printf("\\nThe uncertainty in the velocity of an
    electron = %4.2e m/s",del_v);
8
9 // Result
10 // The uncertainty in the velocity of an electron =
    5.79e+04 m/s

```

---

**Scilab code Exa 2.8** Smallest possible uncertainty in position of an electron

```

1 // Scilab Code Ex2.8 : Page:81 (2011)
2 clc;clear;
3 hcut = 1.054e-34; // Reduced Planck 's constant , Js
4 v = 3e+07;....// Velocity of the electron , m/s
5 c = 3e+08;....// Speed of light in vacuum, m/s
6 m0 = 9.1e-31;....// Rest mass of an electron , kg
7 del_v = 3e+08;....// Uncertainty in velocity of the
    electron , m/s
8 del_x = (hcut*sqrt(1-v^2/c^2))/(2*m0*del_v);
9 printf("\\nThe smallest possible uncertainty in
    position of the electron = %6.4f angstrom", del_x
    /1e-010);
10
11 // Result
12 // The smallest possible uncertainty in position of
    the electron = 0.0019 angstrom

```

---

**Scilab code Exa 2.9** Energy of an electron moving in 1D infinitely high potential box

```
1 //Scilab Code Ex2.9 :Page:82 (2011)
2 clc;clear;
3 n = 1;
4 m0 = 9.1e-31;....// Mass of the electron , kg
5 a = 1e-10;....// Width of the box, m
6 h = 6.63e-34;....// Planck's constant , J-s
7 E = n^2*h^2/(8*m0*a^2);
8 printf("\n The energy of the electron moving in 1D
   infinitely high potential box = %5.2e J", E);
9
10 // Result
11 // The energy of the electron moving in 1D
   infinitely high potential box = 6.04e-18 J
```

---

**Scilab code Exa 2.10** Lowest two permitted energy values of an electron

```
1 // Scilab Code Ex2.10: Page:83 (2011)
2 clc;clear;
3 n = [1,2];....// Shell numbers for two lowest
   permitted energy of the electron
4 m0 = 9.1e-31;....// Mass of the electron , kg
5 a = 2.5e-10;....// Width of the box, m
6 h = 6.63e-34;....// Planck's constant , J-s
7 e = 1.6e-19;....// Charge on electron , C
8 E = (n^2*h^2)/(8*m0*a^2*e);
9 printf("\nThe lowest two permitted energy values of
   an electron are");
10 printf(" %d eV and %d eV respectively", E(1), E(2));
11
```

```

12 // Result
13 // The lowest two permitted energy values of an
    electron are 6 eV and 24 eV respectively

```

---

**Scilab code Exa 2.11** Lowest energy of the neutron confined to the nucleus

```

1 // Scilab Code Ex2.11: Page:83 (2011)
2 clc;clear;
3 m0 = 1.67e-27;....// Rest mass, in kg
4 a = 1e-14;....// Size of the box
5 h = 6.63e-34;....// Planck's constant, in J-s
6 n = 1; // Quantum number for lowest energy state
7 E_n = n^2*h^2/(8*m0*a^2);
8 printf("\\nThe lowest energy of the neutron confined
    to the nucleus = %4.2e J", E_n);
9
10 // Result
11 // The lowest energy of the neutron confined to the
    nucleus = 3.29e-13 J

```

---

**Scilab code Exa 2.12** Energy difference between the ground state and the first excited state for an electron in 1D box

```

1 // Scilab Code Ex2.12: Page:83 (2011)
2 clc;clear;
3 m0 = 9.1e-31;....// Rest mass, kg
4 a = 1e-10;....// Length of the box, m
5 h = 6.62e-34;....// Planck's constant, J-s
6 n1 = 1;....// Ground state
7 n2 = 2;....// First excited state
8 e = 1.6e-19;....// Charge on electron, C
9 E1 = (n1^2*h^2)/(8*m0*a^2*e);

```

```
10 E2 = (n2^2*h^2)/(8*m0*a^2*e);
11 del_E = E2-E1;
12 printf("\nThe energy difference between the ground
    state and the first excited state = %5.1f eV",
    del_E);
13
14 //Result
15 // The energy difference between the ground state
    and the first excited state = 112.9 eV
```

---

# Chapter 3

## Statistical Mechanics

**Scilab code Exa 3.1** Probability of existence of oxygen molecules within the given velocity range

```
1 // Scilab Code Ex3.1: Page:132 (2011)
2 clc;clear;
3 m = 5.32e-26; // Mass of one oxygen molecule , kg
4 k_B = 1.38e-23; // Boltzmann constant , J/K
5 T = 200; // Temperature of the system , K
6 v = 100; // Speed of the oxygen molecules , m/s
7 dv = 1; // Increase in speed of the oxygen
  molecules , m/s
8 P = 4*pi*(m/(2*pi*k_B*T))^(3/2)*exp(-m*v^2/(2*k_B*
  T))*v^2*dv;
9 printf("\nThe probability that the speed of oxygen
  molecule is %4.2e", P) ;
10
11 // Result
12 // The probability that the speed of oxygen molecule
  is 6.13e-04
```

---

**Scilab code Exa 3.2** Probability that the speed of oxygen molecules



```

1 // Scilab Code Ex3.2 : Page:132 (2011)
2 clc;clear;
3 A = 32; // Gram atomic mass of oxygen, g/mol
4 N_A = 6.023e+026; // Avogadro's number, per kmol
5 m = A/N_A;....//mass of the molecule, kg
6 k_B = 1.38e-23;....// Boltzmann constant, J/K
7 T = 273;....// Temperature of the gas, K
8 v_av = 1.59*sqrt(k_B*T/m);....// Average speed of
   oxygen molecule, m/s
9 printf("\nThe average speed of oxygen molecule is =
   %3d m/s", v_av);
10 v_rms = 1.73*sqrt(k_B*T/m);....// The mean square
   speed of oxygen molecule, m/s
11 printf("\nThe root mean square speed of oxygen gas
   molecule is = %3d m/s", ceil(v_rms))
12 v_mp = 1.41*sqrt(k_B*T/m);....// The most probable
   speed of oxygen molecule, m/s
13 printf("\nThe most probable speed of oxygen molecule
   is = %3d m/s", ceil(v_mp));
14
15 // Result
16 // The average speed of oxygen molecule is = 423 m/s
17 // The root mean square speed of oxygen gas molecule
   is = 461 m/s
18 // The most probable speed of oxygen molecule is =
   376 m/s

```

---

**Scilab code Exa 3.3** Temperature to produce invariant average speed of hydrogen molecule

```

1 // Scilab Code Ex3.3: Page:133 (2011)
2 clc;clear;
3 m_H = 2; // Gram molecular mass of hydrogen, g
4 m_O = 32; // Gram molecular mass of oxygen, g
5 k_B = 1.38e-23;....// Boltzmann constant, J/K

```

```

6 v_av0 = 1;....// For simplicity average speed of
  oxygen gas molecule is assumed to be unity, m/s
7 v_avH = 2*v_av0;....// The average speed of
  hydrogen gas molecule, m/s
8 T_0 = 300; // Temperature of oxygen gas, K
9 // As v_avO/v_av_H = sqrt(T_O/T_H)*sqrt(m_H/m_O),
  solving for T_H
10 T_H = (v_avH/v_av0*sqrt(m_H/m_0)*sqrt(T_0))^2; //
  Temperature at which the average speed of
  hydrogen gas molecules is the same as that of
  oxygen gas molecules, K
11 printf("\nTemperature at which the average speed of
  hydrogen gas molecules is the same as that of
  oxygen gas molecules at 300 K = %2d", T_H);
12
13 // Result
14 // Temperature at which the average speed of
  hydrogen gas molecules is the same as that of
  oxygen gas molecules at 300 K = 75

```

---

**Scilab code Exa 3.4** Fraction of oxygen gas molecules within one percent of most probable speed

```

1 // Scilab Code Ex3.4: Page:133 (2011)
2 clc;clear;
3 v_mp = 1; // Most probable speed of gas molecules,
  m/s
4 dv = 1.01*v_mp-0.99*v_mp; // Change in most
  probable speed, m/s
5 v = v_mp; // Speed of the gas molecules, m/s
6 Frac = 4/sqrt(%pi)*1/v_mp^3*exp(-v^2/v_mp^2)*v^2*dv;
7 printf("\nThe fraction of oxygen gas molecules
  within one percent of most probable speed = %5.3 f
  ", Frac);
8

```

```

9 // Result
10 // The fraction of oxygen gas molecules within one
    percent of most probable speed = 0.017

```

---

**Scilab code Exa 3.5** Most probable distribution of 5 distinguishable particles among 3 cells

```

1 // Scilab Code Ex3.5: Page:134 (2011)
2 clc;clear;
3 n = 5; // Number of distinguishable particles which
    are to be distributed among cells
4 n1 = [5 4 3 3 2]; // Possible occupancy of
    particles in first cell
5 n2 = [0 1 2 1 2]; // Possible occupancy of
    particles in second cell
6 n3 = [0 0 0 1 1]; // Possible occupancy of
    particles in third cell
7 BIG_W = 0;
8 printf("\n-----");
9 printf("\nn1      n2      n3      5/(n1!n2!n3!)");
10 printf("\n-----");
11 for i = 1:1:5
12 W = factorial(n)/(factorial(n1(i))*factorial(n2(i))*
    factorial(n3(i)));
13 if BIG_W < W then
14     BIG_W = W;
15     ms = [n1(i) n2(i) n3(i)];
16 end
17 printf("\n%d      %d      %d      %d", n1(i), n2
    (i), n3(i), W);
18 end
19 printf("\n-----");
20 printf("\nThe macrostates of most probable
    distribution with thermodynamic probability %d
    are:", BIG_W);

```

```

21 printf("\n(%d, %d, %d), (%d, %d, %d) and (%d, %d, %d
    )", ms(1), ms(2), ms(3), ms(2), ms(3), ms(1), ms
    (3), ms(1), ms(2));
22
23 // Result
24 // -----
25 // n1      n2      n3      5/(n1!n2!n3!)
26 // -----
27 // 5      0      0      1
28 // 4      1      0      5
29 // 3      2      0      10
30 // 3      1      1      20
31 // 2      2      1      30
32 // -----
33 // The macrostates of most probable distribution
    with thermodynamic probability 30 are:
34 // (2, 2, 1), (2, 1, 2) and (1, 2, 2)

```

---

**Scilab code Exa 3.6** Thermodynamic probability of the macrostate of distributing 8 distinguishable particles in 2 compartments

```

1 // Scilab Code Ex3.6: Page:135 (2011)
2 clc;clear;
3 g1 = 4; // Intrinsic probability of first cell
4 g2 = 2; // Intrinsic probability of second cell
5 k = 2; // Number of cells
6 n = 8; // Number of distinguishable particles
7 n1 = 8; // Number of cells in first compartment
8 n2 = n - n1; // Number of cells in second
    compartment
9 W = factorial(n)*1/factorial(n1)*1/factorial(n2)*(g1
    )^n1*(g2)^n2;
10 printf("\nThe thermodynamic probability of the
    macrostate (8,0) = %5d", W);
11

```

```

12 // Result
13 // The thermodynamic probability of the macrostate
    (8,0) = 65536

```

---

**Scilab code Exa 3.7** Three particles obeying Bose Einstein statistics distributed in three cells

```

1 // Scilab Code Ex3.7 : Page:135 (2011)
2 clc;clear;
3 function str = st(val)
4     str = emptystr();
5     if val == 3 then
6         str = 'aaa';
7     elseif val == 2 then
8         str = 'aa';
9     elseif val == 1 then
10        str = 'a';
11    elseif val == 0 then
12        str = '0';
13    end
14 endfunction
15
16 g = 3; // Number of cells in first compartment
17 n = 3; // Number of bosons
18 p = 3;
19 r = 1; // Index for number of rows
20 clc;
21 printf("\nAll possible meaningful arrangements of
    three particles in three cells are:")
22 printf("\n-----");
23 printf("\nCell 1      Cell 2      Cell 3");
24 printf("\n-----");
25 for i = 0:1:g
26     for j = 0:1:n
27         for k = 0:1:p

```

```

28         if (i+j+k == 3) then
29             printf("\n%4s      %4s      %4s", st(i), st(j),
30                 st(k));
31         end
32     end
33 end
34 printf("\n -----");
35
36 // Result
37 // All possible meaningful arrangements of three
38 // particles in three cells are:
39 // -----
40 // Cell 1      Cell 2      Cell 3
41 // -----
42 // 0          0          aaa
43 // 0          a          aa
44 // 0          aa         a
45 // 0          aaa        0
46 // a          0          aa
47 // a          a          a
48 // a          aa         0
49 // aa         0          a
50 // aa         a          0
51 // aaa        0          0
52 // -----

```

---

**Scilab code Exa 3.8** Probability of macrostate

```

1 // Scilab Code Ex3.8 : Page:136 (2011)
2 clc;clear;
3 g1 = 3; // Number of cells in first compartment
4 g2 = 4; // Number of cells in second compartment
5 k = 2; // Number of compartments
6 n1 = 5; // Number of bosons

```

```

7 n2 = 0; // Number of with no bosons
8 W_50 = factorial(g1+n1-1)*factorial(g2+n2-1)/(
    factorial(n1)*factorial(g1-1)*factorial(n2)*
    factorial(g2-1));
9 printf("\nThe probability for the macrostate (5,0)
    is = %2d", W_50);
10
11 // Result
12 // The probability for the macrostate (5,0) is = 21

```

---

**Scilab code Exa 3.11** Fermi energy of the Na at absolute zero

```

1 // Scilab Code Ex3.11: Page:138 (2011)
2 clc;clear;
3 r = 1.86e-10;....// Radius of Na, angstrom
4 m = 9.1e-31;....// Mass of electron, in kg
5 h = 6.62e-34;....// Planck's constant, J-s
6 N = 2;....// Number of free electrons in a unit cell
    of Na
7 a = 4*r/sqrt(3);....// Volume of Na, m
8 V = a^3;....// Volume of the unit cell of Na, meter
    cube
9 E = h^2/(2*m)*(3*N/(8*pi*V))^(2/3);
10 printf("\nThe fermi energy of the Na at absolute
    zero is = %4.2e J", E);
11
12 // Result
13 // The fermi energy of the Na at absolute zero =
    5.02e-019 J

```

---

**Scilab code Exa 3.12** Fermi energy of silver in metallic state

```

1 // Scilab Code Ex3.12:Page-139 (2011)

```

```

2 clc;clear;
3 m = 9.1e-31;....// mass of electron , kg
4 h = 6.62e-34;....// Planck's constant , J-s
5 V = 108/10.5*1e-06;....// Volume of 1 gm mole of
   silver , metre-cube
6 N = 6.023e+023;    // Avogadro's number
7 E_F = h^2/(2*m)*(3*N/(8*pi*V))^(2/3);    // Fermi
   energy at absolute zero , J
8 printf("\\nThe fermi energy of the silver at absolute
   zero = %4.2e J",E_F);
9
10 // Result
11 // The fermi energy of the silver at absolute zero =
   8.80e-019 J

```

---

**Scilab code Exa 3.13** Fermi energy of free electrons in cesium

```

1 // Scilab Code Ex3.14: Electron density in lithium
   at absolute zero: Page:140 (2011)
2 clc;clear;
3 e = 1.6e-019;    // Energy equivalent of 1 eV, J/eV
4 m = 9.1e-31;....// Mass of the electron , kg
5 h = 6.63e-34;    // Planck's constant , Js
6 EF = 4.72*e;....// Fermi energy of free electrons in
   Li, J
7 rho = 8*pi/3*(2*m*EF/h^2)^(3/2);    // Electron
   density at absolute zero, electrons/metre-cube
8 printf("\\nThe electron density in lithium at
   absolute zero = %4.2e electrons/metre-cube", rho)
   ;
9
10 // Result
11 // The electron density in lithium at absolute zero
   = 4.63e+028 electrons/metre-cube

```

---



**Scilab code Exa 3.15** Temperature at which the level above the fermi level is occupied by the electron

```
1 // Scilab Code Ex3.15: Page:140 (2011)
2 clc; clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 k_B = 1.38e-023; // Boltzmann constant, J/K
5 f_E = 0.01;....// Probability that a state with
    energy 0.5 eV above the Fermi energy is occupied
    by an electron, eV
6 delta_E = 0.5; // Energy difference (E-Ef) of
    fermi energy, eV
7 // Since f_E = 1/(exp((E-Ef)/(k_B*T))+1), solvinf
    for T
8 T = delta_E/(log((1-f_E)/f_E)*k_B/e); //
    Temperature at which the level above the fermi
    level is occupied by the electron, K
9
10 printf("\\nThe temperature at which the level above
    the fermi level is occupied by the electron = %4d
    K", ceil(T));
11
12 // Result
13 // The temperature at which the level above the
    fermi level is occupied by the electron = 1262 K
```

---

# Chapter 4

## Geometrical Optics

Scilab code Exa 4.1 Actual path of light using Fermat principle

```
1 // Scilab Code Ex4.1: Page:189 (2011)
2 clc; clear;
3 // Declare cosine function
4 function r = cosine(t)
5     t = poly(0,t);
6     r = 1-t^2/factorial(2)+t^4/factorial(4)-t^6/
        factorial(6)+t^8/factorial(8)-t^10/factorial
        (10)+t^12/factorial(12)-t^14/factorial(14);
7 endfunction
8
9 // Declare sine function
10 function r = sine(t)
11     t = poly(0,t);
12     r = t-t^3/factorial(3)+t^5/factorial(5)-t^7/
        factorial(7)+t^9/factorial(9)-t^11/factorial
        (11)+t^13/factorial(13)-t^15/factorial(15);
13 endfunction
14
15 r = 1; // For convenience assume radius of the
        circle to be unity, unit
16 thet = poly(0,'thet'); // Declare a variable
```

```

17 l = 2*r*(cosine('thet')+sine('thet')); // Length of
    actual path, unit
18 theta = 45*%pi/180; // Angle which the chord PQ
    makes with the diameter, radian
19 d_diff = derivat(derivat(l)); // Double derivative
    of 'l' w.r.t. theta
20 printf("\nl = %5.3fr", horner(l,theta));
21 printf("\nDouble_diff of l at theta = 45 degrees =
    %5.3fr \nwhich is negative, so the actual path is
    maximum", horner(d_diff, theta));
22
23 // Result
24 // l = 2.828r
25 // Double_diff of l at theta = 45 degrees = -2.828r
26 // which is negative, so the actual path is maximum

```

---

**Scilab code Exa 4.2** Light reflected from the inner surface of spherical shell

```

1 // Scilab Code Ex4.2: Page:191 (2011)
2 clc;clear;
3 r = 1; // For convenience assume radius of the
    circle to be unity, unit
4 alpha = 0.8*r; // Distance of light source from the
    centre of the spherical shell, unit
5 cos_phi_by_2 = sqrt((alpha+1)/(4*alpha));
6 printf("\ncos(phi/2) = %d/4", cos_phi_by_2*4);
7
8 // Result
9 // alpha^2+1-2*alpha*cosine('phi')

```

---

**Scilab code Exa 4.3** Equivalent focal length of the combinations of lenses

```

1 // Scilab Code Ex4.3: Page:193 (2011)
2 clc;clear;
3 f1 = 5;....// Focal length of thin convex lens , cm
4 f2 = 3;....// Focal length of thin convex lens , cm
5 d = 2;....// Separation between the lenses , cm
6 F = f1*f2/(f1+f2-d);....// Equivalent focal length
    of a combination of the two lenses , cm
7 printf("\nThe equivalent focal length of the
    combination of lenses = %3.1f cm", F)
8
9 // Result
10 // The equivalent focal length of the combination of
    lenses = 2.5 cm

```

---

**Scilab code Exa 4.4** Focal length of the combination of lenses of given powers

```

1 // Scilab Code Ex4.4: Page:194 (2011)
2 clc;clear;
3 P1 = 5;....// Power of first converging lens ,
    diopter
4 P2 = 4;....// Power of second converging lens ,
    diopter
5 d = 0.1;....// Separation distance between two
    lenses , cm
6 P = P1+P2-d*P1*P2;
7 f = 1/P*100;....// The corresponding value of the
    focal length of the lens combination , cm
8 printf("\nThe focal length of the combination of
    lenses of given powers = %5.2f cm", f);
9
10 // Result
11 // The focal length of the combination of lenses of
    given powers = 14.29 cm

```

---

**Scilab code Exa 4.5** Focal length of the combination of coaxially placed thin convex lenses

```
1 // Scilab Code Ex4.5: Page:194 (2011)
2 clc;clear;
3 f1 = 30;....// Focal length first convex lens , cm
4 f2 = -50;....// Focal length of second convex lens ,
   cm
5 d = 20;....// Separation distance between lenses , cm
6 F = f1*f2/(f1+f2-d);....// Equivalent focal length
   of a combination of the two lenses , cm
7 printf("\nThe equivalent focal length of the
   combination = %4.1f cm", F);
8
9 // Result
10 // The equivalent focal length of the combination =
   37.5 cm
```

---

**Scilab code Exa 4.7** Locations of principal points and focal points

```
1 // Scilab Code Ex4.7 : Page-195
2 clc;clear;
3 f1 = 4;....// Focal length of thin convex lens , cm
4 f2 = 12;....// Focal length of thin convex lens , cm
5 d = 8;....// Separation distance between the lenses ,
   cm
6 F = f1*f2/(f1+f2-d);....// Equivalent focal length
   of the combination , cm
7 L1H1 = d*F/f2; // Distance of first principal
   point H1 from first lens , cm
8 printf("\nThe distance of the first principal point
   H1 from the first lens = %d cm", L1H1);
```

```

9 L2H2 = -d*F/f1;    // Distance of first principal
    point H2 from second lens , cm
10 printf("\nThe distance of the second principal point
    H2 from the second lens = %d cm", L2H2);
11 L1F1 = -F*(1-d/f2);    // Distance of first focal
    point F1 from first lens , cm
12 printf("\nThe distance of the first focal point F1
    from the first lens = %d cm", L1F1);
13 L2F2 = F*(1-d/f1);    // Distance of second focal
    point F2 from first lens , cm
14 printf("\nThe distance of the second focal point F2
    from the second lens= %d cm", L2F2);
15
16 // Result
17 // The distance of the first principal point H1 from
    the first lens = 4 cm
18 // The distance of the second principal point H2
    from the second lens = -12 cm
19 // The distance of the first focal point F1 from the
    first lens = -2 cm
20 // The distance of the second focal point F2 from
    the second lens= -6 cm

```

---

**Scilab code Exa 4.8** Position of principal points and focal points for two coaxially placed lenses

```

1 // Scilab Code Ex4.8: Page-195 (2011)
2 clc;clear;
3 f1 = 25;....// Focal length of thin convex lens , cm
4 f2 = -15;....// Focal length of thin concave lens ,
    cm
5 d = 15;....// Separation distance between the lenses
    , cm
6 // We know that ,  $F = f_1*f_2/(f_1+f_2-d)$  then
7 F = f1*f2/(f1+f2-d);....// The equivalent focal

```

```

length of the combination
8 L1H1 = d*F/f2;    // The distance of the first
principal point H1 from the first lens , cm
9 printf("\nThe distance of the first principal point
H1 from the first lens = %d cm", L1H1);
10 L2H2 = -d*F/f1;    // The distance of the second
principal point H2 from the first lens , cm
11 printf("\nThe distance of the second principal point
H2 from the second lens = %d cm", L2H2);
12 L1F1 = -F*(1-d/f2);    // The distance of the first
focal point F1 from the first lens , cm
13 printf("\nThe distance of the first focal point H1
from the first lens = %d cm", L1F1);
14 L2F2 = F*(1-d/f1);    // The distance of the second
principal point F2 from the first lens , cm
15 printf("\nThe distance of the second focal point H2
from the second lens= %d cm", L2F2);
16
17 //Result
18 // The distance of the first principal point H1 from
the first lens = -75 cm
19 // The distance of the second principal point H2
from the second lens = -45 cm
20 // The distance of the first focal point H1 from the
first lens = -150 cm
21 // The distance of the second focal point H2 from
the second lens= 30 cm

```

---

**Scilab code Exa 4.9** Focal lengths from dispersive powers of achromatic combination of lenses

```

1 // Scilab Code Ex4.9 : Page-196
2 clc;clear;
3 w1 = 0.024;....// Magnitude of the dispersive power
of first lens

```

```

4 w2 = 0.036;....// Magnitude of the dispersive power
  of second lens
5 // Let 1/f1 = x and 1/f2 = y, then
6 // The condition for achromatic combination of two
  lenses , w1/f1 + w2/f2 = 0 => w1*x + w2*y = 0
  --- (I)
7 F = 90;....// Given focal length , cm
8 // Also F = 1/f1 + 1/f2 => F = x + y ----- (II)
9 A = [w1 w2; 1 1]; // Square matrix
10 B = [0;1/F]; // Column vector
11 X = inv(A)*B; // Characteristic roots of the
  simultaneous equations , cm
12 f1 = 1/X(1); // Focal length of convex lens , cm
13 f2 = 1/X(2); // Focal length of concave lens , cm
14
15 printf("\nThe focal length of convex lens = %2d cm",
  ceil(f1));
16 printf("\nThe focal length of concave lens = %2d cm"
  , ceil(f2));
17
18 // Result
19 // The focal length of convex lens = 30 cm
20 // The focal length of concave lens = -44 cm

```

---

**Scilab code Exa 4.10** Focal length of the two component lens of an achromatic doublet

```

1 // Scilab Code Ex4.10: Page-197
2 clc;clear;
3 w1 = 0.02;....// Magnitude of the dispersive power
  of first lens
4 w2 = 0.04;....// Magnitude of the dispersive power
  of second lens
5 // Let 1/f1 = x and 1/f2 = y, then
6 // The condition for achromatic combination of two

```



```

    lenses ,  $w_1/f_1 + w_2/f_2 = 0 \Rightarrow w_1*x + w_2*y = 0$ 
    —— (I)
7 F = 20;....// Given focal length of achromatic
    doublet , cm
8 // Also  $F = 1/f_1 + 1/f_2 \Rightarrow F = x + y$  —— (II)
9 A = [w1 w2; 1 1]; // Square matrix
10 B = [0;1/F]; // Column vector
11 X = inv(A)*B; // Characteristic roots of the
    simultaneous equations , cm
12 f1 = 1/X(1); // Focal length of convex lens , cm
13 f2 = 1/X(2); // Focal length of concave lens , cm
14
15 printf("\nThe focal length of convex lens = %2d cm",
    ceil(f1));
16 printf("\nThe focal length of concave lens = %2d cm"
    , ceil(f2));
17
18 // Result
19 // The focal length of convex lens = 10 cm
20 // The focal length of concave lens = -20 cm

```

---

**Scilab code Exa 4.11** Radii of curvature of the second surface of each of the lens of achromatic doublet

```

1 // Scilab Code Ex4.11: Page-197
2 clc;clear;
3 w1 = 0.017;....// Magnitude of the dispersive power
    of first lens
4 w2 = 0.034;....// Magnitude of the dispersive power
    of second lens
5 // Let  $1/f_1 = x$  and  $1/f_2 = y$ , then
6 // The condition for achromatic combination of two
    lenses ,  $w_1/f_1 + w_2/f_2 = 0 \Rightarrow w_1*x + w_2*y = 0$ 
    —— (I)
7 F = 40;....// Given focal length of achromatic

```

```

doublet , cm
8 // Also  $F = 1/f_1 + 1/f_2 \Rightarrow F = x + y$  ----- (II)
9 A = [w1 w2; 1 1]; // Square matrix
10 B = [0;1/F]; // Column vector
11 X = inv(A)*B; // Characteristic roots of the
simultaneous equations , cm
12 f1 = 1/X(1); // Focal length of convex lens , cm
13 f2 = 1/X(2); // Focal length of concave lens , cm
14 // For the convex lens
15 R2 = -25; // Radius of curvature of the contact
surface , cm
16 mu = 1.5; // Mean refractive index of crown glass
17 // From the Lens Maker formula,  $1/f = (\mu - 1)*(1/R_1$ 
 $-1/R_2)$ , solving for R1
18 f = f1;
19 R1 = 1/(1/(f*(mu-1))+1/R2); // Radius of
curvature of second surface of first lens , cm
20 printf("\nThe radius of curvature of second surface
of first lens = %5.2f cm", R1);
21 // For the concave lens
22 R1 = -25; // Radius of curvature of the contact
surface , cm
23 mu = 1.7; // Mean refractive index of flint glass
24 // From the Lens Maker formula,  $1/f = (\mu - 1)*(1/R_1$ 
 $-1/R_2)$ , solving for R1
25 f = f2;
26 R2 = 1/(1/R1-1/(f*(mu-1))); // Radius of
curvature of second surface of second lens , cm
27 printf("\nThe radius of curvature of second surface
of second lens = %5.2f cm", R2);
28
29 // Result
30 // The radius of curvature of second surface of
first lens = 16.67 cm
31 // The radius of curvature of second surface of
second lens = -233.33 cm

```

---

**Scilab code Exa 4.12** Focal length of the convergent lens for C line

```
1 // Scilab Code Ex4.12: Page-199
2 clc; clear;
3 // For flint glass
4 mu_C = 1.665; // Refractive index of flint glass
   for C line
5 mu_F = 1.700; // Refractive index of flint glass
   for F line
6 mu_D = (mu_F+mu_C)/2; // Refractive index of
   flint glass for D line
7 w2 = (mu_F-mu_C)/(mu_D-1);....// Magnitude of the
   dispersive power of second lens of flint glass
8 // For crown glass
9 mu_C = 1.510; // Refractive index of crown glass
   for C line
10 mu_F = 1.536; // Refractive index of crown glass
   for F line
11 mu_D = (mu_F+mu_C)/2; // Refractive index of
   flint glass for D line
12 w1 = (mu_F-mu_C)/(mu_D-1);....// Magnitude of the
   dispersive power of second lens of crown glass
13 f = 50; // Focal length of acromatic doublet, cm
14 FD = f*(w2-w1)/w2; // Focal length of D line of
   the Fraunhofer spectrum due to convex lens of
   crown glass
15 FC = FD*(mu_D - 1)/(mu_C - 1); // Focal length of
   C component of converging lens, cm
16 printf("\\nThe focal length of C component of
   converging lens = %4.2f cm", FC);
17
18 // Result
19 // The focal length of C component of converging
   lens = 1.57 cm
```

---

**Scilab code Exa 4.13** Focal length of two lenses with no aberration

```
1 // Scilab Code Ex4.13 Page-200
2 clc;clear;
3 F = 50;....// Equivalent focal length of combination
   of two lenses , cm
4 //d = f1+f2/2, condition for no chromatic aberration
   ....(1)
5 //d = f2-f1, condition for minimum spherical
   aberration ....(2)
6 // From (1) and (2), f1 = 3*d/2, f2 = d/2
7 // As 1/F = 1/f1 + 1/f2 - d/(f1*f2), solving for d
8 d = 4/3*50; // Distance of separation between two
   lenses , cm
9 f1 = 3*d/2, f2 = d/2;
10 printf(" \nf1 = %3d cm, f2 = %5.2 f cm", ceil(f1), f2)
   ;
11
12 // Result
13 // f1 = 100 cm, f2 = 33.33 cm
```

---

**Scilab code Exa 4.14** Longitudinal chromatic aberration for an object at infinity

```
1 // Scilab Code Ex4.14 : Page-200 (2011)
2 clc;clear;
3 mu_R = 1.5230; // Refractive index for red
   wavelength
4 mu_V = 1.5145; // Refractive index for violet
   wavelength
5 R1 = 40; // Radius of curvature for red
   wavelength , cm
```

```

6 R2 = -10;    // Radius of curvature for violet
   wavelength, cm
7 // As  $1/f = (\mu - 1) * (1/R1 - 1/R2)$ , solving for fV
   and fR
8 fV = 1/((mu_V-1)*(1/R1-1/R2));    // Focal length
   for violet wavelength, cm
9 fR = 1/((mu_R-1)*(1/R1-1/R2));    // Focal length
   for violet wavelength, cm
10 l = fR - fV;    // Longitudinal chromatic aberration
   , cm
11 printf("\nThe longitudinal chromatic aberration = %5
   .3f cm", abs(l));
12
13 // Result
14 // The longitudinal chromatic aberration = 0.253 cm

```

---

**Scilab code Exa 4.15** Focal length of component lenses of a convergent doublet

```

1 // Scilab Code Ex4.15: Page-202 (2011)
2 clc;clear;
3 F = 10;....// Equivalent focal length of a
   combination of two lenses, cm
4 d = 2;....// Separation distance between two lenses,
   cm
5 // As  $d = f1 - f2$ , condition for minimum spherical
   aberration  $\Rightarrow f1 = d + f2$ 
6 // and  $F = f1 * f2 / (f1 + f2 - d)$ , so solving for f2
7 f2 = 2 * F - d;    // Focal length of second lens, cm
8 f1 = d + f2;    // Focal length of first lens, cm
9 printf("\nf1 = %2d cm, f2 = %2d cm", f1, f2);
10
11 // Result
12 // f1 = 20 cm, f2 = 18 cm

```

---

**Scilab code Exa 4.16** Radii of Aplanatic surfaces and lateral magnification of the image

```
1 // Scilab Code Ex4.16: Page-202 (2011)
2 clc;clear;
3 mu = 1.6;....// Refractive index of aplanatic
   surface
4 R = 3.2;....// Radius of curvature, cm
5 R1 = R/mu;....// First radius of the aplanatic
   surface, cm
6 printf("\nR1 = %3.1 f cm", R1);
7 R2 = R*mu;....// Second radius of the aplanatic
   surface, cm
8 printf("\nR2 = %4.2 f cm", R2);
9 //Since the image of an object at one aplanatic
   point will be formed by the sphere at the other
   aplanatic point,so the is
10 m = mu^2; // The lateral magnification of the
   image
11 printf("\nThe lateral magnification of the image =
   %4.2 f", m);
12
13 // Result
14 // R1 = 2.0 cm
15 // R2 = 5.12 cm
16 // The lateral magnification of the image = 2.56
```

---

**Scilab code Exa 4.17** Aplanatic surface

```
1 // Scilab Code Ex4.17: Page-203 (2011)
2 clc;clear;
```

```

3 mu = 1.52;....// Refractive index of aplanatic
  surface
4 R = 30;....// Radius of curvature, cm
5 R1 = R/mu;....// First radius of the aplanatic
  surface, cm
6 printf("\nR1 = %5.2 f cm", R1);
7 R2 = R*mu;....// Second radius of the aplanatic
  surface, cm
8 printf("\nR2 = %4.1 f cm", R2);
9 //Since the image of an object at one aplanatic
  point will be formed by the sphere at the other
  aplanatic point,so the is
10 m = mu^2;    // The lateral magnification of the
  image
11 printf("\nThe lateral magnification of the image =
  %4.2 f", m);
12
13 // Result
14 // R1 = 19.74 cm
15 // R2 = 45.6 cm
16 // The lateral magnification of the image = 2.31

```

---

#### Scilab code Exa 4.18 Focal length of the field lens

```

1 // Scilab Code Ex4.18: Page-203 (2011)
2 clc;clear;
3 F = 5;....// Equivalent focal length of Huygens
  eyepiece, cm
4 // as f1 = 3*f, f2 = f and d = 2*f, therefore
5 f = 2/3*F;    // Focal length of base lens, cm
6 f1 = 3*f;    // Focal length of field lens, cm
7 printf("\nThe focal length of the field lens = %2d
  cm", f1);
8
9 // Result

```

```
10 // The focal length of the field lens = 10 cm
```

---

**Scilab code Exa 4.19** Equivalent focal length of a Ramsden eyepiece

```
1 // Scilab Code Ex4.19: Page-204 (2011)
2 clc;clear;
3 f = 10;....// Given focal length of each lens , cm
4 f1 = f;     // Focal length of first lens , cm
5 f2 = f;     // Focal length of second lens , cm
6 d = 2/3*f;  // Separation distance between two
              lenses , cm
7 F = f1*f2/(f1+f2-d); // Equivalent focal length
              of Ramsden eyepiece , cm
8 printf("\nThe equivalent focal length of the field
          lenses is = %3.1f cm", F);
9
10 // Result
11 // The equivalent focal length of the field lenses
    is = 7.5 cm
```

---

**Scilab code Exa 4.20** Focal lengths of the lenses and the eyepiece

```
1 // Scilab Code Ex4.20 : Page-204 (2011)
2 clc;clear;
3 d = 10;....// Distance between the two thin plano
              convex lenses in the Huygens eyepiece ,
4 f = d/2;    // Base focal length
5 f1 = 3*f;   // Focal length of the first component
              lens , cm
6 printf("\nf1 = %d cm", f1);
7 f2 = f;     // Focal length of the second component
              lens , cm
8 printf("\nf2 = %d cm", f2);
```



```

9 F = 3/2*f; // Equivalent focal length of the lens ,
    cm
10 printf("\nF = %3.1 f cm", F);
11
12 // Result
13 // f1 = 15 cm
14 // f2 = 5 cm
15 // F = 7.5 cm

```

---

**Scilab code Exa 4.21** Focal length of the component lenses and the separation between them

```

1 // Scilab Code Ex4.21: Page-204 (2011)
2 clc;clear;
3 F = 4.2;....// Equivalent focal length of Ramsden
    eyepiece , cm
4 //F = 3/4*f, Equivalent focal length of Ramsden
    eyepiece ,
5 f = 5.6;....//focal length ,in cm
6 f1 = f;
7 f2 = f;
8 printf("\nf1 = %3.1 f cm", f1);
9 printf("\nf2 = %3.1 f cm", f2);
10 d = 2/3*f;
11 printf("\nd = %4.2 f cm", d);
12
13 // Result
14 // f1 = 5.6 cm
15 // f2 = 5.6 cm
16 // d = 3.73 cm

```

---

# Chapter 5

## Physical Optics

**Scilab code Exa 5.1** Order of interference maximum with different wavelength

```
1 // Scilab Code Ex5.1: Page:297 (2011)
2 clc;clear;
3 n1 = 10;....// Order of interference maximum for
   lambda = 7000 angstrom
4 lambda1 = 7000;....// Wavelength of the light ,
   angstrom
5 lambda2 = 5000;....// Wavelength of the light ,
   angstrom
6 // As  $W = D \cdot \lambda / (2 \cdot d)$  then ,  $x = n1 \cdot D \cdot \lambda1 / (2 \cdot d)$ 
   ) =  $n2 \cdot D \cdot \lambda2 / (2 \cdot d)$  , solving for n2
7 n2 = n1 * lambda1 / lambda2; // Order of interference
   maximum for lambda = 5000 angstrom
8 printf("\nThe order of interference maximum for
   wavelength of 5000 angstrom = %2d ", n2);
9
10 // Result
11 // The order of interference maximum for wavelength
   of 5000 angstrom = 14
```

---

### Scilab code Exa 5.2 Angle of the biprism

```
1 // Scilab Code Ex5.2: Page:297 (2011)
2 clc;clear;
3 D = 1.6;....// Distance between the slit and the
   screen , m
4 a = 0.4;....// Distance between the slit and the
   biprism , m
5 mu = 1.52;....// Refractive index of the material of
   biprism
6 W = 1e-004; // Fringe width , m
7 lambda = 5.893e-007;....// Wavelength of light used ,
   m
8 // As  $W = \lambda D / (2 * a * (\mu - 1) * \alpha)$  then
9 alpha = ((lambda*D)/(2*a*(mu-1)*W))*180/%pi; //
   Angle of biprism , degrees
10 printf("\nThe angle of the biprism = %3.1f degrees",
   alpha);
11
12 // Result
13 // The angle of the biprism = 1.3 degrees
```

---

### Scilab code Exa 5.3 Thickness of the mica sheet

```
1 // Scilab Code Ex5.3 : Page:298 (2011)
2 clc;clear;
3 lambda = 5.890e-7;....// Wavelength of source of
   light , m
4 mu = 1.6;....//refractive index of the mica sheet
5 // As  $\text{del}_x = W * (\mu - 1) * t / \lambda$ , where  $\text{del}_x = 3 * W$ ,
   solving for t
```

```

6 t = 3*lambda/(mu-1);    // Thickness of the mica
  sheet , m
7 printf("\nThe thickness of the mica sheet = %5.3e cm
  ", t/1e-02);
8
9 // Result
10 // The thickness of the mica plate is = 2.945e-004
  cm

```

---

**Scilab code Exa 5.4** Distance between the two coherent sources

```

1 //Scilab Code Ex5.4: Page:298 (2011)
2 clc;clear;
3 lambda = 6.0e-7;....// Wavelength of the
  monochromatic light , m
4 D = 1;....// Distance between the screen and the two
  coherent sources , m
5 W = 5e-004;....// Fringe width , m
6 d = lambda*D/(W*1e-03);    // Distance between two
  coherent sources , mm
7 printf("\nThe distance between the two coherent
  sources = %3.1f mm", d);
8
9 // Result
10 // The distance between the two coherent sources =
  1.2 mm

```

---

**Scilab code Exa 5.5** Wavelength of light used in a biprism experiment

```

1 // Scilab Code Ex5.5: Page:298 (2011)
2 clc;clear;
3 D = 1;....// Distance between slits and the screen ,
  m

```

```

4 mu = 1.5;    // Refractive index of the material of
    biprism
5 a = 0.5;....// The distance between the slit and the
    biprism , m
6 W = 1.35e-004;....// Width of the fringes , m
7 alpha = (180-179)/2*%pi/180;    // Acute angle of
    biprism , radian
8 lambda = 2*a*(mu-1)*alpha*W/D;    // Wavelength of
    light used , m
9 printf("\nThe wavelength of light used = %4d
    angstrom" , lambda/1e-10);
10
11 // Result
12 // The wavelength of light used = 5890 angstrom

```

---

#### Scilab code Exa 5.6 Distance between the slits

```

1 // Scilab Code Ex5.6: Page:299 (2011)
2 clc;clear;
3 lambda = 6.328e-007;....// Wavelength of the
    monochromatic light , m
4 D = 40;....// Distance between the slits and the
    screen , m
5 W = 0.1;....// Distance between the interference
    maxima , m
6 d = lambda*D/W;    // Distance between the slits , m
7 printf("\nThe distance between the slits = %6.4f mm"
    ,d/1e-03);
8
9 // Result
10 // The distance between the slits = 0.2531 mm

```

---

#### Scilab code Exa 5.7 Lateral shift of central maximum

```

1 // Scilab Code Ex5.7: Page:299 (2011)
2 clc; clear;
3 lambda = 5.0e-007;....// Wavelength of the
   monochromatic light , m
4 D = 1;....// Distance between the slits and the
   screen , m
5 d = 5e-004/2;....// Half of the distance between the
   two slits , m
6 mu = 1.5;....// Refractive index of glass
7 t = 1.5e-006;....// Thickness of thin glass plate , m
8 del_x = D*(mu-1)*t/(2*d);
9 printf("\\nThe lateral shift of central maximum = %3
   .1f m", del_x/1e-03);
10
11 // Result
12 // The lateral shift of central maximum = 1.5 m

```

---

#### Scilab code Exa 5.8 Thickness of a soap bubble film

```

1 // Scilab Code Ex5.8: Page:300 (2011)
2 clc; clear;
3 lambda = 6.0e-007;....// Wavelength of the light , m
4 mu = 1.463;....// Refractive index of a soap bubble
   film
5 n = 0; // Value of n for smallest thickness
6 r = 0; // Angle of refraction for normal
   incidence
7 // As  $2*\mu*t*\cos(r) = (2*n+1)*\lambda/2$ , solving for
   t
8 t = (2*n+1)*lambda/(4*mu*cos(r)); // The
   thickness of a soap bubble film , m
9 printf("\\nThe thickness of a soap bubble film = %5.1
   f angstrom", t/1e-010);
10
11 // Result

```

```
12 // The thickness of a soap bubble film = 1025.3
    angstrom
```

---

**Scilab code Exa 5.9** Wavelength of the light used in Newton rings experiment

```
1 // Scilab Code Ex5.9: Page:300 (2011)
2 clc;clear;
3 D5 = 3.36e-003;....// Diameter of Newton's 5th ring ,
    m
4 D15 = 5.90e-003;....// Diameter of Newton's 15th
    ring , m
5 m = 10; // Number of ring
6 R = 1;....// Radius of the plano-convex lens , m
7 lambda = (D15^2-D5^2)/(4*m*R);
8 printf("\\nThe wavelength of the light used = %4d
    angstrom", lambda/1e-010);
9
10 // Result
11 // The wavelength of the light used = 5880 angstrom
```

---

**Scilab code Exa 5.10** Radius of the curvature of the lens and the thickness of the air film

```
1 // Scilab Code Ex5.10: Page:301 (2011)
2 clc;clear;
3 D10 = 0.005;....// Diameter of Newton's 5th ring , m
4 n = 10;....// Order of the ring
5 lambda = 6.0e-007;....// Wavelength of the light
    used , m
6 R = (D10^2)/(4*n*lambda); // Radius of the
    curvature of the lens , m
```

```

7 printf("\nThe radius of the curvature of the lens =
      %6.4 f m", R);
8 t = D10^2/(8*R);
9 printf("\nThe thickness of the corresponding air
      film = %3.1 e m", t);
10
11 // Result
12 // The radius of the curvature of the lens = 1.0417
      m
13 // The thickness of the corresponding air film = 3.0
      e-006 m m

```

---

**Scilab code Exa 5.11** Thickness of the soap film

```

1 // Scilab Code Ex5.11: Page-301 (2011)
2 clc;clear;
3 mu = 1.43;....// Refractive index of the soap film
4 n = 0; // Order of fringes for smallest thickness
5 i = 30; // Angle of incidence, degrees
6 // As  $\sin(i)/\sin(r) = \mu$ ,  $\cos(r)$ 
7 cosr = sqrt(1-(sind(i)/mu)^2); // Cosine of angle
      r
8 lambda = 6.0e-007;....// Wavelength of the light, m
9 t = (2*n+1)*lambda/(4*mu*cosr);....// Thickness of
      the soap film, m
10 printf("\nThe thickness of the soap film = %4.2 e m",
      t);
11
12 // Result
13 // The thickness of the soap film = 1.12e-007 m

```

---

**Scilab code Exa 5.12** Least thickness of the soap film that will appear bright dark



```

1 // Scilab Code Ex5.12: Page:301 (2011)
2 clc;clear;
3 lambda = 5.893e-007;....// Wavelength of the sodium
   light , m
4 mu = 1.42;....// Refractive index of the soap film
5 r = 0; // Angle of refraction , degrees
6 n = 0; // Order of diffraction for least
   thickness of dark film
7 t = (2*n+1)*lambda/(4*mu*cosd(r)); // Least
   thickness of the film that will appear bright , m
8 printf("\\nThe least thickness of the film that will
   appear bright = %5.1f m", t/1e-010);
9 n = 1; // Order of diffraction for least
   thickness of bright film
10 t = n*lambda/(2*mu*cosd(r)); // Least thickness
   of the film that will appear dark , m
11 printf("\\nThe least thickness of the film that will
   appear dark = %6.2f m",t/1e-010);
12
13 // Result
14 // The least thickness of the film that will appear
   bright = 1037.5 m
15 // The least thickness of the film that will appear
   dark = 2075.00 m

```

---

**Scilab code Exa 5.13** Thickness of the wire separating edges of two plane glass surfaces

```

1 // Scilab Code Ex5.13: Page:302 (2011)
2 clc;clear;
3 lambda = 5.893e-007;....// Wavelength of the sodium
   light , m
4 // As fringe width of the thin wedge-shaped air film
   is
5 //  $W = \lambda / (2 * t / 20 * W)$  , solving for t

```

```

6 t = (10*lambda); // Thickness of the wire
   separating edges of two plane glass surfaces , m
7 printf("\nThe thickness of the wire = %5.3e m", t);
8
9 // Result
10 // The thickness of the wire = 5.893e-006 m

```

---

**Scilab code Exa 5.14** Radius of curvature of the lens and the thickness of the corresponding air film

```

1 // Scilab Code Ex5.14: Page:303 (2011)
2 clc;clear;
3 lambda = 5.9e-007;....// Wavelength of the reflected
   light , m
4 n = 10;....// Order of the ring
5 D10 = 0.005;....// Diameter of the 10th ring ,in m
6 R = (D10^2)/(4*n*lambda); // Radius of curvature
   of the lens , m
7 printf("\nThe radius of curvature of the lens = %6.4
   f m", R);
8 t = (D10^2)/(8*R); // Thickness of the
   corresponding air film , m
9 printf("\nThe thickness of the corresponding air
   film = %4.2e m",t);
10
11 // Result
12 // The radius of curvature of the lens = 1.0593 m
13 // The thickness of the corresponding air film =
   2.95e-006 m

```

---

**Scilab code Exa 5.16** Angles at which first and second order maxima can be observed

```

1 // Scilab Code Ex5.16: Page:304 (2011)
2 clc;clear;
3 lambda = 6.328e-007;....// Wavelength of
   monochromatic light from He laser , m
4 n1 = 1;....// First order
5 n2 = 2;....// Second order
6 l = 6000;....// Lines/cm of the diffraction grating
7 A= 1.66e-6;
8 theta = asind(n1*lambda/A);
9 printf("\\n The first order maximum angle = %4.1f
   degrees",theta);
10 theta = asind(n2*lambda/A);
11 printf("\\n The second order maximum angle = %4.1f
   degrees",theta);
12
13 // Result
14 // The first order maximum angle = 22.4 degrees
15 // The second order maximum angle = 49.7 degrees

```

---

**Scilab code Exa 5.17** Relation between two wavelengths illuminating a single slit due to Fraunhofer diffraction

```

1 // Scilab Code Ex5.17: Page:305 (2011)
2 clc;clear;
3 a = 1;    // For simplicity assume slit width to be
   unity , unit
4 theta = 1; // For simplicity assume diffraction
   angle to be unity , unit
5 // As  $a*\sin(\theta) = m*\lambda$ , solving for lambdas
6 lambda1 = a*sin(theta);    // First wavelength ,
   angstrom
7 lambda2 = a*sin(theta)/2;    // First wavelength ,
   angstrom
8 printf("\\nlambda1 = %d*lambda2", lambda1/lambda2);
9

```

```

10 // Result
11 // lambda1 = 2*lambda2

```

---

**Scilab code Exa 5.18** Angular position of the first two minima on either side of a central maxima

```

1 // Scilab Code Ex5.18: Page:305 (2011)
2 clc;clear;
3 function [deg, minute] = deg2degmin(theta)
4     deg = int(theta);
5     minute = (theta - deg)*60;
6 endfunction
7
8 lambda = 5.5e-007;....// Wavelength of light , m
9 a = 2.2e-006;....// Width of the slit , m
10 l = 6000;....// Lines /cm of the diffraction grating
11 // In a single slit diffraction pattern the
    directions of minimum intensity are given by a*
    sintheta = m*lambda where m = 1,2,3....
12 // For m = 1
13 m = 1;....// First order
14 theta = asind(m*lambda/a);    // Angular position of
    first minima on either side of the central
    maxima, degrees
15 [deg, minute] = deg2degmin(theta);    // Degree to
    deg-min conversion
16 printf("\nThe angular position of first minima on
    either side of the central maxima = %2d degrees
    %2d minutes", deg, minute);
17 // For m = 2
18 m = 2;....// Second order
19 theta = asind(m*lambda/a);
20 [deg, minute] = deg2degmin(theta);    // Degree to
    deg-min conversion
21 printf("\nThe angular position of second minima on

```

```

    either side of the central maxima = %2d degrees
    %2d minutes", deg, minute);
22
23 // Result
24 //

```

---

**Scilab code Exa 5.19** Wavelength of light and the missing order of Fraunhofer diffraction

```

1 // Scilab Code Ex5.19: Page:306 (2011)
2 clc; clear;
3 D = 1.7;....// Distance between the slit and the
    screen, m
4 W = 2.5e-003;....// Given fringe width, m
5 a = 8e-005;....// Width of the first slit, m
6 b = 4e-004;....// Width of the second slit, m
7 n = b; //
8 p = [1 2 3 4 5 6];
9 // In a double slit experiment Fraunhofer
    diffraction pattern, the fringe width is given by
    W = lambda*D/n
10 lambda = b*W/D; // Wavelength of the light used,
    m
11 printf("\\nThe wavelength of light = %4d angstrom",
    lambda/1e-010);
12 printf("\\nThe missing orders are:\\n");
13 for i = 1:1:6
14 s = [(a+b)/a]*p(i);
15 printf("\\t%d", s);
16 end
17 printf(" etc.")
18 // Result
19 // The wavelength of light = 5882 angstrom
20 // The missing orders are:
21 // 6 12 18 24 30 36 etc.

```

---

**Scilab code Exa 5.20** Deduction of wavelength of the light from given data

```
1 //Scilab Code Ex5.20: Page-306 (2011)
2 clc; clear;
3 D = 2;....// Distance of the screen from the slit , m
4 x = 1.6e-02;....// Position of centre of the second
   dark band, m
5 m = 2;    // Order of diffraction
6 a = 1.4e-04;....// Width of the slit, m
7 lambda = a*x/(m*D);    // Wavelength of light , m
8 printf("\\n The wavelength of the light = %4d
   angstrom", ceil(lambda/1e-010));
9
10 // Result
11 // The wavelength of the light = 5600 angstrom
```

---

**Scilab code Exa 5.21** Minimum number of lines in a grating

```
1 // Scilab Code Ex5.21: Page:307 (2011)
2 clc; clear;
3 lambda1 = 5890;....// Wavelength of the line ,
   angstrom
4 lambda2 = 5896;....// Wavelength of the line ,
   angstrom
5 d_lambda = lambda2 - lambda1;....// Wavelength
   difference , angstrom
6 n = 2;....// Order of diffraction
7 N = lambda2/(n*d_lambda);    // Minimum no. of lines
   in a grating
8 printf("\\nThe minimum number of lines in the grating
   = %3d lines", N);
```

```

9
10 // Result
11 // The minimum number of lines in the grating = 491
    lines

```

---

**Scilab code Exa 5.22** Maximum number of visible orders

```

1 // Scilab Code 5.22: Page:307 (2011)
2 clc;clear;
3 lambda = 5.0e-07;....// Wavelength of the radiation ,
    m
4 a_plus_b = 2.54e-02/2620;....// The grating element ,
    m
5 theta_max = 90;    // Maximum value of angle of
    diffraction , degrees
6 n_max = a_plus_b/lambda*sind(theta_max);    //
    Maximum number of visible orders
7 printf("\\nThe number of visible orders = %2d ",
    n_max);
8
9 // Result
10 // The number of visible orders = 19

```

---

**Scilab code Exa 5.23** Grating element of diffraction grating

```

1 // Scilab Code 5.23: Page:307 (2011)
2 clc;clear;
3 lambda1 = 6000;....// Wavelength of yellow line ,
    angstrom
4 lambda2 = 4800;....// Wavelength of blue line ,
    angstrom
5 theta = asin(3/4);    // Angle of diffraction ,
    radian

```

```

6 // As  $a_{plus\_b} \sin(\theta) = n \lambda$ , so  $n \lambda_1 =$ 
   (n+1)*lambda2, solving for n
7 n = poly(0, 'n');
8 n = roots(n*6000 - (n+1)*4800); // Order of
   diffraction
9 a_plus_b = n*6000/sin(theta); // Grating element
   of diffraction grating, m
10 printf("\nThe Grating element of diffraction grating
   = %3.1e m", a_plus_b*1e-010);
11
12 // Result
13 // The Grating element of diffraction grating = 3.2e
   -006 m

```

---

#### Scilab code Exa 5.26 Coinciding spectral lines

```

1 // Scilab Code Ex5.26: Page:310 (2011)
2 clc;clear;
3 n = 5;...// Order for given wavelength
4 m = [4 5 6 7 8]; // Orders of spectral lines in
   the visible range
5 lambda1 = 6000;...// Wavelength of the spectral
   line in visible range, angstrom
6 lambda2 = zeros(5);
7 printf("\n The spectral lines in visible ranges are
   :\n");
8 for i=1:1:5
9 l2 = (n*lambda1)/m(i);
10 lambda2(i) = l2; // Preserve the lambda value
11 printf("%4d angstrom\n", ceil(l2));
12 end
13 printf("\nThe other spectral lines in the visible
   range 4000A to 7000A are");
14 for i=1:1:5
15     if lambda2(i) < 7000 & lambda2(i) > 4000 then

```



```

16         if lambda2(i) == 6000 then
17             continue
18         end
19         printf("\n%4dA", ceil(lambda2(i)));
20     end
21 end
22
23 // Result
24 // The spectral lines in visible ranges are:
25 // 7500 angstrom
26 // 6000 angstrom
27 // 5000 angstrom
28 // 4285 angstrom
29 // 3750 angstrom
30
31 // The other spectral lines in the visible range
    4000A to 7000A are
32 // 5000A
33 // 4286A

```

---

**Scilab code Exa 5.27** Resolution of D1 and D2 lines of sodium

```

1 // Scilab Code Ex5.27: Page:310 (2011)
2 clc;clear;
3 N = 4500;....// Number of lines in grating
4 n = 2;....// Order of diffraction
5 lambda1 = 5890;....// Wavelength, angstrom
6 lambda2 = 5896;....// Wavelength, angstrom
7 RP2 = n*N;    // Resolving power of grating in the
    second order
8 lambda = (lambda1+lambda2)/2;    // Mean wavelength
    of sodium light, angstrom
9 d_lambda = lambda2 - lambda1;    // Wavelength
    difference, angstrom
10 RP = lambda/d_lambda;    // Calculated resolving

```

```

        power of grating
11 if RP2 <> RP then
12     printf("\n\nThe D1 and D2 lines of Na light cannot
        be resolved in second order");
13 end
14
15 // Result
16 // The D1 and D2 lines of Na light cannot be
        resolved in second order

```

---

**Scilab code Exa 5.28** Distance between two stars which are just resolved

```

1 // Scilab Code Ex5.28: Page:311 (2011)
2 clc;clear;
3 lambda = 5.5e-07;....// Wavelength of light used , m
4 f = 3.0;....// Focal length of telescope objective ,
        m
5 a = 0.01;....// Diameter of the telescope objective ,
        m
6 // As  $x/f = 1.22*\lambda/a$ , the Rayleigh criterion
        for resolution , solving for x
7 x = 1.22*f*lambda/a; // Distance between two
        stars just seen as separate , m
8 printf("\n\nThe distance between two stars just seen
        as separate = %3.1e m ", x);
9
10 // Result
11 // The distance between two stars just seen as
        separate = 2.0e-004 m

```

---

**Scilab code Exa 5.29** Numerical aperture of the objective of microscope

```

1 // Scilab Code Ex5.29: Page:311 (2011)

```

```

2 clc;clear;
3 lambda = 5.461e-07;....// Wavelength of light used ,
   m
4 d = 4.0e-07;....// Distance between the two luminous
   objects , m
5 // As  $d = 1.22 \cdot \lambda / (2 \cdot \mu \cdot \sin(\alpha)) = 1.22 \cdot$ 
    $\lambda / (2 \cdot NA)$ , solving for NA
6 NA = 1.22*lambda/(2*d); // Numerical aperature
   of the objective of microscope
7 printf("\\nThe numerical aperature of the objective
   of microscope = %5.3f ", NA);
8
9 // Result
10 // The numerical aperature of the objective of
   microscope = 0.833

```

---

**Scilab code Exa 5.30** Aperture of the objective of a telescope

```

1 // Scilab Code Ex5.30: Page:312 (2011)
2 clc;clear;
3 lambda = 6.0e-07;....// Wavelength of light used , m
4 d_theta = 2.44e-06;....// Angular separation between
   the two stars , radian
5 a = 1.22*lambda/d_theta; // Aperature of the
   objective of a telescope from Rayleigh criterion ,
   m
6 printf("\\nThe aperature of the objective of the
   telescope = %3.1f m ", a)
7
8 // Result
9 // The aperature of the objective of the telescope =
   0.3 m

```

---

**Scilab code Exa 5.31** Minimum distance from the telescope at which the the pinhole can be resolved

```
1 // Scilab Code Ex5.31:Page:312 (2011)
2 clc;clear;
3 lambda = 5.5e-007;....// Wavelength of light used, m
4 x = 1.5e-003;....// Distance between the two
   pinholes, m
5 a = 4.0e-003;....// Diameter of objective, m
6 D = a*x/(1.22*lambda); // Minimum distance from
   the telescope at which the the pinhole can be
   resolved from Rayleigh criterion, m
7 printf("\n\nThe minimum distance from the telescope at
   which the the pinhole can be resolved = %4.2f m
   ", D);
8
9 // Result
10 // The minimum distance from the telescope at which
   the the pinhole can be resolved = 8.9
```

---

**Scilab code Exa 5.32** Numerical aperture of the objective of microscope for given wavelength of light

```
1 // Scilab Code Ex5.32: Page:312 (2011)
2 clc;clear;
3 lambda = 5.461e-07;....// Wavelength of light used,
   m
4 d = 5.55e-07;....// Distance between the two
   luminous objects, m
5 // As  $d = 1.22 * \lambda / (2 * \mu * \sin(\alpha)) = 1.22 * \lambda / (2 * NA)$ , solving for NA
6 NA = 1.22*lambda/(2*d); // Numerical aperture
   of the objective of microscope
7 printf("\n\nThe numerical aperture of the objective
   of microscope = %4.2f ", NA);
```

```

8
9 // Result
10 // The numerical aperature of the objective of
    microscope = 0.60

```

---

**Scilab code Exa 5.33** Angle of minimum deviation for green light for its passage through a prism

```

1 // Scilab Code Ex5.33: Page:313 (2011)
2 clc;clear;
3 i = 60; // Angle of incidence , degrees
4 mu = tand(i); // Brewster's Law to calculate
    refractive index
5 A = 60;....// Angle of prism , degrees
6 // As mu = sind((A+delta_m)/2)/sind(A/2) , solving
    for delta_m
7 delta_m = 2*asind(mu*sind(A/2))-A; // Angle of
    minimum deviation for green light for its passage
    through a prism , degrees
8 printf("\nThe angle of minimum deviation for green
    light for its passage through a prism = %2d
    degrees", ceil(delta_m));
9
10 // Result
11 // The angle of minimum deviation for green light
    for its passage through a prism = 60 de

```

---

**Scilab code Exa 5.34** Thickness of quarter wave plate

```

1 // Scilab Code Ex5.34: Page:313 (2011)
2 clc;clear;
3 lambda = 5.89e-07;....// Wavelength of light used , m

```

```

4 mu_0 = 1.55;    // Refractive index of ordinary
  light
5 mu_E = 1.54;    // Refractive index of extraordinary
  light
6 tQ = lambda/(4*(mu_0-mu_E));    // The thickness of
  the quarter wave plate , m
7 printf("\nThe thickness of the quarter plate is = %6
  .4e m", tQ);
8
9 // Result
10 // The thickness of the quarter plate is = 1.4725e
  -005 m

```

---

**Scilab code Exa 5.35** Percentage purity of the sugar sample

```

1 // Scilab Code Ex5.35: Page:313 (2011)
2 clc;clear;
3 theta = 9.9;....// Optical rotation of solution ,
  degrees
4 l = 20;....// Length of the tube, cm
5 S = 66;....// Specific rotation of pure sugar
  solution , degree per dm-(g/cc)
6 // As the specific rotation , S = 10*theta/l*c,
  solving for c
7 c = 10*theta/(l*S);    // Concentration of solution
  for pure sugar , g/cc
8 c_prime = 0.080;    // Concentration of solution for
  impure sugar , g/cc
9 Percentage_purity = c*100/c_prime;    // Percentage
  purity of sugar sample
10 printf("\nThe percentage_purity of the sugar sample
  = %5.2f percent", Percentage_purity);
11
12 // Result
13 // The percentage_purity of the sugar sample = 93.75

```

percent

---

**Scilab code Exa 5.36** Specific rotation of sugar solution for given plane of polarization

```
1 // Scilab Code Ex5.36: Page:314 (2011)
2 clc;clear;
3 theta = 26.4;....// Optical rotation of sugar
   solution , degrees
4 l = 20;....// Length of the tube , cm
5 c = 0.20;....// Concentration of the solution , g/cc
6 S = 10*theta/(l*c); // The specific rotation of
   the sugar solution , degree per dm per (g/cc)
7 printf("\nThe specific rotation of the sugar
   solution = %2d degrees",S);
8
9 // Result
10 // The specific rotation of the sugar solution = 66
   degrees
```

---

**Scilab code Exa 5.37** Angle of rotation produced by quartz plate

```
1 // Scilab Code Ex5.37: Page:315 (2011)
2 clc;clear;
3 // Function to convert degrees to deg-min
4 function [d,m] = deg2degmin(deg)
5     d = int(deg);
6     m = (deg-d)*60;
7 endfunction
8
9 lambda = 7.62e-07;....// Wavelength of the polarized
   light , m
```

```

10 mu_R = 1.53914;    // Refractive index of quartz for
    right-handed circularly polarized light
11 mu_L = 1.53920;    // Refractive index of quartz for
    left-handed circularly polarized light
12 t = 5.0e-004;....// Thickness of the plate, m
13 theta = %pi*t*(mu_L-mu_R)/lambda;    // The angle
    of optical rotation, radian
14 [d,m] = deg2degmin(theta*180/%pi);    // Call the
    conversion function
15 printf("\nThe angle of rotation produced by its
    plate = %6.4f radians = %d degrees %d minutes",
    theta, d, m);
16
17 // Result
18 // The angle of rotation produced by its plate =
    0.1237 radians = 7 degrees 5 minutes

```

---

**Scilab code Exa 5.38** Optical rotation produced by new length of sugar solution

```

1 // Scilab Code Ex5.38: Page:315 (2011)
2 clc;clear;
3 theta = 13;....// Optical rotation of the solution,
    degrees
4 l = 20;....// Length of the tube, cm
5 l_prime = 30;....// New length of the tube, cm
6 c = 1;    // For simplicity assume concentration of
    sugar solution to be unity, g/cc
7 c_prime = c/3;    // New concentration of sugar
    solution, g/cc
8 // As,  $S = 10*\theta/(l*c)$  so  $10*\theta/(l*c) = 10*$ 
     $\theta\_prime/(l\_prime*c\_prime)$ 
9 // Solving for theta_prime
10 theta_prime = theta/(l*c)*l_prime*c_prime;    // The
    optical rotation produced by new length of sugar

```



```

    solution , degrees
11 printf("\nThe optical rotation of %d cm length of
    sugar solution = %3.1f degrees", l_prime,
    theta_prime);
12
13 // Result
14 // The optical rotation of 30 cm length of sugar
    solution = 6.5 degrees

```

---

**Scilab code Exa 5.39** Strength of the solution

```

1 // Scilab Code Ex5.39: Page:315 (2011)
2 clc;clear;
3 theta = 11;....// Optical rotation of sugar solution
    , degrees
4 l = 20;....// Length of the tube, cm
5 S = 66;....// Specific rotation of sugar solution ,
    degrees
6 c = theta*10/(l*S); // The concentration of sugar
    solution , g/cc
7 printf("\nThe strength of the solution = %6.4f g/cc"
    , c);
8
9 // Result
10 // The strength of the solution = 0.0833 g/cc

```

---

**Scilab code Exa 5.40** Length of sugar solution for given concentration and optical rotation

```

1 // Scilab Code Ex5.40: Page:316 (2011)
2 clc;clear;
3 theta = 20;....// Optical rotation of sugar solution
    , degrees

```

```
4 theta_prime = 35;....// New optical rotation of
   sugar solution , degrees
5 c = 5;....// Percentage concentration of the
   solution
6 c_prime = 10; // New percentage concentration of
   the solution
7 l = 1; // For simplicity assume length of the
   sugar solution to be unity
8 l_prime = theta_prime*l*c/(c_prime*theta);
9 printf("\nThe length of sugar solution for %d
   percent concentration and %d degrees optical
   rotation = %5.3f*l ", c_prime, theta_prime,
   l_prime);
10
11 // Result
12 // The length of sugar solution for 10 percent
   concentration and 35 degrees optical rotation =
   0.875*l
```

---

# Chapter 6

## X Rays

Scilab code Exa 6.1 Electrons striking the target in X ray coolidge tube

```
1 // Scilab Code Ex6.1: Page-369 (2011)
2 clc; clear;
3 i = 2e-003;....// Current through X-ray tube , A
4 e = 1.6e-019;....// Charge on an electron , C
5 V = 12.4e+003;....// Potential difference applied
   across X-ray tube , V
6 m0 = 9.1e-031;....// Rest mass of the electron , Kg
7 n = i/e; // Number of electrons striking the
   target per second
8 printf("\\nThe number of electrons striking the
   target per sec = %4.2e electrons", n);
9 v = sqrt(2*e*V/m0);....// Velocity of the electrons ,
   m/s
10 printf("\\nThe speed with which electrons strike the
   target = %4.2e m/s", v);
11
12 // Result
13 // The number of electrons striking the target per
   sec = 1.25e+016 electrons
14 // The speed with which electrons strike the target
   = 6.60e+007 m/s
```

---

**Scilab code Exa 6.2** Maximum speed of the electron striking the target

```
1 // Scilab Code Ex6.2: Page-370 (2011)
2 clc;clear;
3 e = 1.6e-019;....// Charge on an electron , C
4 V = 13.6e+003;....// Potential difference applied
   across X-ray tube , V
5 m0 = 9.1e-031;....// Rest mass of the electron , Kg
6 v = sqrt(2*e*V/m0);....// Velocity of the electron ,
   m/s
7 printf("\nThe maximum speed with which the electrons
   strike the target = %4.2e m/s", v);
8
9 // Result
10 // The maximum speed with which the electrons strike
   the target = 6.92e+007 m/s
```

---

**Scilab code Exa 6.3** Longest wavelength that can be analysed by a rock salt crystal

```
1 // Scilab Code Ex6.3: Page-370 (2011)
2 clc;clear;
3 d = 2.82e-010;....// Spacing of the rock-salt , m
4 n = 2;....// Order of diffraction
5 theta = %pi/2;    // Angle of diffraction , radian
6 // Braggs equation for X-rays of wavelength lambda
   is n*lambda = 2*d*sin(theta), solving for lambda
7 lambda = 2*d*sin(theta)/n;    // Wavelength of X-ray
   using Bragg's law , m
8 printf("\nThe longest wavelength that can be
   analysed by a rock-salt crystal = %4.2f angstrom"
   , lambda/1e-010);
```

```

9
10 // Result
11 // The longest wavelength that can be analysed by a
    rock-salt crystal = 2.82 angstrom

```

---

**Scilab code Exa 6.4** Angles at which the second and the third Bragg diffraction maxima are observed

```

1 // Scilab Code Ex6.4: Page-371 (2011)
2 clc;clear;
3 lambda = 3e-011;....// Wavelength of the X-ray, m
4 d = 5e-011;....// Lattice spacing, m
5 n = [2 3];....// Orders of diffraction
6 // Bragg's equation for X-rays of wavelength lambda
    is n*lambda = 2*d*sin(theta), solving for thetas
7 for i = 1:1:2
8 theta = asind(n(i)*lambda/(2*d));
9 printf("\nFor n = %d, theta = %4.1f degrees", n(i),
    theta);
10 end
11
12 // Result
13 // For n = 2, theta = 36.9 degrees
14 // For n = 3, theta = 64.2 degrees

```

---

**Scilab code Exa 6.5** Interplanar separation of atomic planes in the crystal

```

1 // Scilab Code Ex6.5: Page-371 (2011)
2 clc;clear;
3 lambda = 3.6e-011;....// Wavelength of X-rays, m
4 n = 1; // Order of diffraction
5 theta = 4.8; // Angle of diffraction, degrees

```

```

6 // Braggs equation for X-rays is  $n \cdot \lambda = 2 \cdot d \cdot \sin(\theta)$ 
   // solving for d
7 d = n*lambda/(2*sind(theta)); // Interplanar
   // spacing, m
8 printf("\nThe interplanar separation of atomic
   // planes in the crystal = %4.2f angstrom", d/1e
   // -010);
9
10 // Result
11 // The interplanar separation of atomic planes in
   // the crystal = 2.15 angstrom

```

---

**Scilab code Exa 6.6** Wavelength of K alpha radiation in copper for its given value in Mo

```

1 // Scilab Code Ex6.6: Page-371 (2011)
2 clc;clear;
3 lambda1 = 0.71;...// Wavelength of k alpha line in
   // molybdenum, angstrom
4 Z1 = 42; // Atomic number of Mo
5 Z2 = 29; // Atomic number of Cu
6 // Wavelength of characteristic X-ray for K-alpha
   // spectral line is given by
7 //  $1/\lambda = 3/4 \cdot R \cdot (Z-1)^2$  then
8 lambda2 = lambda1*(Z1-1)^2/(Z2-1)^2; // The
   // wavelength of K alpha radiation in copper, m
9 printf("\nThe wavelength of K-alpha radiation in
   // copper = %4.2f angstrom", lambda2);
10
11 // Result
12 // The wavelength of K-alpha radiation in copper =
   // 1.52 angstrom

```

---

**Scilab code Exa 6.7** Wavelength of gamma radiation at 90 degree

```
1 // Scilab Code Ex6.7: Page-372 (2011)
2 clc;clear;
3 phi = %pi/2; // Scattering angle, degrees
4 m0 = 9.1e-031;....// Rest mass of an electron, kg
5 h = 6.62e-034;....// Planck's constant, J-s
6 c = 3e+008;....// Speed of light in vacuum, m/s
7 E = 8.16e-014;....// Energy of gamma radiation, J
8 lambda = h*c/(E*1e-010); // Wavelength of
   incident photon, angstrom
9 lambda_prime = lambda+h*(1-cos(phi))/(m0*c*1e-010);
   // Wavelength of scattered photon, angstrom
10 printf("\nThe wavelength of radiation at 90 degrees
   = %6.4f angstrom", lambda_prime);
11
12 // Result
13 // The wavelength of radiation at 90 degrees =
   0.0486 angstrom
```

---

**Scilab code Exa 6.8** Compton shift from a carbon block

```
1 // Scilab Code Ex6.8: Page-372 (2011)
2 clc;clear;
3 phi = %pi/2; // Scattering angle, radian
4 m0 = 9.1e-031;....// Rest mass of the electron, kg
5 h = 6.62e-034;....// Planck's constant, J-s
6 c = 3e+008;....// Speed of light in vacuum, m/s
7 lambda = 1.00 ;....// Wavelength of incident photon,
   in angstrom
8 del_lambda = h*(1-cos(phi))/(m0*c*1e-010); //
   Compton shift, angstrom
9 printf("\nThe Compton shift = %6.4f angstrom",
   del_lambda);
10
```

```
11 // Result
12 // The Compton shift = 0.0242 angstrom
```

---

**Scilab code Exa 6.9** Wavelength of incident photon

```
1 // Scilab Code Ex6.9: Page-373 (2011)
2 clc;clear;
3 phi = %pi/2; // Scattering angle, radian
4 m0 = 9.1e-031;...// Rest mass of the electron, kg
5 h = 6.62e-034;...// Planck's constant, J-s
6 c = 3e+008;...// Speed of light in vacuum, m/s
7 // As Compton shift = del_lambda = lambda, so
8 lambda = h*(1-cos(phi))/(m0*c*1e-010); //
   Wavelength of incident photon, angstrom
9 printf("\nThe wavelength of incident radiation = %6
   .4f angstrom", lambda);
10
11 // Result
12 // The wavelength of incident radiation = 0.0242
   angstrom
```

---



# Chapter 7

## Lasers and Holography

Scilab code Exa 7.1 Energy of the laser pulse

```
1 // Scilab Code Ex 7.1: Page-411 (2011)
2 clc; clear;
3 e = 1.6e-019;....// Charge on an electron , eV
4 h = 6.62e-034;....// Planck's constant , J-s
5 c = 3e+008;....// Speed of light in vacuum, m/s
6 n = 2.8e+019;....// Number of photons in laser pulse
7 lambda = 7e-007;....// Wavelength of the radiation
   emitted by the laser , m
8 E = h*c/(lambda*e);....// Energy of the photon in
   the laser light , eV
9 del_E = E*n;....// The energy of laser pulse having
   n photons , eV
10 printf("\\nThe energy of the laser pulse = %4.2e eV",
   del_E);
11
12 // Result
13 // The energy of the laser pulse = 4.97e+019 eV
```

---

**Scilab code Exa 7.2** Coherence length resultant bandwidth and line width of laser beam

```
1 // Scilab Code Ex7.2: Page-411 (2011)
2 clc;clear;
3 c = 3e+008;....// Speed of light in vacuum, m/s
4 lambda = 6.5e-007;....// Wavelength of the pulse, m
5 t = 0.5e-009;....// Time interval between successive
    pulses, s
6 L_c = c*t;....// Coherence length of laser pulse, m
7 printf("\nThe coherence length of the pulse = %4.2f
    m", L_c);
8 del_nu = 1/t;....// Resultant bandwidth of laser
    pulse, Hz
9 printf("\nThe bandwidth of the laser pulse = %1.0e
    Hz", del_nu);
10 del_lambda = lambda^2*del_nu/c;....// Linewidth of
    laser beam, m
11 printf("\nThe linewidth of the pulse = %5.3f
    angstrom", del_lambda/1e-010);
12
13 // Result
14 // The coherence length of the pulse = 0.15 m
15 // The bandwidth of the laser pulse = 2e+009 Hz
16 // The linewidth of the pulse = 0.028 angstrom
```

---

**Scilab code Exa 7.3** Angular spread and areal spread of laser beam

```
1 // Scilab Code Ex7.3: Page-411 (2011)
2 clc;clear;
3 a = 4e-003;....// Coherence width of laser source, m
4 lambda = 6e-007;....// Wavelength of the pulse, m
5 D = 100;....// Distance of the surface from laser
    source, m
6 A = 2*lambda/a; // Angular spread of laser beam,
```

```
        radian
7 printf("\nThe angular spread = %1.0e radian", A);
8 theta = A/2;    // Semi angle, radian
9 A_s = %pi*(D*theta)^2;....// Areal spread of laser
    beam, Sq.m
10 printf("\nThe areal spread = %1.0e Sq.m", A_s);
11
12 // Result
13 // The angular spread = 3e-004 radian
14 // The areal spread = 7e-004 Sq.m
```

---

# Chapter 8

## Ultrasonics

**Scilab code Exa 8.1** Frequency of the fundamental mode of ultrasonic wave

```
1 // Scilab Code Ex8.1: Page-429 (2011)
2 clc; clear;
3 d = 8e-004;....// Thickness of the piece of
   piezoelectric crystal, m
4 v = 5760;....// Velocity of ultrasonic waves in the
   piece of piezoelectric crystal, m/s
5 n = v/(2*d); // The frequency of the fundamental
   mode of ultrasonic wave, Hz
6 printf("\nThe frequency of the fundamental mode of
   ultrasonic wave = %3.1f MHz", n/1e+006);
7
8 4// Result
9 // The frequency of the fundamental mode of
   ultrasonic wave = 3.6 MHz
```

---

**Scilab code Exa 8.2** Fundamental frequency of quartz crystal

```
1 // Scilab Code Ex8.2: Page-430 (2011)
```

```

2  clc;clear;
3  d = 2e-003;....// Thickness of the piece of quartz
    crystal, m
4  rho = 2650;....// Density of the crystal, kg/meter-
    cube
5  Y = 7.9e+010;....// Value of Youngs Modulus, N/metre
    -square
6  n = 1/(2*d)*sqrt(Y/rho);    //The frequency of the
    fundamental mode of vibration, Hz
7  printf("\\nThe frequency of the fundamental mode of
    vibration in quatrz crystal = %5.3f Hz", n/1e
    +006);
8
9  // Result
10 // The frequency of the fundamental mode of
    vibration in quatrz crystal = 1.365 Hz

```

---

**Scilab code Exa 8.3** Thickness of steel plate using ultrasonic beam

```

1  // Scilab Code Ex8.3: Page-430 (2011)
2  clc;clear;
3  v = 5e+003;....// Velocity of ultrasonic beam in
    steel plate, m/s
4  n = 25e+003;....// Difference between two
    neighbouring harmonic frequencies (Nm - Nm_minus1
    ), Hz
5  d = v/(2*n);    // The thickness of steel plate, m
6  printf("\\nThe thickness of steel plate = %3.1f m", d
    );
7
8  // Result
9  // The thickness of steel plate = 0.1 m

```

---

**Scilab code Exa 8.4** Inductance of an inductor to produce ultrasonic waves

```
1 // Scilab Code Ex8.4: Page-430 (2011)
2 clc;clear;
3 n = 1e+006;....// Frequency of Ultrasonic waves, Hz
4 C = 2.5e-014;....// Capacitance of capacitor, F
5 // Frequency of electric oscillations is given by n =
   1/(2*%pi)*sqrt(1/(L*C)), solving for L
6 L = 1/(4*%pi^2*n^2*C); // The inductance of an
   inductor to produce ultrasonic waves, henry
7 printf("\\nThe inductance of an inductor to produce
   ultrasonic waves = %d henry", L);
8
9 // Result
10 // The inductance of an inductor to produce
   ultrasonic waves = 1 henry
```

---

**Scilab code Exa 8.5** Position of imperfection and the velocity of pulse inside the rod

```
1 // Scilab Code Ex8.5: Page-431 (2011)
2 clc;clear;
3 d = 50e-002;....// Thickness of the metallic rod, m
4 t1 = 30e-006;....// Arrival time for first pulse, s
5 t2 = 80e-006;.... // Arrival time for second
   pulse, s
6 v = 2*d/t2;....// Velocity of ultrasonic waves, m/s
7 printf("\\nThe velocity of pulse inside the rod = %4
   .2e m/s", v);
8 x = t1*v/2;
9 printf("\\nThe position of pulse inside the rod = %6
   .4f m", x);
10
11 // Result
12 // The velocity of pulse inside the rod = 1.25e+004
```

```
13 // The position of pulse inside the rod = 0.1875 m
    m/s
```

---

**Scilab code Exa 8.6** Maximum acceleration and displacement of a quartz ultrasonic transducer

```
1 // Scilab Code Ex8.6: Page-431 (2011)
2 clc;clear;
3 I = 2.5e+004;....// Sound intensity , W/meter-square
4 v = 1480;....// Sound velocity , m/s
5 rho_w = 1000;....// Density of water , kg/meter-cube
6 rho_c = 2650;....// Density of crystal of transducer
    , kg/meter-cube
7 d = 0.001;....// Thickness of the quartz , m
8 f = 20e+003;....// Frequency of sound in water , Hz
9 // As sound intensity ,  $I = p^2/(2*\rho_0*v)$ , solving
    for p
10 p = sqrt(2*rho_w*v*I); // Pressure in the medium,
    N/metre-square
11 a = p/(d*rho_c); // Maximum acceleration of the
    quartz ultrasonic transducer , metre/second-square
12 printf("\\nThe maximum acceleration produced in
    quartz transducer = %4.2e metre/second-square", a
    );
13 y = a/(2*%pi*f)^2; // Maximum displacement of the
    quartz transducer , m
14 printf("\\nThe maximum displacement of quartz
    transducer = %3.1f micron", y/1e-006);
15
16 // Result
17 // The maximum acceleration produced in quartz
    transducer = 1.03e+005 metre/second-square
18 // The maximum displacement of quartz transducer =
    6.5 micron
```

---

**Scilab code Exa 8.7** Fundamental frequency of a magnetostrictive hydrophone

```
1 // Scilab Code Ex8.7: Page-432 (2011)
2 clc;clear;
3 L = 0.2;....// Length of a magnetostrictive
   hydrophone, m
4 lambda = 2*L;....// Wavelength of ultrasonic wave, m
5 v = 4900;....// Velocity of ultrasonic beam in water
   , m/s
6 f = v/lambda;....// Fundamental frequency of
   ultrasonic, KHz
7 printf("\nThe fundamental frequency of a
   magnetostrictive hydrophone = %4.2f KHz", f/1e
   +03);
8
9 // Result
10 // The fundamental frequency of a magnetostrictive
   hydrophone = 12.25 KHz
```

---

**Scilab code Exa 8.8** Length of the copper wire used to introduce ultrasonic delay

```
1 // Scilab Code Ex8.8: Page-432 (2011)
2 clc;clear;
3 v = 3700;....// Velocity of ultrasonic beam in
   copper, m/s
4 t = 1e-006;....// Delay time for ultrasonic beam, s
5 L = v*t; // // Length of a copper wire required
   for a delay, m
6 printf("\nThe length of a copper wire required for a
   delay = %6.4f m", L);
7
```



```
8 // Result
9 // The length of a copper wire required for a delay
   = 0.0037 m
```

---

# Chapter 9

## Fibre Optics

Scilab code Exa 9.1 NA and the acceptance angle of optical fibre

```
1 // Scilab Code Ex9.1: Page-463 (2011)
2 clc;clear;
3 mu_1 = 1.55;....// Refractive index of the core
4 mu_2 = 1.50;....// Refractive indices of cladding
5 NA = mu_1*sqrt(2*(mu_1-mu_2)/mu_1);
6 printf("\nThe NA of the optical fibre = %5.3f", NA);
7 theta_a = asind(NA); // The acceptance angle of
   optical fibre , degrees
8 printf("\nThe acceptance angle of the optical fibre
   is = %4.1f degrees", theta_a);
9
10 // Result
11 // The NA of the optical fibre = 0.394
12 // The acceptance angle of the optical fibre is =
   23.2 degrees
```

---

Scilab code Exa 9.2 NA acceptance angle and the critical angle of optical fibre

```

1 // Scilab Code Ex9.2: Page-463 (2011)
2 clc;clear;
3 mu_1 = 1.50;....// Refractive index of the core
4 mu_2 = 1.45;....// Refractive index cladding
5 NA = mu_1*sqrt(2*(mu_1-mu_2)/mu_1); // Numerical
    aperture of optical fibre
6 printf("\n The NA of the optical fibre = %5.3f", NA)
    ;
7 theta_a = asind(NA); // The acceptance angle
    of optical fibre , degrees
8 printf("\n The acceptance angle of the optical fibre
    = %5.2f degrees", theta_a);
9 theta_c = asind(mu_2/mu_1); // The critical angle of
    the optical fibre , degrees
10 printf("\n The acceptance angle of the optical fibre
    = %4.1f degrees", theta_c);
11
12 // Result
13 // The NA of the optical fibre = 0.387
14 // The acceptance angle of the optical fibre = 22.8
    degrees
15 // The acceptance angle of the optical fibre = 75.2
    degrees

```

---

### Scilab code Exa 9.3 Acceptance angle for the optical fibre in water

```

1 // Scilab Code Ex9.3: Page-464 (2011)
2 clc;clear;
3 mu0 = 1;....// Refractive index of fibre in air
4 mu2 = 1.59;....// Refractive index of the cladding
5 NA = 0.2;....// Numerial aperture of optical fibre
6 mu1 = sqrt(NA^2+mu2^2); // Refractive index of core
7 mu0 = 1.33; // Refractive index of fibre in water
8 NA = sqrt(mu1^2-mu2^2)/mu0; // Numerial aperture
    of optical fibre in water

```

```

9 theta_a = asind(NA);    // Acceptance angle for the
   fibre in water
10 printf("\nThe acceptance angle for the optical fibre
   in water = %3.1f degrees", theta_a);
11
12 // Result
13 // The acceptance angle for the optical fibre in
   water = 8.6 degrees

```

---

#### Scilab code Exa 9.4 The characteristics of glass clad fibre

```

1 // Scilab Code Ex9.4: Page-464 (2011)
2 clc;clear;
3 mu0 = 1;           // Refractive index of air
4 mu1 = 1.50;.... // Refractive index of glass core '
5 del = 0.005;.... // Fractional change in refractive
   index
6 mu2 = mu1*(1-del); // Refractive index of
   cladding
7 printf("\nThe refractive index of cladding =%6.4f",
   mu2);
8 theta_c = asind(mu2/mu1); // Critical angle ,
   degrees
9 printf("\nThe critical angle = %5.2f degrees",
   theta_c);
10 theta_a = asind(sqrt(mu1^2-mu2^2)/mu0); //
   Acceptance angle , degrees
11 printf("\nThe value of acceptance angle is = %4.2f
   degrees", theta_a);
12 NA = mu1*sqrt(2*del); // Numerical aperture of
   optical fibre
13 printf("\nThe NA of the optical fibre = %4.2f", NA);
14
15 // Result
16 // The refractive index of cladding =1.4925

```

```

17 // The critical angle = 84.27 degrees
18 // The value of acceptance angle is = 8.62 degrees
19 // The NA of the optical fibre = 0.15

```

---

**Scilab code Exa 9.5** Refractive index of core and cladding of an optical fibre

```

1 // Scilab Code Ex9.5: Page-465 (2011)
2 clc; clear;
3 NA = 0.22; // Numerical aperture of the optical
  fibre
4 del = 0.012; ... // Fractional difference between the
  refractive index of core and cladding
5 mu1 = NA/sqrt(2*del); // The refractive index of
  core of optical fibre
6 printf("\\nThe refractive index of core = %4.2f", mu1
  );
7 mu2 = mu1*(1-del); // The refractive index of
  cladding of optical fibre
8 printf("\\nThe refractive index of cladding = %4.2f",
  mu2);
9
10 // Result
11 // he refractive index of core = 1.42
12 // The refractive index of cladding = 1.40

```

---

**Scilab code Exa 9.6** NA and the core radius of an optical fibre

```

1 // Scilab Code Ex9.6: Page-466 (2011)
2 clc; clear;
3 mu1 = 1.466; // Refractive index of core
4 mu2 = 1.460; // Refractive index of cladding

```

```

5 v = 2.4;....// Cut-off parameter of the optical
  fibre
6 lambda = 0.8e-006;....// Operating wavelength , m
7 NA = sqrt(mu1^2-mu2^2);
8 printf("\nThe NA of optical fibre = %4.2f", NA) ;
9 // Asthe cut-off parameter v of the optical fibre , v
  = 2*%pi*a*sqrt(mu1^2-mu2^2)/lambda, solving for
  a
10 a = lambda*v/(2*%pi*sqrt(mu1^2-mu2^2));
11 printf("\nThe core radius of the optical fibre = %4
  .2e micron", a/1e-006);
12
13 // Result
14 // The NA of optical fibre = 0.13
15 // The core radius of the optical fibre = 2.31e+00
  micron

```

---

**Scilab code Exa 9.7** v number and the number of modes supported by the optical fibre

```

1 // Scilab Code Ex9.7: Page-466 (2011)
2 clc;clear;
3 mu1 = 1.54; // The refractive index of core
4 mu2 = 1.50; // The refractive index of cladding
5 lambda = 1.3e-006;....// Operating wavelength of
  optical fibre , m
6 a = 25e-006;....// Radius of fibre core , m
7 v = 2*%pi*a*sqrt(mu1^2-mu2^2)/lambda; // V-number
  of optical fibre
8 printf("\nThe cut-off parameter of the optical fibre
  = %5.2f", v);
9 n = v^2/2; // The number of modes supported by
  the fibre
10 printf("\nThe number of modes supported by the fibre
  = %3d", ceil(n));

```

```

11
12 // Result
13 // The cut-off parameter of the optical fibre =
    42.14
14 // The number of modes supported by the fibre = 888

```

---

**Scilab code Exa 9.8** Maximum values of refractive index of cladding and the fractional change in refractive index

```

1 // Scilab Code Ex9.8: Page-466 (2011)
2 clc; clear;
3 mu1 = 1.54;....// Refractive index of core
4 v = 2.405;....// Cut-off parameter of optical fibre
5 lambda = 1.3e-006;....// Operating wavelength of
    optical fibre , m
6 a = 1e-006;....// Radius of the core ,
7 NA = v*lambda/(2*%pi*a);    // Numerical aperture of
    optical fibre
8 del = 1/2*(NA/mu1)^2;    // Fractional change in
    refractive index of core and cladding
9 printf("\\nThe fractional difference of refractive
    indices of core and cladding = %7.5f", del);
10 mu2 = mu1*(1-del);    // Maximum value of
    refractive index of cladding
11 printf("\\nThe maximum refractive index of cladding =
    %5.3f", mu2);
12
13 // Result
14 // The fractional difference of refractive indices
    of core and cladding = 0.05220
15 // The maximum refractive index of cladding = 1.460

```

---

**Scilab code Exa 9.9** Normalized frequency for the optical fibre

```

1 // Scilab Code Ex9.9: Page-467 (2011)
2 clc;clear;
3 mu1 = 1.45;....// Index of refraction of core
4 NA = 0.16;....// Numerical aperture of step index
   fibre
5 a = 3e-006;....// Radius of the core, m
6 lambda = 0.9e-006;....// Operating wavelength of
   optical fibre, m
7 v = 2*%pi*a*NA/lambda;      // The normalized
   frequency or v-number of optical fibre
8 printf("\\nThe normalized frequency of the optical
   fibre = %5.2f", v);
9
10 // Result
11 // The normalized frequency of the optical fibre =
   3.35

```

---

**Scilab code Exa 9.10** Cut off parameter or v number of modes supported by the fibre

```

1 // Scilab Code Ex9.10: Page-467 (2011)
2 clc;clear;
3 mu1 = 1.52;....// Refractive index of core
4 a = 14.5e-006;....// Radius of the fibre core, m
5 del = 0.0007;....// Fractional index difference
6 lambda = 1.3e-006;....// Operating wavelength of
   optical fibre, m
7 mu2 = mu1*(1-del);      // Refractive index of
   cladding
8 v = 2*%pi*a*sqrt(mu1^2-mu2^2)/lambda;      // Cut-off
   parameter v of the optical fibre
9 printf("\\nThe cut-off parameter of the optical fibre
   = %5.3f", v);
10 //The is number of modes supported by the fibre
   given by,

```



```

11 n = v^2/2;
12 printf("\nThe number of modes supported by the fibre
    = %d", ceil(n));
13
14 // Result
15 // The cut-off parameter of the optical fibre =
    3.985
16 // The number of modes supported by the fibre = 8

```

---

**Scilab code Exa 9.11** Power output through optical fibre

```

1 // Scilab Code Ex9.11: Page-468 (2011)
2 clc;clear;
3 alpha = 3.5;....// Attenuation of the optical fibre ,
    dB/km
4 Pi = 0.5;....// Input power of optical fibre , mW
5 L = 4;.... // Distance through the optical wave
    transmits through the fibre , km
6 // As alpha = 10/L*log10(Pi/Po), solving for Po
7 Po = Pi/exp(alpha*L*2.3026/10); // Output power of
    optical fibre , mW
8 printf("\nThe output power of optical fibre = %4.1f
    micro-watt", Po/1e-003);
9
10 // Result
11 // The output power of optical fibre = 19.9 micro-
    watt

```

---

**Scilab code Exa 9.12** Attenuation of power through optical fibre

```

1 // Scilab Code Ex9.12: Page-468 (2011)
2 clc;clear;
3 Pi =1;....// Input power of optical fibre , mW

```

```

4 Po = 0.85;....// Outptu power of optical fibre , mW
5 L = 0.5;....//The distance through the optical wave
   transmits through the fibre , km
6 alpha = (10/L)*log10(Pi/Po);      // The attenuation
   of power through the optical fibre
7 printf("\nThe attenuation of power through the
   optical fibre = %5.3f dB/km", alpha);
8
9 // Result
10 // The attenuation of power through the optical
   fibre = 1.412 dB/km

```

---

**Scilab code Exa 9.13** Minimum optical power input to an optical fibre

```

1 // Scilab Code Ex9.13: Page-469 (2011)
2 clc;clear;
3 C = 0.8;      // Connector loss per km, dB
4 F = 1.5;      // Fibre loss per km, dB
5 alpha = C + F;....// Attenuation of power the
   optical fibre , dB/km
6 Po = 0.3e-006;....// Output power of optical fibre ,
   W
7 L = 15;....// The distance through the optical wave
   transmits through the fibre , km
8 //As the attenuation , alpha = 10/L*log(Pi/Po) ,
   solving for Pi
9 Pi = Po*exp(2.3026*alpha*L/10);      // Input power
   of optical fibre , mW
10 printf("\nThe minimum input power to optical fibre =
   %5.3f mW", Pi/1e-003);
11
12 // Result
13 // The minimum input power to optical fibre = 0.846
   mW

```

---

# Chapter 10

## Electrostatics

**Scilab code Exa 10.1** Potential difference between the two charged horizontal plates

```
1 // Scilab Code Ex10.1: Page-507 (2011)
2 clc;clear;
3 m = 4e-013;....// Mass of the particle , kg
4 q = 2.4e-019;....// Charge on particle , C
5 d = 2e-002;....// Distance between the two
   horizontally charged plates , m
6 g = 9.8;....// Acceleration due to gravity , m/sec-
   square
7 E = m*g/q ;....// Electric field strength , N/C
8 V = E*d;.....// Potential difference between the two
   charged horizontal plates , V
9 printf("\nThe potential difference between the two
   horizontally charged plates = %3.1e V", V);
10
11 // Result
12 // The potential difference between the two
   horizontally charged plates = 3.3e+005 V
```

---

**Scilab code Exa 10.2** Electric potential at a point equidistant from the three corners of a triangle

```
1 // Scilab Code Ex10.2: Page-507 (2011)
2 clc;clear;
3 q1 = 1e-009; // Charge at first corner, C
4 q2 = 2e-009; // Charge at second corner, C
5 q3 = 3e-009; // Charge at third corner, C
6 d = 1;....// Side of the equilateral triangle, m
7 theta = 30;....// Angle at which line joining the
   observation point to the source charge makes with
   the side, degrees
8 r = (d/2)/cosd(theta);....// Distance of observation
   point from the charges, m
9 //since,1/4*%pi*%eps = 9e+009;
10 V = (q1+q2+q3)*(9e+009)/r;.....// Electric potential,
   V
11 printf("\nThe electric potential at the point
   equidistant from the three corners of the
   triangle = %4.1f V", V);
12
13 // Result
14 // The electric potential at the point equidistant
   from the three corners of the triangle = 93.5 V
```

---

**Scilab code Exa 10.3** Electric potential at the centre of a square

```
1 // Scilab Code Ex10.3: Page-507 (2011)
2 clc;clear;
3 q = 2e-008;
4 q1 = q; // Charge at first corner, C
5 q2 = -2*q; // Charge at second corner, C
6 q3 = 3*q; // Charge at third corner, C
7 q4 = 2*q; // Charge at fourth corner, C
8 d = 1;.... // Side of the square, m
```

```

9 r = d*sin(2*%pi/8);....// Distance of centre of the
    square from each corner, m
10 V = (q1+q2+q3+q4)*(9e+009)/r;.....// Electric
    potential at the centre of the square, V
11 printf("\nThe electric potential at the centre of
    the square = %4d V", V);
12
13 // Result
14 // The electric potential at the centre of square =
    1018 V

```

---

**Scilab code Exa 10.6** New potential when the two charged drops coalesce to form a bigger drop

```

1 // Scilab Code Ex10.6: Page-512 (2011)
2 clc;clear;
3 V = 60;....// Electric potential of smaller drop,
    volt
4 r = 1;....// For simplicity assume radius of each
    small drop to be unity, unit
5 q = 1;....// For simplicity assume charge on smaller
    drop to be unity, C
6 k = 1;....// For simplicity assume Coulomb's
    constant to be unity, unit
7 R = 2^(1/3)*r;.....// Radius of bigger drop, unit
8 Q = 2*q;....// Charge on bigger drop, C
9 V_prime = k*Q/R*V;....// Electric potential of
    bigger drop, volt
10 printf("\nThe electric potential of new drop = %4.1 f
    V", V_prime);
11
12 // Result
13 // The electric potential of new drop = 95.2 V

```

---

**Scilab code Exa 10.7** Magnitude and the direction of electric field which would balance the weight of an electron placed in it

```
1 // Scilab Code Ex10.7: Page-512 (2011)
2 clc;clear;
3 m = 9.1e-31;....// Mass of the electron , kg
4 e = 1.6e-19;....// Charge on an electron , C
5 g = 9.8;....// Acceleration due to gravity , m/sec-
    square
6 // Electric force ,  $F = e \cdot E$ , where  $F = m \cdot g$  or  $e \cdot E = m$ 
    *g
7 E = m*g/e; // Electric field which would balance
    the weight of an electron placed in it , N/C
8 printf("\nThe required electric field strength = %3
    .1e N/C", E);
9 printf("\nThis field acts opposite to the direction
    of weight");
10
11 // Result
12 // The required electric field strength = 5.6e-011 N
    /C
13 // This field acts opposite to the direction of
    weight
```

---

**Scilab code Exa 10.8** Magnitude and the direction of electric field at a point midway between two charges

```
1 // Scilab Code Ex10.8: Page-512 (2011)
2 clc;clear;
3 q1 = 8e-007;....// First Charge , C
4 q2 = -8e-007;....// Second Charge , C
```

```

5 r = 15e-002;....// Distance between the two charges ,
  m
6 k = 9e+009; // Coulomb's constant , N-metre-square/
  coulomb-square
7 E1 = k*q1/r^2;....// Electric field strength due to
  charge 8e-007 C
8 printf("\nThe electric field strength at midpoint =
  %3.1e N/C", E1);
9 E2 = abs(k*q2/r^2);....// Electric field strength -8
  e-007 C
10 printf("\nThe electric field strength at midpoint =
  %3.1e N/C", E2);
11 // Total electric field strength at the mid-point is
12 E = E1+E2; // Net electric field at mid point ,
  N/C
13 printf("\nThe net electric field strength at
  midpoint = %3.1e N/C", E);
14
15 // Result
16 // The electric field strength at midpoint = 3.2e+05
  N/C
17 // The electric field strength at midpoint = 3.2e+05
  N/C
18 // The net electric field strength at midpoint = 6.4
  e+05 N/C

```

---

#### Scilab code Exa 10.9 Electric field strength at a point

```

1 // Scilab Code Ex10.9: Page:513 (2011)
2 clc;clear;
3 x = poly(0, 'x');
4 V = 1000/x+1500/x^2+500/x^3; // Given electric
  potential at a point (x,0,0), V
5 E = -1*derivat(V); // Electric field at a point as
  gradient of scalar potential, N/C

```

```

6 E_x = horner(E, 1);      // Electric field at the
    point x = 1, N/C
7 printf("\nThe electric field strength at point x = 1
    is %4di V/m", E_x);
8
9 // Result
10 // The electric field strength at point x = 1 is
    5500i V/m

```

---

**Scilab code Exa 10.11** Electric field strength due to spherical charge distribution

```

1 // Scilab Code Ex10.11: Page:514 (2011)
2 clc;clear;
3 function e = E(r)
4     a = 1;      // For convenience assume radius of
    sphere to be unity
5     r = poly(0, 'r');
6     e = r/3-r^3/(5*a^2);
7 endfunction
8
9 rho_0 = 1;     // For convenience assume charge
    density to be unity
10 epsilon_0 = 1; // For convenience assume
    permittivity to be unity
11 r = poly(0, 'r');
12 E_int = rho_0/epsilon_0*E('r');
13 delta_E = derivat(E_int);
14 r = roots(delta_E);
15 printf("\nThe electric field strength is maximum at
    an internal point at a distance r = sqrt(%g)a/3
    from the centre", (3*r(1))^2);
16
17 // Result
18 // The electric field strength is maximum at an

```



internal point at a distance  $r = \sqrt{5}a/3$  from  
the centre

---

**Scilab code Exa 10.12** Maximum electric field strength at an internal point

```
1 // Scilab Code Ex10.12: Page:517 (2011)
2 clc;clear;
3 function e = E(r)
4     a = 1; // For convenience assume radius of
5           sphere to be unity
6     r = poly(0, 'r');
7     e = r/3-r^2/(4*a);
8 endfunction
9 rho_0 = 1; // For convenience assume charge
10          density to be unity
11 epsilon_0 = 1; // For convenience assume
12              permittivity to be unity
13 r = poly(0, 'r');
14 E_int = rho_0/epsilon_0*E('r');
15 delta_E = derivat(E_int);
16 r = roots(delta_E);
17 printf("\nThe electric field strength is maximum at
18        an internal point at a distance r = %da/3 from
19        the centre", 3*r);
20
21 // Result
22 // The electric field strength is maximum at an
23 // internal point at a distance r = 2a/3 from the
24 // centre
```

---

# Chapter 11

## Electromagnetic Theory

Scilab code Exa 11.1 Amplitude of field vector E in free space

```
1 // Scilab Code Ex11.1: Page-559 (2011)
2 clc; clear;
3 H_0 = 1;....// Amplitude off field vector ,in A/m
4 mu_0 = 12.56e-7;....// Permeability ,in weber/A-m
5 eps = 8.85e-12;....// Permittivity in free space ,in
   C/N-meter-square
6 // From the relation between the amplitude of the
   field vector E and vector H of an EM wave in free
   space
7 E_0 = H_0*(sqrt(mu_0/eps));
8 printf("\\nThe amplitude of field vector E in free
   space = %5.1f V/m" ,E_0);
9
10
11 // Result
12 // The amplitude of field vector E in free space =
   376.7 V/m
```

---

Scilab code Exa 11.2 Maximum value of magnetic induction vector

```

1 // Scilab Code Ex11.2: Page-560 (2011)
2 clc;clear;
3 E_o = 1e+3;....// Amplitude field vector in free
   space,N/C
4 c = 3e+8;....// Speed of light ,in m/s
5 // From the relation between the amplitude of the
   field vector E and vector H of an EM wave in free
   space E_o = H_o*(sqrt(mu_o/eps))and B_o = mu_o*
   H_o, we have
6 B_o = E_o/c;
7 printf("\nThe maximum value of magnetic induction
   vector = %4.2e weber/A-m",B_o);
8
9 // Result
10 // The maximum value of magnetic induction vector =
   3.33e-006 weber/A-m

```

---

**Scilab code Exa 11.3** Conduction and displacement current densities in the conducting medium

```

1 // Scilab Code Ex11.3: Page-560 (2011)
2 clc;clear;
3 sigma = 5;....// Conductivity of the conducting
   medium, mho/m
4 eps_r = 8.85e-12;....// Relative electrical
   permittivity of medium, F/m
5 eps_0 = 1; // Electrical permittivity of free
   space, F/m
6 E0 = 250; // Amplitude of applied electric field,
   V/m
7 J = sigma*E0; // Amplitude of conduction current
   density, A/metre-square
8 J_D = eps_r*eps_0*E0*1e+010; // Amplitude of
   displacement current density, A/metre-square
9 omega = sigma/(eps_0*eps_r); // Frequency at

```

```

    which J = J_D
10 printf("\nThe conduction current density = %3dsin
    (10^10t) A/metre-square", J);
11 printf("\nThe displacement current density = %5.3
    fcos(10^10t) A/metre-square", J_D);
12 printf("\nThe frequency at which J = J_D is %3.1e Hz
    ", omega);
13
14 // Result
15 // The conduction current density = 1250 sin(10^10t)
    A/metre-square
16 // The displacement current density = 22.125 cos
    (10^10t) A/metre-square
17 // The frequency at which J = J_D is 5.6e+11 Hz

```

---

**Scilab code Exa 11.8** Average value of the intensity of electric field of radiation

```

1 // Scilab Code Ex11.8 :Page-565 (2011)
2 clc;clear;
3 P = 1000;....// Energy radiated by the lamp, watt
4 r = 2;....// Distance from the source at which the
    electric field intensity is given, m
5 S = P/(4*pi*r^2); // Magnitude of Poynting
    vector, W/metre-square
6 // As wave impedance, Z0 = E/H = 377 and H = E/377,
    so that with E*H = S we have
7 E = poly(0, 'E');
8 E = roots(E*E/377-S);
9 printf("\nThe average value of the intensity of
    electric field of radiation = %4.1f V/m", E(2));
10
11 // Result
12 // The average value of the intensity of electric
    field of radiation = 86.6 V/m

```

---

**Scilab code Exa 11.9** Amplitude of electric and magnetic fields of radiation

```
1 // Scilab Code Ex11.9: Page-566 (2011)
2 clc;clear;
3 S = 2*4.186/60*1e+04;....// Solar constant , J/s/
    metre-square
4 // From the poynting vector S = E*H
5 C = 377;....// Wave Impedence , ohm
6 E = sqrt(S*C);      // Electric field of radiation ,
    V/m
7 H = E/C;           // Magnetic field of radiation ,
    A/m
8 E0 = E*sqrt(2);    // Amplitude of electric field
    of radiation , V/m
9 H0 = H*sqrt(2);    // Amplitude of magnetic field
    of radiation , A/m
10 printf("\nThe amplitude of electric field of
    radiation = %6.1f V/m", E0);
11 printf("\nThe amplitude of magnetic field of
    radiation = %5.3f V/m", H0);
12
13
14 // Result
15 // The amplitude of electric field of radiation =
    1025.7 V/m
16 // The amplitude of magnetic field of radiation =
    2.721 V/m
```

---

**Scilab code Exa 11.10** Phase difference between electric and magnetic field vectors of an EM wave

```

1 // Scilab Code Ex 11.10: Page-568 (2011)
2 clc;clear;
3 function s = sine(x)
4     s = x - x^3/factorial(3) + x^5/factorial(5) - x
        ^7/factorial(7) + x^9/factorial(9);
5 endfunction
6
7 function s = cosine(x)
8     s = 1 - x^2/factorial(2) + x^4/factorial(4) - x
        ^6/factorial(6) + x^8/factorial(8);
9 endfunction
10
11 k = 1; // For simplicity assume constant of
        proportionality to be unity, units
12 for theta = 1:1:45
13     alpha = k*cosd(theta);
14     b = k*sind(theta);
15     if alpha == b then
16         phi = atand(b/alpha);
17         break;
18     end
19 end
20 //printf("\nThe phase difference between electric
        and magnetic field vectors = %4.2f rad", phi);
21
22
23 // Result
24 // The skin depth of an EM-wave in Al = 0.000010 m

```

---

**Scilab code Exa 11.12** Skin depth of an EM wave in Al

```

1 // Scilab Code Ex11.12: Page-569 (2011)
2 clc;clear;
3 sigma = 3.54e+007;....// Electrical conductivity of
        Al, mho per metre

```

```

4 mu = 12.56e-007;....// Permeability of the medium,
   weber/A-m
5 f = 71.6e+06;      // Frequency of the wave, Hz
6 omega = 2*pi*f;....// Angular frequency of the wave
   , rad per sec
7 delta = sqrt(2/(sigma*mu*omega));    // Skin depth
   of the EM wave in Al, m
8 printf("\nThe skin depth of an EM-wave in Al = %2.0 f
   micron", delta/1e-06);
9
10 // Result
11 // The skin depth of an EM-wave in Al = 10 micron

```

---

**Scilab code Exa 11.14** Skin depth and attenuation constant of sea water

```

1 // Scilab Code Ex11.14: Page-571 (2011)
2 clc;clear;
3 sigma = 5;....// Electrical conductivity, mho per
   metre
4 mu = 12.56e-007;....// Permeability of the medium,
   weber/A-m
5 eps_0 = 8.85e-012;....// Electric permittivity of
   free space, C-square/N-m-square
6 eps = 70*eps_0;    // Electric permittivity of the
   medium, C-square/N-m-square
7 delta = 2/sigma*sqrt(eps/mu); // The skin depth and
   attenuation constant of sea water
8 printf("\nThe skin depth of an EM-wave in sea water
   = %6.4 f m", delta);
9 Beta = 1/delta;    // The attenuation constant of
   sea water, per metre
10 printf("\nThe attenuation constant of the sea water
   = %6.2 f m", Beta);
11
12 // Result

```

- 13 // The skin depth of an EM-wave in sea water =  
0.0089 m
- 14 // The attenuation constant of the sea water =  
112.57 m
-



## Chapter 12

# Magnetic Properties of Materials

Scilab code Exa 12.1 Current through the solenoid

```
1 // Scilab Code Ex12.1 Page-603 (2011)
2 clc; clear;
3 H = 5e+3;....// Coercivity of a bar magnet, A/m
4 L = 0.1;....// Length of the solenoid, m
5 N = 50;....// Turns in solenoid
6 n = 500;....// Turns/m
7 // Using the relation
8 I = H/n;....// where I is the current through the
   solenoid
9 printf("\n\nThe current through the solenoid is = %2d
   A", I);
10
11 // Result
12 // The current through the solenoid is = 10 A
```

---

Scilab code Exa 12.2 Magnetic moment of the iron rod

```

1 // Scilab Code Ex12.2 : Page-603 (2011)
2 clc; clear;
3 n = 500;....// Number of turns wound per metre on
   the solenoid
4 i = 0.5;....// Current through the solenoid , A
5 V = 1e-03;....// Volume of iron rod , per metre cube
6 mu_r = 1200;    // Relative permeability of the iron
7 H = n*i;    // Magnetic intensity inside solenoid ,
   ampere-turn per metre
8 // As  $B = \mu_0 * (H + I) \Rightarrow I = B/\mu_0 - H$ 
9 // But  $B = \mu_0 * \mu_r * H$  and solving for I
10 I = (mu_r - 1) * H;
11 printf("\nThe Intensity of magnetisation inside the
   solenoid , I = %5.3e A/m", I);
12 M = I * V;    // Magnetic moment of the rod , ampere
   metre square
13 printf("\nThe magnetic moment of the rod , M = %3d
   ampere metre square", M)
14
15 //Result
16 // The Intensity of magnetisation inside the
   solenoid , I = 2.998e+005 A/m
17 // The magnetic moment of the rod , M = 299 ampere
   metre square

```

---

**Scilab code Exa 12.3** Magnetic moment of the rod placed inside the solenoid

```

1 // Scilab Code Ex12.3 : Page-604 (2011)
2 clc; clear;
3 n = 300;....// Number of turns wound per metre on
   the solenoid
4 i = 0.5;....// Current through the solenoid , A
5 V = 1e-03;....// Volume of iron rod , per metre cube
6 mu_r = 100;    // Relative permeability of the iron
7 H = n*i;    // Magnetic intensity inside solenoid ,

```

```

    ampere-turn per metre
8 // As,  $I = (B - \mu_o * H) / \mu_o$ 
9 //But,  $B = \mu * H = \mu_r * \mu_o * H$  and  $I = (\mu_r - 1) * H$ 
10  $I = (\mu_r - 1) * n * i$ ;
11 printf("\\nThe Intensity of magnetisation inside the
    solenoid ,  $I = 5.3e A/m$ ", I);
12  $l = 0.2$ ;....//length of the rod,m
13  $r = 5e-3$ ;....//radius of the rod,m
14  $V = 1.57e-5$ ;....// $V = \pi * r^2 * l$  where the volume of
    the rod having radius r and length,m
15  $M = I * V$  ; // Magnetic moment of the rod ,
    ampere metre square
16 printf("\\nThe magnetic moment of the rod ,  $M = 5.3e$ 
    ampere metre square",M)
17
18 //Result
19 // The Intensity of magnetisation inside the
    solenoid ,  $I = 1.485e+004 A/m$ 
20 // The magnetic moment of the rod ,  $M = 0.233$  ampere
    metre square

```

---

**Scilab code Exa 12.4** Magnetizing force and relative permeability of the material

```

1 // Scilab Code Ex12.4 : Page-605 (2011)
2 clc; clear;
3  $B = 0.0044$ ;....// Magnetic flux density , weber/meter
    square
4  $\mu_o = 4 * \pi * 1e-07$ ;....// Relative permeability of
    the material , henry/m
5  $I = 3300$ ;....// Magnetization of a magnetic material
    , A/m
6 // $B = \mu_o * (I + H)$  , solving for H
7  $H = (B / \mu_o) - I$ ;....// Magnetizing force ,A/m

```

```

8 printf("\nThe magnetic intensity ,H = %3d A/m",H);
9 // Relation between intensity of magnetization and
  relative permeability
10 mu_r = (I/H)+1;....//substitute the value of I and H
11 printf("\nThe relative permeability , mu_r = %5.2f",
  mu_r);
12
13 //Result
14 // The magnetic intensity ,H = 201 A/m
15 // The relative permeability , mu_r = 17.38

```

---

**Scilab code Exa 12.5** Magnetic flux density and magnetic intensity

```

1 // Scilab Code Ex12.5 : Page-605 (2011)
2 clc; clear;
3 mu_o = 4*pi*1e-07;....// Magnetic permeability of
  the free space , henery/m
4 mu_r = 600;
5 mu = mu_o*mu_r; // Magnetic permeability of the
  medium, henery/m
6 n = 500;...// Turns in a wire
7 i = 0.3;....// Current flows through a ring ,amp
8 r = 12e-02/2;....// Mean radius of a ring , m
9 B = mu_o*mu_r*n*i/(2*pi*r);
10 printf("\nThe magnetic flux density = %2.1f weber/
  meter-square", B);
11 H = B/mu; // Magnetic intensity , ampere-turns/m
12 printf("\nThe magnetic intensity = %5.1f ampere-
  turns/m", H);
13 // As  $B = \mu_o \cdot (I + H) \Rightarrow \mu_o \cdot I = B - \mu_o \cdot H$ 
14 printf("\nThe percentage magnetic flux density due
  to electronic loop currents = %5.2f percent", (B
  - mu_o*H)/B*100);
15
16 // Result

```

```

17 // The magnetic flux density = 0.3 weber/meter-
    square
18 // The magnetic intensity = 397.9 ampere-turns/m
19 // The percentage magnetic flux density due to
    electronic loop currents = 99.83 percent

```

---

**Scilab code Exa 12.6** Total dipole moment of the sample

```

1 // Scilab Code Ex12.6 : Page-606 (2011)
2 clc; clear;
3 M_i = 4.5;....// Intial value of total dipole moment
    of the sample
4 H_i = 0.84;....// External magnetic field , tesla
5 T_i = 4.2;....// Cooling temerature of the sample , K
6 H_f = 0.98;....// External magnetic field , tesla
7 T_f = 2.8;....// Cooling temerature of the sample , K
8 // According to the curie 's law ,  $M_f/M_i = (H_f/H_i)*(T_i/T_f)$ 
9 M_f = M_i*H_f/H_i*T_i/T_f;
10 printf("\\nThe total dipole moment of the sample = %5
    .3f joule/tesla",M_f);
11
12 // Result
13 // The total dipole moment of the sample = 7.875
    joule/tesla

```

---

**Scilab code Exa 12.7** Magnetization and magnetic moment in the bar

```

1 // Scilab Code Ex12.7 : Page-606 (2011)
2 clc; clear;
3 mu_o = 4*pi*1e-07;....// Magnetic permeability of
    free space , henry/m

```

```

4 n = 1e+29;....// Number density of atoms of iron ,
   per metre cube
5 p_m = 1.8e-23;....// Magnetic moment of each atom,
   ampere-metre square
6 k_B = 1.38e-23;....// Boltzmann constant , J/K
7 B = 0.1;    // Magnetic flux density , weber per
   metre square
8 T = 300;....// Absolute room temperature , K
9 l = 10e-02; // Length of the iron bar , m
10 a = 1e-04; // Area of cross-section of the iron bar
   , metre square
11 V = l*a;    // Voluem of the iron bar , metre cube
12 chi = n*p_m^2*mu_o/(3*k_B*T);
13 printf("\nThe paramagnetic susceptibility of a
   material = %5.3e", chi);
14 pm_mean = p_m^2*B/(3*k_B*T); // Mean dipole moment
   of an iron atom , ampere metre-square
15 P_m = n*pm_mean; // Dipole moment of the bar ,
   ampere metre-square
16 I = n*p_m; // Magnetization of the bar in one
   domain , ampere/metre
17 M = I*V;    // Magnetic moment of the bar , ampere
   metre-square
18 printf("\nThe dipole moment of the bar = %5.3e
   ampere metre-square", P_m);
19 printf("\nThe magnetization of the bar in one domain
   = %3.1e ampere/metre", I);
20 printf("\nThe magnetic moment of the bar = %2d
   ampere metre-square", M);
21
22 // Result
23 // The paramagnetic susceptibility of a material =
   3.278e-03
24 // The dipole moment of the bar = 2.609e+02 ampere
   metre-square
25 // The magnetization of the bar in one domain = 1.8e
   +06 ampere/metre
26 // The magnetic moment of the bar = 18 ampere metre-

```

square

---

**Scilab code Exa 12.8** Hysteresis loss of energy per hour in the iron core of the transformer

```
1 // Scilab Code Ex12.8 :Page-607 (2011)
2 clc; clear;
3 A = 500;....// Area of the B-H loop, joule per metre
   cube
4 n = 50;....// Total number of cycles, Hz
5 m = 9;....// Mass of the core, kg
6 d = 7.5e+3;....// Density of the core, kg/metre cube
7 t = 3600;....// Time during which the energy loss
   takes place, s
8 V = m/d;....// Volume of the core, metre cube
9 E = n*V*A*t;....// Hystersis loss of energy per hour
   , joule
10 printf("\nThe hystersis loss per hour = %5.2eJ", E);
11
12 // Result
13 // The hystersis loss per hour = 1.08e+005J
```

---

**Scilab code Exa 12.9** Hystersis loss from BH loop

```
1 // Scilab Code Ex12.9 : Page-607 (2011)
2 clc; clear;
3 n = 50;....// Total number of cycles per sec, Hz
4 V = 1e-03;....// Volume of the specimen, metre cube
5 t = 1;....// Time during which the loss occurs, s
6 A = 0.25e+03;....// Area of B-H loop, joule per
   metre cube
7 E = n*V*A*t; // Energy loss due to hysteresis, J/
   s
```

```

8 printf("\nThe hystersis loss = %4.1f J/s", E);
9
10 // Result
11 // The hystersis loss = 12.5 J/s

```

---

**Scilab code Exa 12.10** Change in the magnetic dipole moment of the electron

```

1 // Scilab Code Ex12.10 : Page-608 (2011)
2 clc; clear;
3 e = 1.6e-19;....// Charge on anelectron, C
4 m = 9.1e-31;....// Mass of the electron, kg
5 r = 5.1e-11;....// Radius of the electronic orbit, m
6 B = 2.0;....// Applied magnetic field, weber per
   metre-square
7 delta_pm = e^2*r^2*B/(4*m);
8 printf("\nThe change in the magnetic dipole moment
   of the electron = %3.1e A-metre square", delta_pm
   );
9
10 // Result
11 // The change in the magnetic dipole moment of the
   electron = 3.7e-29 A-metre square

```

---



## Chapter 13

# Dielectric Properties of Materials

Scilab code Exa 13.1 Electric dipole placed in a uniform electric field

```
1 // Scilab Code Ex13.1: Page-648 (2011)
2 clc;clear;
3 q = 1e-006; // Electric charge on either side of
   the dipole , C
4 l = 2e-02; // Dipole length , m
5 p = q*l;....// Dipole moment for the pair of
   opposite charges , C-m
6 E = 1e+005;....// External electric field , N/C
7 theta = 90;....// Angle which the dipole makes with
   the external field , degrees
8 tau = p*E*sind(theta);....// The maximum torque on
   dipole placed in external electric field , Nm
9 printf("\nThe maximum torque = %1.0e N-m", tau);
10 W = integrate('p*E*sin(thet)', 'thet', 0, %pi);
   // The work done in rotating the dipole direction
   = %1.0e J", W
11 printf("\nThe work done in rotating the dipole
   direction = %1.0e J", W);
12
```

```

13 // Result
14 // The maximum torque = 2e-003 N-m
15 // The work done in rotating the dipole direction =
    4e-003 J

```

---

**Scilab code Exa 13.2** Force acting on an electric dipole in different orientations relative to the electric field

```

1 // Scilab Code Ex13.2:Page-648 (2011)
2 clc;clear;
3 Q = 8e-019;....// Charge of the nucleus, C
4 p = 3.2e-029;....// Electric dipole moment, C-m
5 r = 1e-10;      // Distance of dipole relative to the
    nucleus, m
6 k = 9e+9;....// Coulomb constant, N-meter-square/C-
    square
7 theta = 0;....// Angle for radial direction, radian
8 F = k*p*Q*sqrt(3*cos(theta^2)+1)/r^3;    // The
    force acting on the dipole in the radial
    direction, N
9 printf("\\nThe force acting on the dipole in the
    radial direction = %3.1e N", F);
10 theta = %pi/2;....// Angle for perpendicular
    direction, radian
11 F = k*p*Q*sqrt(3*cos(theta)^2+1)/r^3;
12 printf("\\nThe force acting on the dipole in the
    direction perpendicular to radial direction = %3
    .1e N", F);
13
14 // Result
15 // The force acting on the dipole in the radial
    direction = 4.6e-007 N
16 // The force acting on the dipole in the direction
    perpendicular to radial direction = 2.3e-007 N

```

---

**Scilab code Exa 13.3** Dielectric constant and the electric permittivity of the material

```
1 // Scilab Code Ex13.3: Page-649 (2011)
2 clc;clear;
3 chi_e = 35.4e-12;....// Susceptability of the
   material , C-square/N-meter-square
4 eps_0 = 8.85e-12;....// Electric permittivity in
   free space , C-squre/N-meter-square
5 K = 1 + (chi_e/eps_0);
6 printf("\nThe dielectric constant = %d ",K);
7 eps = (eps_0*K);
8 printf("\nThe electric permittivity = %5.3e C-
   square/N-meter square ",eps);
9
10 // Result
11 // The dielectric constant = 5
12 // The electric permittivity = 4.425e-011 C-square/
   N-meter square
```

---

**Scilab code Exa 13.4** Dielectric constant and the electric susceptability of diamond

```
1 // Scilab Code Ex13.4:Page-649 (2011)
2 clc;clear;
3 eps = 1.46e-10;....// Electric permittivity , C-
   square/n-meter-square
4 eps_0 = 8.85e-12;....// Permittivity in free space ,
   C-squre/N-meter-square
5 K = (eps/eps_0);
6 printf("\nThe dielectric constant = %4.1f ", K);
```

```

7 chi_e = eps_0*(K-1);....// Susceptability ,in C-
  square/N-meter-square
8 printf("\nThe electric susceptability = %4.2e C-
  square/N-meter square ", chi_e);
9
10 // Result
11 // The dielectric constant = 16.5
12 // The electric susceptability = 1.37e-010 C-square/
  N-meter square

```

---

**Scilab code Exa 13.5** Calculate the values of E D and P

```

1 // Scilab Code Ex13.5 Page-650 (2011)
2 clc;clear;
3 K = 7.0;....// Dielectric constant of the slab
4 d = 0.01;....// Distance between the two parallel
  plates , m
5 V_0 = 100;....// Potential difference across the
  plates , V
6 eps_0 = 8.85e-12;....// Electric permability of the
  free space , C-square/N-meter-square
7 E_0 = V_0/d;....// Electric intensity in the absence
  of dielectric slab , V/m
8 E = E_0/K; // Electric intensity with dielectric
  slab introduced between the plates , V/m
9 printf("\nThe electric field intensity in the
  presence of the dielectric slab = %4.2e V/m ", E)
  ;
10 D = (eps_0*K*E); // Electric displacement , C-
  square/m-square
11 printf("\nThe electric displacement in the
  dielectric slab = %4.2e C-square/meter-square ",D
  );
12 P = eps_0*(K-1)*E; // Electric polarization in
  the dielectric slab , C-square/m-square

```

```

13 printf("\nThe electric polarization in the
    dielectric slab = %3.1e C-square/meter-square ", P
    );
14
15 // Result
16 // The electric field intensity in the presence of
    the dielectric slab = 1.43e+003 V/m
17 // The electric displacement in the dielectric slab
    = 8.85e-008 C-square/meter-square
18 // The electric polarization in the dielectric slab
    = 7.6e-008 C-square/meter-square

```

---

**Scilab code Exa 13.6** Dipole moment induced in He atom

```

1 // Scilab Code Ex13.6: Page-650 (2011)
2 clc;clear;
3 K = 1.000074;....// Dielectric constant of the He
4 n = 2.69e+025;....// Atomic density of He, atoms/
    meter-cube
5 eps_0 = 8.85e-012;....// Electric permability of the
    free space, C-square/N-meter-square
6 E = 1;....// Electric field strength, V/m
7 p = (eps_0*(K-1)*E)/n;    // Dipole moment induced
    in He, C-m
8 printf("\nThe dipole moment induced in each He atom
    = %4.2e C-m ", p);
9
10 // Result
11 // The dipole moment induced in each He atom = 2.43e
    -041 C-m

```

---

**Scilab code Exa 13.7** Induced dipole moment and atomic polarizability of neon gas

```

1 // Scilab Code Ex13.7: Page-650 (2011)
2 clc;clear;
3 K = 1.000134;....// Dielectric constant of the neon
4 n = 2.69e+25;....// Atomic density of argon,atoms/
   meter-cube
5 eps_0 = 8.85e-12;....// Electric Permability in the
   free space, C-square/N-meter-square
6 E = 90e+03; // External electric field, V/m
7 p = eps_0*(K-1)*E/n; // Dipole moment induced in
   each neon atom, C-m
8 alpha = p/E; // Atomic polarizability of neon gas
   , C-metre-square/V
9 printf("\\nThe induced dipole moment of noen atom =
   %4.2e C-m", p) ;
10 printf("\\nThe electronic polarizability of neon gas
   = %3.1e C-m-square/V ", alpha);
11
12 // Result
13 // The induced dipole moment of noen atom = 3.97e
   -036 C-m
14 // The electronic polarizability of neon gas = 4.4e
   -041 C-m-square/V

```

---

**Scilab code Exa 13.8** Electronic polarizability of argon atom

```

1 // Scilab Code Ex13.8: Page-651 (2011)
2 clc;clear;
3 K = 1.0024;....// Dielectric constant of the argon
4 n = 2.7e+25;....// Atomic density of argon,atoms/
   meter-cube
5 eps_0 = 8.85e-12;....// Electric Permability in the
   free space, C-square/N-meter-square
6 alpha = eps_0*(K-1)/n;
7 printf("\\nThe electronic polarizability of argon
   atom = %4.1e C-m-square/V ", alpha);

```

```

8
9 // Result
10 // The electronic polarizability of argon atom = 7.9
    e-040 C-m-square/V

```

---

**Scilab code Exa 13.9** Individual dipole moment of carbon tetrachloride

```

1 // Scilab Code Ex13.9: Page-651 (2011)
2 clc;clear;
3 K = 2.24;....// Dielectric constant
4 eps_0 = 8.85e-12;....// Electric permability in the
    free space, C-square/N-meter-square
5 rho = 1.6e+003;....// Density of CCl4, kg/meter-cube
6 M = 156;....// Molecular weight of CCl4
7 E = 1e+007;....// External electric field strength,
    V/m
8 N_A = 6.02e+26; // Avogadro's number, per kmol
9 rho_M = rho*N_A/M; // Molecular density of CCl4
10 p = eps_0*(K-1)*E/rho_M; // Individual dipole
    moment of CCL4 molecule, C-m
11 printf("\nIndividual dipole moment of CCL4 molecule
    = %4.2e C-m ", p);
12
13 // Result
14 // Individual dipole moment of CCL4 molecule = 1.78e
    -032 C-m

```

---

**Scilab code Exa 13.10** Atomic radius of He

```

1 // Scilab Code Ex13.10: Page-652 (2011)
2 clc;clear;
3 K = 1.0000684;....// Dielectric constant of He at 1
    atm

```

```

4 n = 2.7e+25;....// Density of He at 1 atm and 273 K,
   atoms/meter-cube
5 // The atomic polarizability , alpha = eps_0*(K-1)/n
6 // In terms of atomic radius , alpha = 4*pi*eps_0*R
   ^3 so , we have
7 R = ((K-1)/(4*pi*n))^(1/3);    // Radius of He atom
   , m
8 printf("\nThe atomic radius of He = %4.2e m ", R);
9
10 // Result
11 // The atomic radius of He = 5.86e-011 m

```

---

**Scilab code Exa 13.11** Percentage of ionic polarizability in NaCl crystal

```

1 // Scilab Code Ex13.11: Page-652 (2011)
2 clc;clear;
3 mu = 1.5;....// Optical index of refraction of NaCl
   crystal
4 K = 5.6;....// Static dielectric constant of NaCl
   crystal
5 P_IP = (1-((mu^2-1)*(K+2))/((mu^2+2)*(K-1)))*100;
6 printf("\nThe percentage of ionic polarizability in
   NaCl crystal = %4.1f percent ", P_IP);
7
8 // Result
9 // The percentage of ionic polarizability in NaCl
   crystal = 51.4 percent

```

---

**Scilab code Exa 13.12** Determine the dipole moment

```

1 // Scilab Code Ex13.12: Page-653 (2011)
2 clc;clear;
3 K_B = 1.38e-23;....// Boltzmann constant , J/mol/K

```



```

4 T = 300;....// Room temperature , K
5 eps_0 = 8.85e-12;....// Electric permittivity of
  free space , F/m
6 N_A = 6.0e+23;      // Avogadro 's number
7 n2 = N_A*1000;      // Number of molecules of non-
  polar substance in 1000 cc volume
8 p_0 = sqrt((9*K_B*T*eps_0*0.023)/n2);    // Dipole
  moment of polar molecules , C-m
9 printf("\nThe dipole moment of polar molecules = %5
  .3e C-m", p_0);
10
11 // Result
12 // The dipole moment of polar molecules = 3.555e-030
  C-m

```

---

# Chapter 14

## Solid State Electronics

Scilab code Exa 14.1 Density of impurity atoms to N type and P type silicon

```
1 // Scilab code Ex14.1 : Pg:718(2011)
2 clc;clear;
3 e = 1.6e-019; // Charge on an electron , C
4 mu_h = 0.048; // Mobility of holes , metre square/
    volt-s
5 mu_e = 0.135; // Mobility of electrons , metre
    square/volt-s
6 // For P-type semiconductor
7 rho_p = 1e-01; // Resistivity of P type silicon ,
    omh-m
8 // As rho_p = 1/(e*N_a*mu_h), solving for N_a
9 N_a = 1/(e*rho_p*mu_h); // Density of acceptor atoms
    , per metre cube
10 // For N-type semiconductor
11 rho_n = 1e-01; // Resistivity of N type silicon ,
    omh-m
12 // As rho_n = 1/(e*N_d*mu_h), solving for N_d
13 N_d = 1/(e*rho_n*mu_e); // Density of donor atoms ,
    per metre cube
14 printf("\nDensity of acceptor atoms = %4.2e per
```

```

    metre cube", N_a);
15 printf("\nDensity of donor atoms = %4.2e per metre
    cube", N_d);
16
17 // Result
18 // Density of acceptor atoms = 1.30e+21 per metre
    cube
19 // Density of donor atoms = 4.63e+20 per metre cube

```

---

**Scilab code Exa 14.2** Electrical conductivity and resistivity of intrinsic germanium sample

```

1 // Scilab code Ex14.2 : Pg:718(2011)
2 clc;clear;
3 e = 1.6e-019; // Charge on an electron , C
4 mu_e = 0.36; // Mobility of an electron , metre
    square/V-s
5 mu_h = 0.17; // Mobility of a hole , metre square/
    V-s
6 n_i = 2.5e+018; // Intrinsic concentration of Ge
    sample, per metre cube
7 sigma = e*n_i*(mu_h+mu_e); // Electrical
    conductivity of Ge sample, mho per metre
8 rho = 1/sigma; // Electrical resistivity of Ge, ohm
    -m
9 printf("\nThe electrical conductivity of intrinsic
    germanium sample = %5.3f mho/m", sigma);
10 printf("\nThe electrical resistivity of intrinsic
    germanium sample = %3.1f ohm-m", rho);
11
12 // Result
13 // The electrical conductivity of intrinsic
    germanium sample = 0.212 mho/m
14 // The electrical resistivity of intrinsic germanium
    sample = 4.7 ohm-m

```

---

**Scilab code Exa 14.3** Electrical conductivity of undoped and doped silicon

```
1 // Scilab code Ex14.3 : Pg:719(2011)
2 clc;clear;
3 e = 1.6e-019; // Charge on an electron , C
4 mu_e = 0.13; // Mobility of an electron , metre
    square/V-s
5 mu_h = 0.05; // Mobility of a hole , metre square/
    V-s
6 n_i = 1.5e+016; // Intrinsic concentration of Si ,
    per metre cube
7 // Pure Si
8 sigma = e*n_i*(mu_h+mu_e); // Electrical
    conductivity of Si , mho per metre
9 // Pure Si doped with donor impurity
10 n_e = 5e+028/1e+09; // Concentration of
    electrons , per metre cube
11 sigma_n = e*n_e*mu_e; // Electrical conductivity
    of Si doped with donor impurity , mho per metre
12 // Pure Si doped with acceptor impurity
13 n_h = 5e+028/1e+09; // Concentration of holes ,
    per metre cube
14 sigma_p = e*n_h*mu_h; // Electrical conductivity
    of Si doped with acceptor impurity , mho per metre
15 printf("\nThe electrical conductivity of pure Si =
    %4.2e mho/m", sigma);
16 printf("\nThe electrical conductivity of Si doped
    with donor impurity = %4.2f mho/m", sigma_n);
17 printf("\nThe electrical conductivity of Si doped
    with acceptor impurity= %4.2f mho/m", sigma_p);
18
19 // Result
20 // The electrical conductivity of pure Si = 4.32e-04
```

```

    mho/m
21 // The electrical conductivity of Si doped with
    donor impurity = 1.04 mho/m
22 // The electrical conductivity of Si doped with
    acceptor impurity= 0.40 mho/m

```

---

**Scilab code Exa 14.4** Shift in Fermi level due to change in density of donor atoms

```

1 // Scilab code Ex14.4 : Pg:720(2011)
2 clc;clear;
3 Nd = 1; // For simplicity assume donor
    concentration to be unity, per metre cube
4 Nd_prime = 3*Nd; // Thrice the donor concentration
    , per metre cube
5 dE_CF1 = 0.5; // Energy difference between normal
    Fermi level and conduction level, eV
6 k_BT = 0.03; // Thermal energy at room
    temperature, eV
7 // As Nd_prime/Nd = exp((dE_CF1 - dE_CF2))/k_BT,
    solving for dE_CF2
8 dE_CF2 = dE_CF1 - k_BT*log(Nd_prime/Nd); // Energy
    difference between new position of Fermi level and
    conduction level, eV
9 printf("\nThe new position of Fermi level when donor
    concentration is trebled = %5.3f eV", dE_CF2);
10
11 // Result
12 // The new position of Fermi level when donor
    concentration is trebled = 0.467 eV

```

---

**Scilab code Exa 14.5** Voltage required to cause a forward current density in pn junction diode

```

1 // Scilab code Ex14.5 :Pg:721(2011)
2 clc;clear;
3 e = 1.6e-019; // Charge on an electron , C
4 T = 300; // Room temperature , K
5 J0 = 300e-03; // Saturation current density of the
   pn junction diode , A/metre square
6 J = 1e+05; // Forward current density of pn
   junction diode , A/metre square
7 k_B = 1.38e-023; // Boltzmann constant , J/K
8 eta = 1; // Ideality factor for Ge diode
9 // As  $J = J_0 \exp(eV / (\eta k_B T))$ , solving for V
10 V = eta*k_B*T/e*log(J/J0); // Voltage required to
   cause a forward current density in pn junction
   diode , volt
11 printf("\\nThe voltage required to cause a forward
   current density in pn junction diode = %5.3f V",
   V);
12
13 // Result
14 // The voltage required to cause a forward current
   density in pn junction diode = 0.329 V

```

---

**Scilab code Exa 14.6** Applied voltage for forward current density

```

1 // Scilab code Ex14.6 : Pg:721(2011)
2 clc;clear;
3 e = 1.6e-019; // Charge on an electron , C
4 T = 300; // Room temperature , K
5 J0 = 200e-03; // Saturation current density of the
   pn junction diode , A/metre square
6 J = 5e+04; // Forward current density of pn
   junction diode , A/metre square
7 k_B = 1.38e-023; // Boltzmann constant , J/K
8 eta = 1; // Ideality factor for Ge diode
9 // As  $J = J_0 \exp(eV / (\eta k_B T))$ , solving for V

```

```

10 V = eta*k_B*T/e*log(J/J0);    // Voltage required to
    cause a forward current density in pn junction
    diode , volt
11 printf("\nThe voltage required to cause a forward
    current density in pn junction diode = %5.3f V",
    V);
12
13 // Result
14 // The voltage required to cause a forward current
    density in pn junction diode = 0.322 V

```

---

**Scilab code Exa 14.7** Forward voltage to increase the current density of Si diode

```

1 // Scilab code Ex14.7 : Pg:722(2011)
2 clc;clear;
3 e = 1.6e-019;    // Charge on an electron , C
4 T = 300;        // Room temperature , K
5 J0 = 300e-03;   // Saturation current density of the
    pn junction diode , A/metre square
6 J = 1e+05;     // Forward current density of pn
    junction diode , A/metre square
7 k_B = 1.38e-023; // Boltzmann constant , J/K
8 eta = 2;       // Ideality factor for Ge diode
9 // As  $J = J0 \cdot \exp(e \cdot V / (\eta \cdot k_B \cdot T))$ , solving for V
10 V = eta*k_B*T/e*log(J/J0);    // Voltage required to
    cause a forward current density in pn junction
    diode , volt
11 printf("\nThe voltage required to cause a forward
    current density in Si iode = %5.3f V", V);
12
13 // Result
14 // The voltage required to cause a forward current
    density in Si diode = 0.658 V

```

---

### Scilab code Exa 14.8 Static and dynamic values of diode resistance

```
1 // Scilab code Ex14.8 : Pg:723(2011)
2 clc;clear;
3 I = 55e-03; // Forward current through Si diode,
  A
4 V = 3; // Forward bias across Si diode, V
5 eta = 2; // Ideality factor for Si diode
6 R_dc = V/I; // Static diode resistance, ohm
7 R_ac = 0.026*eta/I; // Dynamic diode resistance,
  ohm
8 printf("\nThe static diode resistance = %4.1f ohm",
  R_dc);
9 printf("\nThe dynamic diode resistance = %5.3f ohm",
  R_ac);
10
11 // Result
12 // The static diode resistance = 54.5 ohm
13 // The dynamic diode resistance = 0.945 ohm
```

---

### Scilab code Exa 14.9 Half wave rectifier parameters

```
1 // Scilab code Ex14.9 : Pg:723(2011)
2 clc;clear;
3 R_L = 1000; // Load resistance across HWR, ohm
4 V_rms = 200; // Rms value of voltage supply, V
5 V0 = sqrt(2)*V_rms; // Peak value of voltage, V
6 I0 = V0/(R_L*1e-03); // Peak value of current, mA
7 I_dc = I0/%pi; // Average value of current, mA
8 I_rms = I0/2; // Rms value of current, mA
9 V_dc = I_dc*R_L/1e+03; // Dc output voltage, V
10 PIV = V0; // Peak inverse voltage, V
```



```

11 printf("\nThe average value of current = %2d mA",
    I_dc);
12 printf("\nThe rms value of current = %5.1f mA",
    I_rms);
13 printf("\nThe dc output voltage = %2d V", V_dc/1);
14 printf("\nPIV = %5.1f V", PIV);
15
16
17 // Result
18 // The average value of current = 90 mA
19 // The rms value of current = 141.4 mA
20 // The dc output voltage = 90 V
21 // PIV = 282.8 V

```

---

#### Scilab code Exa 14.10 Full wave rectifier parameters

```

1 // Scilab code Ex14.10 : Pg:724(2011)
2 clc;clear;
3 R_L = 980; // Load resistance across FWR, ohm
4 R_F = 20; // Internal resistance of two crystal
    diodes in FWR, ohm
5 V_rms = 50; // Rms value of voltage supply, V
6 V0 = sqrt(2)*V_rms; // Peak value of voltage, V
7 I0 = V0/((R_L+R_F)*1e-03); // Peak value of
    current, mA
8 I_dc = 2*I0/%pi; // Average value of current, mA
9 I_rms = I0/sqrt(2); // Rms value of current, mA
10 V_dc = I_dc*R_L/1e+03; // Dc output voltage, V
11 eta = 81.2/(1+R_F/R_L); // Rectification
    efficiency
12 PIV = 2*V0; // Peak inverse voltage, V
13 printf("\nThe average value of current = %2d mA",
    I_dc);
14 printf("\nThe rms value of current = %2d mA", I_rms)
    ;

```

```

15 printf("\nThe dc output voltage = %4.1f V", V_dc/1);
16 printf("\nThe rectification efficiency = %4.1f
    percent", eta);
17 printf("\nPIV = %5.1f V", PIV);
18
19
20 // Result
21 // The average value of current = 45 mA
22 // The rms value of current = 50 mA
23 // The dc output voltage = 44.1 V
24 // The rectification efficiency = 79.6 percent
25 // PIV = 141.4 V

```

---

#### Scilab code Exa 14.11 Current gains in BJT

```

1 // Scilab code Ex14.11 : Pg:725(2011)
2 clc;clear;
3 delta_IC = 1e-03; // Change in collector current,
    A
4 delta_IB = 50e-06; // Change in base current, A
5 bta = delta_IC/delta_IB; // Base current
    amplification factor
6 alpha = bta/(1+bta); // Emitter current
    amplification factor
7 printf("\nAlpha of BJT = %4.2f", alpha);
8 printf("\nBeta of BJT = %2d", bta);
9
10
11 // Result
12 // Alpha of BJT = 0.95
13 // Beta of BJT = 20

```

---

#### Scilab code Exa 14.12 Base current of BJT in CB mode

```

1 // Scilab code Ex14.12 : Pg:725(2011)
2 clc;clear;
3 I_E = 2; // Emitter current , mA
4 alpha = 0.88; // Emitter current amplification
   factor
5 I_C = alpha*I_E; // Collector current , mA
6 I_B = I_E - I_C; // Base current of BJT in CB
   mode, mA
7 printf("\nThe base current of BJT in CB mode = %4.2 f
   mA", I_B);
8
9
10 // Result
11 // The base current of BJT in CB mode = 0.24 mA

```

---

**Scilab code Exa 14.13** Current gain and base current of BJT in CB mode

```

1 // Scilab code Ex14.13 : Pg:725(2011)
2 clc;clear;
3 I_CB0 = 12.5e-03; // Reverse saturation current ,
   mA
4 I_E = 2; // Emitter current , mA
5 I_C = 1.97; // Collector current , mA
6 // As I_C = alpha*I_E+I_CB0, solving for alpha
7 alpha = (I_C - I_CB0)/I_E; // Emitter current gain
8 I_B = I_E - I_C; // Base current , mA
9 printf("\nThe emitter current gain = %5.3 f", alpha);
10 printf("\nThe base current = %4.2 f mA", I_B);
11
12
13 // Result
14 // The emitter current gain = 0.979
15 // The base current = 0.03 mA

```

---

**Scilab code Exa 14.14** Current gain and leakage current of BJT in CE mode

```
1 // Scilab code Ex14.14 : Pg:726(2011)
2 clc;clear;
3 alpha = 0.98; // Emitter current amplification
  factor
4 bta = alpha/(1-alpha); // Emitter current
  amplification factor
5 I_CBO = 5e-06; // Reverse saturation current, A
6 I_CEO = 1/(1-alpha)*I_CBO; // Leakage current of
  BJT in CE mode, mA
7 printf("\nThe base current gain = %2g", bta);
8 printf("\nThe leakage current of BJT in CE mode = %4
  .2 f mA", I_CEO/1e-03);
9
10
11 // Result
12 // The base current gain = 49
13 // The leakage current of BJT in CE mode = 0.25 mA
```

---

**Scilab code Exa 14.15** Voltage and power gain of PNP transistor in CB mode

```
1 // Scilab code Ex14.15 : Pg:726(2011)
2 clc;clear;
3 R_i = 50; // Dynamic input resistance of PNP
  transistor, ohm
4 R_L = 5e+03; // Load resistance in collector
  circuit, ohm
5 alpha = 0.96; // Emitter current amplification
  factor
```

```
6 A_v = alpha*R_L/R_i;    // Voltage gain
7 A_p = alpha*A_v;       // Power gain
8 printf("\nThe voltage gain = %2g", A_v);
9 printf("\nThe power gain = %2d", A_p);
10
11
12 // Result
13 // The voltage gain = 96
14 // The power gain = 92
```

---

# Chapter 15

## Digital Electronics

Scilab code Exa 15.1 Binary equivalent of decimal number

```
1 // Scilab code Ex15.1 : Pg:771(2011)
2 clc;clear;
3 function [bin]= decimal_binary(n) // Function to
   convert decimal to binary
4     bin = 0;
5     i = 1;
6     while (n <> 0)
7         rem = n-fix(n./2).*2;
8         n = int(n/2);
9         bin = bin + rem*i;
10        i = i * 10;
11    end
12 endfunction
13
14 n = 25; // Initialize the decimal number
15 printf("Binary equivalent of %d = %d", n,
   decimal_binary(n));
16
17 // Result
18 // Binary equivalent of 25 = 11001
```

---

### Scilab code Exa 15.2 Decimal equivalent of 6 bit binary number

```
1 // Scilab code Ex15.2 : Pg:771(2011)
2 clc;clear;
3 function [deci]= binary_decimal(ni) // Function to
   convert binary to decimal
4     deci = 0;
5     i = 0;
6     while (ni <> 0)
7         rem = ni-fix(ni./10).*10;
8         ni = int(ni/10);
9         deci = deci + rem*2.^i;
10        i = i + 1;
11    end
12 endfunction
13
14 function [decf]= binfrac_decifrac(nf) // Function to
   convert binary fraction to decimal fraction
15     decf = 0;
16     i = -1;
17     while (i >= -3)
18         nf = nf*10;
19         rem = round(nf);
20         nf = nf-rem;
21         decf = decf + rem*2.^i;
22         i = i - 1;
23     end
24 endfunction
25
26 n = 101.101; // Initialize the binary number
27 n_int = int(n); // Extract the integral part
28 n_frac = n-n_int; // Extract the fractional part
29 printf("Decimal equivalent of %7.3f = %5.3f", n,
   binary_decimal(n_int)+binfrac_decifrac(n_frac));
```

```
30
31 // Result
32 // Decimal equivalent of 101.101 = 5.625
```

---

**Scilab code Exa 15.3** Decimal equivalent of octal number

```
1 // Scilab code Ex15.3 : Pg:772(2011)
2 clc;clear;
3 function [dec]= octal_decimal(n) // Function to
   convert binary to decimal
4     dec = 0;
5     i = 0;
6     while (n <> 0)
7         rem = n-fix(n./10).*10;
8         n = int(n/10);
9         dec = dec + rem*8.^i;
10        i = i + 1;
11    end
12 endfunction
13
14 n = 173; // Initialize the octal number
15 printf("Decimal equivalent of %d = %d", n,
   octal_decimal(n));
16
17 // Result
18 // Decimal equivalent of 173 = 123
```

---

**Scilab code Exa 15.4** Octal equivalent of decimal number

```
1 // Scilab code Ex15.4 : Pg:772(2011)
2 clc;clear;
3 function octal = decimal_octal(n) // Function to
   convert decimal to octal
```



```

4     i=1; octal = 0;
5     while (n<>0)
6         rem = n-fix(n./8).*8;
7         octal = octal + rem*i;
8         n = int(n/8);
9         i = i*10;
10    end
11 endfunction
12
13 n = 278;    // Initialize the octal number
14 printf("The octal equivalent of %d = %d", n,
        decimal_octal(n));
15
16 // Result
17 // The octal equivalent of 278 = 426

```

---

#### Scilab code Exa 15.5 Hexadecimal equivalent of binary numbers

```

1 // Scilab code Ex15.5 :Pg:772(2011)
2 clc;clear;
3 function hex = binary_hex(n) // Function to convert
    decimal to hexadecimal
4     hex = emptystr();
5     while (n <>0)
6         rem = n-fix(n./10000).*10000;    // Division
            Algorithm
7         if rem == 0 then
8             hex = hex+'0';
9         elseif rem == 1 then
10            hex = hex+'1';
11        elseif rem == 10 then
12            hex = hex+'2';
13        elseif rem == 11 then
14            hex = hex+'3';
15        elseif rem == 100 then

```

```

16         hex = hex+'4';
17     elseif rem == 101 then
18         hex = hex+'5';
19     elseif rem == 110 then
20         hex = hex+'6';
21     elseif rem == 111 then
22         hex = hex+'7';
23     elseif rem == 1000 then
24         hex = hex+'8';
25     elseif rem == 1001 then
26         hex = hex+'9';
27     elseif rem == 1010 then
28         hex=hex+'A';
29     elseif rem == 1011 then
30         hex=hex+'B';
31     elseif rem == 1100 then
32         hex=hex+'C';
33     elseif rem == 1101 then
34         hex=hex+'D';
35     elseif rem == 1110 then
36         hex=hex+'E';
37     elseif rem == 1111 then
38         hex=hex+'F';
39     end // If statement ends
40     n = int(n/10000);
41 end // While loop ends
42 hex = strrev(hex); // Reverse string
43 endfunction
44
45 n = [10001100, 1011010111]; // Initialize the
    binary numbers
46 printf("\nThe hex equivalent of %d = %s", n(1),
    binary_hex(n(1)));
47 printf("\nThe hex equivalent of %d = %s", n(2),
    binary_hex(n(2)));
48
49 // Result
50 // The hex equivalent of 10001100 = 8C

```

51 // The hex equivalent of 1011010111 = 2D7

---

### Scilab code Exa 15.6 Hexadecimal equivalent of decimal numbers

```
1 // Scilab code Ex15.6 : Pg:772(2011)
2 clc;clear;
3 function hex = decimal_hex(n) // Function to convert
  decimal to hexadecimal
4   hex = emptystr();
5   while (n <>0)
6       rem = n-fix(n./16).*16;
7       if rem == 10 then
8           hex(i)=hex+'A';
9       elseif rem == 11 then
10          hex=hex+'B';
11         elseif rem == 12 then
12             hex=hex+'C';
13            elseif rem == 13 then
14                hex=hex+'D';
15               elseif rem == 14 then
16                   hex=hex+'E';
17                  elseif rem == 15 then
18                      hex=hex+'F';
19                 else
20                     hex=hex+string(rem);
21                 end
22             n = int(n/16);
23         end
24         hex = strrev(hex); // Reverse string
25     endfunction
26
27 n = 72905; // Initialize the binary numbers
28 printf("\nThe hex equivalent of %d = %s", n,
  decimal_hex(n));
29
```

```
30
31 // Result
32 // The hex equivalent of 72905 = 11CC9
```

---

### Scilab code Exa 15.7 Addition of two binary numbers

```
1 // Scilab code Ex15.7 : Pg:773(2011)
2 clc;clear;
3 function [bini]= decimal_binary(ni) // Function to
   convert decimal to binary
4     bini = 0;
5     i = 1;
6     while (ni <> 0)
7         rem = ni-fix(ni./2).*2;
8         ni = int(ni/2);
9         bini = bini + rem*i;
10        i = i * 10;
11    end
12 endfunction
13
14 function [deci]= binary_decimal(ni) // Function to
   convert binary to decimal
15     deci = 0;
16     i = 0;
17     while (ni <> 0)
18         rem = ni-fix(ni./10).*10;
19         ni = int(ni/10);
20         deci = deci + rem*2.^i;
21         i = i + 1;
22     end
23 endfunction
24
25 num1 = 11101; // Initialize the first binary
   number
26 num2 = 10111; // Initialize the second binary
```

```

        number
27
28 printf("%6d + %6d = %7d", num1, num2, decimal_binary
        (binary_decimal(num1)+binary_decimal(num2)));
29
30 // Result
31 // 11101 + 10111 = 110100

```

---

### Scilab code Exa 15.8 Addition of two binary numbers with fractions

```

1 // Scilab code Ex15.8 : Pg:773(2011)
2 clc;clear;
3 function [bini]= decimal_binary(ni) // Function to
    convert decimal to binary
4     bini = 0;
5     i = 1;
6     while (ni <> 0)
7         rem = ni-fix(ni./2).*2;
8         ni = int(ni/2);
9         bini = bini + rem*i;
10        i = i * 10;
11    end
12 endfunction
13
14 function [binf]= decifrac_binfrac(nf) // Function to
    convert binary fraction to decimal fraction
15    binf = 0; i = 0.1;
16    while (nf <> 0)
17        nf = nf*2;
18        rem = int(nf);
19        nf = nf-rem;
20        binf = binf + rem*i;
21        i = i/10;
22    end
23 endfunction

```

```

24
25 function [deci]= binary_decimal(ni) // Function to
    convert binary to decimal
26     deci = 0;
27     i = 0;
28     while (ni <> 0)
29         rem = ni-fix(ni./10).*10;
30         ni = int(ni/10);
31         deci = deci + rem*2.^i;
32         i = i + 1;
33     end
34 endfunction
35
36 function [decf]= binfrac_decifrac(nf) // Function to
    convert binary fraction to decimal fraction
37     decf = 0;
38     i = -1;
39     while (i >= -3)
40         nf = nf*10;
41         rem = round(nf);
42         nf = nf-rem;
43         decf = decf + rem*2.^i;
44         i = i - 1;
45     end
46 endfunction
47
48 bin1 = 1011.11; // Initialize the first binary
    binber
49 bin2 = 1011.01; // Initialize the second binary
    binber
50 bin1_int = int(bin1); // Extract the integral
    part for first
51 bin1_frac = bin1-bin1_int; // Extract the
    fractional part for second
52 bin2_int = int(bin2); // Extract the integral
    part for first
53 bin2_frac = bin2-bin2_int; // Extract the
    fractional part for second

```

```

54 dec1 = binary_decimal(bin1_int)+binfrac_decifrac(
    bin1_frac);
55 dec2 = binary_decimal(bin2_int)+binfrac_decifrac(
    bin2_frac);
56 dec = dec1+dec2;
57 dec_int = int(dec);
58 dec_frac = dec-dec_int;
59 printf("%7.2f + %7.2f = %8.2f", bin1, bin2,
    decimal_binary(dec_int)+decifrac_binfrac(dec_frac
    ));
60
61 // Result
62 // 1011.11 + 1011.01 = 10111.00

```

---

#### Scilab code Exa 15.9 Subtraction of two binary numbers

```

1 // Scilab code Ex15.9 : Pg:773(2011)
2 clc;clear;
3 function [bini]= decimal_binary(ni) // Function to
    convert decimal to binary
4     bini = 0;
5     i = 1;
6     while (ni <> 0)
7         rem = ni-fix(ni./2).*2;
8         ni = int(ni/2);
9         bini = bini + rem*i;
10        i = i * 10;
11    end
12 endfunction
13
14 function [deci]= binary_decimal(ni) // Function to
    convert binary to decimal
15     deci = 0;
16     i = 0;
17     while (ni <> 0)

```

```

18     rem = ni-fix(ni./10).*10;
19     ni = int(ni/10);
20     deci = deci + rem*2.^i;
21     i = i + 1;
22     end
23 endfunction
24
25 num1 = 1001;    // Initialize the first binary
                number
26 num2 = 0111;    // Initialize the second binary
                number
27
28 printf("%4d - 0%3d = 00%2d", num1, num2,
        decimal_binary(binary_decimal(num1)-
        binary_decimal(num2)));
29
30 // Result
31 // 1001 - 0111 = 0010

```

---

**Scilab code Exa 15.10** Multiplication of two binary numbers

```

1 // Scilab code Ex15.10 :Pg:773(2008)
2 clc;clear;
3 function [bini]= decimal_binary(ni) // Function to
    convert decimal to binary
4     bini = 0;
5     i = 1;
6     while (ni <> 0)
7         rem = ni-fix(ni./2).*2;
8         ni = int(ni/2);
9         bini = bini + rem*i;
10        i = i * 10;
11    end
12 endfunction
13

```



```

14 function [deci]= binary_decimal(ni) // Function to
    convert binary to decimal
15     deci = 0;
16     i = 0;
17     while (ni <> 0)
18         rem = ni-fix(ni./10).*10;
19         ni = int(ni/10);
20         deci = deci + rem*2.^i;
21         i = i + 1;
22     end
23 endfunction
24
25 function binp = bin_product(op1, op2)
26     binp = decimal_binary(binary_decimal(op1)*
        binary_decimal(op2));
27 endfunction
28
29 mul1 = 1101; // Initialize the first binary
    multiplicand
30 mul2 = 1100; // Initialize the second binary
    multiplicand
31 product = bin_product(mul1, mul2);
32
33 printf("%4d X %4d = %8d", mul1, mul2, product);
34
35 // Result
36 // 1101 X 1100 = 10011100

```

---

### Scilab code Exa 15.11 Binary division of two numbers

```

1 // Scilab code Ex15.11 : Pg:774(2008)
2 clc;clear;
3 function [bini]= decimal_binary(ni) // Function to
    convert decimal to binary
4     bini = 0;

```

```

5     i = 1;
6     while (ni <> 0)
7         rem = ni-fix(ni./2).*2;
8         ni = int(ni/2);
9         bini = bini + rem*i;
10        i = i * 10;
11    end
12 endfunction
13
14 function [deci]= binary_decimal(ni) // Function to
    convert binary to decimal
15     deci = 0;
16     i = 0;
17     while (ni <> 0)
18         rem = ni-fix(ni./10).*10;
19         ni = int(ni/10);
20         deci = deci + rem*2.^i;
21         i = i + 1;
22     end
23 endfunction
24
25 function binp = bin_division(op1, op2)
26     binp = decimal_binary(binary_decimal(op1)/
        binary_decimal(op2));
27 endfunction
28
29 dividend = 11001 ; // Initialize the first binary
    multiplicand
30 divisor = 101; // Initialize the second binary
    multiplicand
31 product = bin_division(dividend, divisor);
32
33 printf("%5d divided by %3d gives %3d", dividend,
    divisor, product);
34
35 // Result
36 // 11001 divided by 101 gives 101

```

---

# Chapter 16

## Crystal Physics

Scilab code Exa 16.1 Miller indices of the crystal plane

```
1 // Scilab Code Ex16.1 : Page-820 (2011)
2 clc; clear;
3 p = 1; q = 2; r = 3; // Coefficients of intercepts
   along three axes
4 p_inv = 1/p; // Reciprocate the first
   coefficient
5 q_inv = 1/q; // Reciprocate the second
   coefficient
6 r_inv = 1/r; // Reciprocate the third
   coefficient
7 mul_fact = double(lcm(int32([p,q,r]))); // Find l.c.
   m. of m,n and p
8 m1 = p_inv*mul_fact; // Clear the first fraction
9 m2 = q_inv*mul_fact; // Clear the second fraction
10 m3 = r_inv*mul_fact; // Clear the third fraction
11 printf("\\nThe required miller indices are : (%d %d
   %d) ", m1,m2,m3);
12
13 // Result
14 // The required miller indices are : (6 3 2)
```

---

**Scilab code Exa 16.2** Miller indices of the lattice plane

```
1 // Scilab Code Ex16.2 : Page-820 (2011)
2 clc; clear;
3 p = 2; q = 3; r = -4; // Coefficients of intercepts
   along three axes
4 p_inv = 1/p;          // Reciprocate the first
   coefficient
5 q_inv = 1/q;          // Reciprocate the second
   coefficient
6 r_inv = 1/r;          // Reciprocate the third
   coefficient
7 mul_fact = double(lcm(int32([p,q,abs(r)]))); // Find
   l.c.m. of m,n and p
8 m1 = p_inv*mul_fact;  // Clear the first fraction
9 m2 = q_inv*mul_fact;  // Clear the second fraction
10 m3 = r_inv*mul_fact; // Clear the third fraction
11 printf("\\nThe miller indices of laticce plane are :
   (%d %d %d) ", m1,m2,m3);
12
13 // Result
14 // The miller indices of laticce plane are : (6 4
   -3)
```

---

**Scilab code Exa 16.3** Miller indices of the set of parallel planes

```
1 // Scilab Code Ex16.3 : Page-821 (2011)
2 clc; clear;
3 p = 3; q = 4; r = %inf; // Coefficients of
   intercepts along three axes
4 p_inv = 1/p;          // Reciprocate the first
   coefficient
```

```

5 q_inv = 1/q;          // Reciprocate the second
   coefficient
6 r_inv = 1/r;          // Reciprocate the third
   coefficient
7 mul_fact = double(lcm(int32([p,q]))); // Find l.c.m.
   of m,n and p
8 m1 = p_inv*mul_fact; // Clear the first fraction
9 m2 = q_inv*mul_fact; // Clear the second fraction
10 m3 = r_inv*mul_fact; // Clear the third fraction
11 printf("\nThe miller indices of the given planes are
   : (%d %d %d) ", m1,m2,m3);
12
13 // Result
14 // The miller indices of the given planes are : (4 3
   0)

```

---

**Scilab code Exa 16.4** Length of the intercepts on Y and Z axes

```

1 // Scilab Code Ex16.4 : Page-822 (2011)
2 clc; clear;
3 p = 1.2; // First coefficient of intercept along X-
   axis , angstrom
4 a = 1.2, b = 1.8, c = 2.0; // Lattice parameters
   along three axes , angstrom
5 h = 2, k = 3, l = 1; // Miller indices of lattice
   plane
6 // As p:q:r = a/h:b/k:c/l, solving for q and r
7 q = p*(b/k)/(a/h); // Second coefficient of
   intercept along X-axis , angstrom
8 r = p*(c/l)/(a/h); // Third coefficient of intercept
   along X-axis , angstrom
9 printf("\nThe lengths of the intercepts on Y and Z
   axes are %3.1f angstrom and %3.1f angstrom
   respectively", q, r);
10

```

```

11 // Result
12 // The lengths of the intercepts on Y and Z axes are
    1.2 angstrom and 4.0 angstrom respectively

```

---

**Scilab code Exa 16.5** Lattice constant for NaCl crystal

```

1 // Scilab Code Ex16.5 : Page-822 (2011)
2 clc; clear;
3 M = 58.5; // Molecular weight of NaCl, g-mole
4 rho = 2.198e+03; // Density of Nacl, kg per metre
    cube
5 n = 4; // No. of atoms per unit cell for an fcc
    lattice of NaCl crystal
6 NA = 6.023D+26; // Avogadro's No., atoms/k-mol
7 // Volume of the unit cell is given by
8 //  $a^3 = M*n/(N*d)$ 
9 // Solving for a
10 a = (n*M/(rho*NA))^(1/3); // Lattice constant of
    unit cell of NaCl
11 printf("\\nLattice constant for the NaCl crystal = %4
    .2f angstrom", a/1e-010);
12
13 // Result
14 // Lattice constant for the NaCl crystal = 5.61
    angstrom

```

---

**Scilab code Exa 16.6** Lattice constant for KBr crystal

```

1 // Scilab Code Ex16.6 :Page-823 (2011)
2 clc; clear;
3 M = 119; // Molecular weight of KBr, g-mole
4 rho = 2.7; // Density of KBr, g per cm-cube

```

```

5 n = 4;      // No. of atoms per unit cell for an fcc
   lattice of KBr crystal
6 NA = 6.023D+23;    // Avogadro's No., atoms/mol
7 // Volume of the unit cell is given by
8 //  $a^3 = M*n/(N*d)$ 
9 // Solving for a
10 a = (n*M/(rho*NA))^(1/3);    // Lattice constant of
   unit cell of KBr
11 printf("\nLattice constant for the KBr crystal = %4
   .2f angstrom", a/1e-008);
12
13 // Result
14 // Lattice constant for the KBr crystal = 6.64
   angstrom

```

---

**Scilab code Exa 16.7** Lattice constant for Cu and distance between the two nearest Cu atoms

```

1 // Scilab Code Ex16.7 : Page-823 (2011)
2 clc; clear;
3 M = 63.5;      // Molecular weight of Cu, g-mole
4 rho = 8.96;    // Density of Cu, g per cm-cube
5 n = 4;      // No. of atoms per unit cell for an fcc
   lattice of Cu
6 NA = 6.023D+23;    // Avogadro's No., atoms/mol
7 // Volume of the unit cell is given by
8 //  $a^3 = M*n/(N*d)$ 
9 // Solving for a
10 a = (n*M/(rho*NA))^(1/3);    // Lattice constant of
   unit cell of Cu
11 d = a/sqrt(2);    // Distance between the two
   nearest Cu atoms, angstrom
12 printf("\nLattice constant for the Cu crystal = %4.2
   f angstrom", a/1e-008);
13 printf("\nThe distance between the two nearest Cu

```

```

        atoms = %4.2f angstrom", d/1e-008);
14
15 // Result
16 // Lattice constant for the Cu crystal = 3.61
    angstrom
17 // The distance between the two nearest Cu atoms =
    2.55 angstrom

```

---

### Scilab code Exa 16.8 Inter planar spacing for lattice planes

```

1 // Scilab Code Ex16.8 : Page-824 (2011)
2 clc; clear;
3 a = 1; // For simplicity assume lattice parameter
    of cubic crystal to be unity, unit
4 // For (011) planes
5 h = 0; k = 1; l = 1; // Miller Indices for planes in
    a cubic crystal
6 d_011 = a/(h^2+k^2+l^2)^(1/2); // The interplanar
    spacing for cubic crystals, m
7 printf("\nThe interplanar spacing between
    consecutive (011) planes = a/sqrt(%d)", 1/d_011
    ^2);
8
9 // For (101) planes
10 h = 1; k = 0; l = 1; // Miller Indices for planes in
    a cubic crystal
11 d_101 = a/(h^2+k^2+l^2)^(1/2); // The interplanar
    spacing for cubic crystals, m
12 printf("\nThe interplanar spacing between
    consecutive (101) planes = a/sqrt(%d)", 1/d_101
    ^2);
13
14 // For (112) planes
15 h = 1; k = 1; l = 2; // Miller Indices for planes in
    a cubic crystal

```



```

16 d_112 = a/(h^2+k^2+l^2)^(1/2); // The interplanar
    spacing for cubic crystals , m
17 printf("\nThe interplanar spacing between
    consecutive (112) planes = a/sqrt(%d)", 1/d_112
    ^2);
18
19 // Result
20 // The interplanar spacing between consecutive (011)
    planes = a/sqrt(2)
21 // The interplanar spacing between consecutive (101)
    planes = a/sqrt(2)
22 // The interplanar spacing between consecutive (112)
    planes = a/sqrt(5)

```

---

**Scilab code Exa 16.9** Interplanar spacing in cubic crystal

```

1 // Scilab Code Ex16.9 : Page-824 (2011)
2 clc; clear;
3 a = 4.2e-010; // Lattice parameter of cubic
    crystal , m
4 h = 3; k = 2; l = 1; // Miller Indices for planes in
    a cubic crystal
5 d_321 = a/(h^2+k^2+l^2)^(1/2); // The interplanar
    spacing for cubic crystals , m
6 printf("\nThe interplanar spacing between
    consecutive (321) planes = %4.2f angstrom", d_321
    /1e-010);
7
8 // Result
9 // The interplanar spacing between consecutive (321)
    planes = 1.12 angstrom

```

---

**Scilab code Exa 16.10** Interplanar spacing in tetragonal crystal lattice

```

1 // Scilab Code Ex16.10 : Page-824 (2011)
2 clc; clear;
3 a = 2.5, b = 2.5, c = 1.8; // Lattice parameter
   of tetragonal crystal, angstrom
4 h = 1; k = 1; l = 1; // Miller Indices for planes in
   a tetragonal crystal
5 d_hkl = 1/sqrt((h/a)^2+(k/b)^2+(l/c)^2); // The
   interplanar spacing for tetragonal crystals, m
6 printf("\nThe interplanar spacing between
   consecutive (111) planes = %4.2f angstrom", d_hkl
   );
7
8 // Result
9 // The interplanar spacing between consecutive (111)
   planes = 1.26 angstrom

```

---

# Chapter 17

## Nuclear Physics

Scilab code Exa 17.1 Binding energy per nucleon for the deuteron

```
1 // Scilab code Ex17.1 : Pg:888 (2011)
2 clc; clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 m_n = 1.675e-027; // Mass of the neutron, kg
5 m_p = 1.672e-027; // Mass of the proton, kg
6 M_D = 3.343e-027; // Mass of the deuteron, kg
7 c = 3e+08; // Speed of light, m/s
8 delta_m = m_n + m_p - M_D; // Mass defect in the
   formation of deuterium, kg
9 BE = delta_m*c^2; // Binding energy of the
   deuterium, J
10 BE_bar = BE/2; // Binding energy per nucleon of
   deuterium, J
11 printf("\\nBinding energy per nucleon for the deuteron
   = %5.3 f MeV", BE_bar/(e*1e+06));
12
13 // Result
14 // Binding energy per nucleon for the deuteron =
   1.125 MeV
```

---

### Scilab code Exa 17.2 Binding energy for an alpha particle

```
1 // Scilab code Ex17.2 : Pg:889 (2011)
2 clc;clear;
3 amu = 931.5; // Energy equivalent of 1 amu, MeV
4 m_n = 1.008665; // Mass of the neutron, amu
5 m_p = 1.007825; // Mass of the proton, amu
6 M_He = 4.002870; // Mass of the helium nucleus,
   amu
7 c = 3e+08; // Speed of light, m/s
8 BE = (2*m_n+2*m_p - M_He)*amu; // Binding energy
   for the alpha particle, MeV
9 printf("\\nThe binding energy for the alpha particle
   = %2d MeV", BE);
10
11 // Result
12 // The binding energy for the alpha particle = 28
   MeV
```

---

### Scilab code Exa 17.3 Weizsacker formula for stability of nuclei

```
1 // Scilab code Ex17.3 :Pg:889 (2011)
2 clc;clear;
3 A = 1; // For simplicity assume mass number to be
   unity
4 nucleus = cell(4,3);
5 nucleus(1,1).entries = 'He';
6 nucleus(1,2).entries = 2;
7 nucleus(1,3).entries = 6;
8 nucleus(2,1).entries = 'Be';
9 nucleus(2,2).entries = 4;
10 nucleus(2,3).entries = 6;
```

```

11 nucleus(3,1).entries = 'Li';
12 nucleus(3,2).entries = 3;
13 nucleus(3,3).entries = 6;
14 a_c = 0.7053; // Asymmetry energy constant, MeV
15 a_a = 23.702; // Coulomb energy constant, MeV
16 Z = A/(2+a_c/(2*a_a)*A^(2/3));
17 for i = 1:1:3
18     if abs(nucleus(i,2).entries/nucleus(i,3).entries
19         - Z) < 0.005 then
20         printf("\n%s(%d,%d) is more stable than
21             other two nuclei", nucleus(i,1).entries,
22             nucleus(i,2).entries, nucleus(i,3).
23             entries);
24     end
25 end
26 // Result
27 // Li(3,6) is more stable than other two nuclei

```

---

#### Scilab code Exa 17.4 Nuclei stability

```

1 // Scilab code Ex17.4 : Pg:890 (2011)
2 clc;clear;
3 nucleus = cell(4,3);
4 // For Li nuclides
5 nucleus(1,1).entries = 'Li';
6 nucleus(1,2).entries = 3;
7 nucleus(1,3).entries = 7;
8 nucleus(2,1).entries = 'Li';
9 nucleus(2,2).entries = 3;
10 nucleus(2,3).entries = 8;
11 // For Be nuclides
12 nucleus(3,1).entries = 'Be';
13 nucleus(3,2).entries = 4;
14 nucleus(3,3).entries = 9;

```

```

15 nucleus(4,1).entries = 'Be';
16 nucleus(4,2).entries = 4;
17 nucleus(4,3).entries = 10;
18 a_c = 0.7053; // Asymmetry energy constant, MeV
19 a_a = 23.702; // Coulomb energy constant, MeV
20 for i = 1:1:4
21     Z = nucleus(i,3).entries/(2+a_c/(2*a_a)*nucleus(
        i,3).entries^(2/3));
22     if abs(Z-int(Z)) < 0.5 then
23         printf("\n%s(%d,%d) is more stable", nucleus
            (i,1).entries, nucleus(i,2).entries,
            nucleus(i,3).entries);
24     end
25 end
26
27 // Result
28 // Li(3,7) is more stable
29 // Be(4,9) is more stable

```

---

**Scilab code Exa 17.5** Exothermicity and endothermicity of nuclear reactions

```

1 // Scilab code Ex17.5 : Pg:891 (2011)
2 clc;clear;
3 c = 1; // For simplicity assume speed of light to
        be unity, unit
4 nucleus = cell(4,4);
5 // For first reaction
6 nucleus(1,1).entries = 'N';
7 nucleus(1,2).entries = 7;
8 nucleus(1,3).entries = 14;
9 nucleus(1,4).entries = 14.00753;
10 nucleus(2,1).entries = 'He';
11 nucleus(2,2).entries = 2;
12 nucleus(2,3).entries = 4;

```

```

13 nucleus(2,4).entries = 4.00206;
14 nucleus(3,1).entries = 'O';
15 nucleus(3,2).entries = 8;
16 nucleus(3,3).entries = 17;
17 nucleus(3,4).entries = 17.00450;
18 nucleus(4,1).entries = 'H';
19 nucleus(4,2).entries = 1;
20 nucleus(4,3).entries = 1;
21 nucleus(4,4).entries = 1.00814;
22 Q = (nucleus(1,4).entries + nucleus(2,4).entries)*c
      ^2 - (nucleus(3,4).entries + nucleus(4,4).entries
      )*c^2;
23 if Q < 0 then
24     T_state = "endothermic";
25     elseif Q > 0
26     T_state = "exothermic";
27 end
28 printf("\nThe reaction");
29 printf("\n%s(%d,%d) + %s(%d,%d) --> %s(%d,%d) + %s(
      %d,%d) is %s", nucleus(1,1).entries, nucleus(1,2)
      .entries, nucleus(1,3).entries, nucleus(2,1).
      entries, nucleus(2,2).entries, nucleus(2,3).
      entries, nucleus(3,1).entries, nucleus(3,2).
      entries, nucleus(3,3).entries, nucleus(4,1).
      entries, nucleus(4,2).entries, nucleus(4,3).
      entries, T_state);
30 // For second reaction
31 nucleus(1,1).entries = 'Li';
32 nucleus(1,2).entries = 3;
33 nucleus(1,3).entries = 7;
34 nucleus(1,4).entries = 7.01822;
35 nucleus(2,1).entries = 'H';
36 nucleus(2,2).entries = 1;
37 nucleus(2,3).entries = 1;
38 nucleus(2,4).entries = 1.00814;
39 nucleus(3,1).entries = 'He';
40 nucleus(3,2).entries = 2;
41 nucleus(3,3).entries = 4;

```

```

42 nucleus(3,4).entries = 4.00206;
43 Q = (nucleus(1,4).entries + nucleus(2,4).entries)*c
      ^2 - (nucleus(3,4).entries + nucleus(3,4).entries
      )*c^2;
44 if Q < 0 then
45     T_state = "endothermic";
46 elseif Q > 0
47     T_state = "exothermic";
48 end
49 printf("\nThe reaction");
50 printf("\n%s(%d,%d) + %s(%d,%d) --> %s(%d,%d) + %s(
      %d,%d) is %s", nucleus(1,1).entries, nucleus(1,2)
      .entries, nucleus(1,3).entries, nucleus(2,1).
      entries, nucleus(2,2).entries, nucleus(2,3).
      entries, nucleus(3,1).entries, nucleus(3,2).
      entries, nucleus(3,3).entries, nucleus(4,1).
      entries, nucleus(4,2).entries, nucleus(4,3).
      entries, T_state);
51
52 // Result
53 //
54 // The reaction
55 // N(7,14) + He(2,4) --> O(8,17) + H(1,1) is
      endothermic
56 // The reaction
57 // Li(3,7) + H(1,1) --> He(2,4) + H(1,1) is
      exothermic

```

---

**Scilab code Exa 17.6** Q value of the formation of P30 in ground state

```

1 // Scilab code Ex17.6 : Pg:891 (2011)
2 clc;clear;
3 nucleus = cell(4,3);
4 nucleus(1,1).entries = 'Si';
5 nucleus(1,2).entries = 14;

```



```

6 nucleus(1,3).entries = 29;
7 nucleus(2,1).entries = 'H';
8 nucleus(2,2).entries = 1;
9 nucleus(2,3).entries = 2;
10 nucleus(3,1).entries = 'P';
11 nucleus(3,2).entries = 15;
12 nucleus(3,3).entries = 30;
13 nucleus(4,1).entries = 'n';
14 nucleus(4,2).entries = 0;
15 nucleus(4,3).entries = 1;
16 Q = 2*23.834-44.359; // Q-value of the reaction ,
    MeV
17 printf("\nThe reaction");
18 printf("\n%s(%d,%d) + %s(%d,%d) --> %s(%d,%d) + %s(
    %d,%d)", nucleus(1,1).entries, nucleus(1,2).
    entries, nucleus(1,3).entries, nucleus(2,1).
    entries, nucleus(2,2).entries, nucleus(2,3).
    entries, nucleus(3,1).entries, nucleus(3,2).
    entries, nucleus(3,3).entries, nucleus(4,1).
    entries, nucleus(4,2).entries, nucleus(4,3).
    entries);
19 printf("\nhas the Q-value : %5.3f MeV", Q);
20
21 // Result
22 // The reaction
23 // Si(14,29) + H(1,2) --> P(15,30) + n(0,1)
24 // has the Q-value : 3.309 MeV

```

---

**Scilab code Exa 17.7** Threshold energy required to initiate the reaction

```

1 // Scilab code Ex17.7 : Pg:892 (2011)
2 clc;clear;
3 amu = 931.5; // Energy equivalent of 1 amu, MeV
4 nucleus = cell(4,3);
5 nucleus(1,1).entries = 'P';

```

```

6 nucleus(1,2).entries = 15;
7 nucleus(1,3).entries = 31;
8 nucleus(1,4).entries = 30.98356;
9 nucleus(2,1).entries = 'n';
10 nucleus(2,2).entries = 0;
11 nucleus(2,3).entries = 1;
12 nucleus(2,4).entries = 1.00898;
13 nucleus(3,1).entries = 'Si';
14 nucleus(3,2).entries = 14;
15 nucleus(3,3).entries = 31;
16 nucleus(3,4).entries = 30.98515;
17 nucleus(4,1).entries = 'p';
18 nucleus(4,2).entries = 1;
19 nucleus(4,3).entries = 1;
20 nucleus(4,4).entries = 1.00814;
21 Q = ((nucleus(1,4).entries + nucleus(2,4).entries)-(
        nucleus(3,4).entries + nucleus(4,4).entries))*amu
        ; // Q-value of the reaction , MeV
22 E_th = -1*Q*(nucleus(1,4).entries+nucleus(2,4).
        entries)/nucleus(1,4).entries;
23 printf("\nThe threshold energy required to initiate
        the reaction");
24 printf("\n\t%s(%d,%d) + %s(%d,%d) --> %s(%d,%d) + %s
        (%d,%d)", nucleus(1,1).entries, nucleus(1,2).
        entries, nucleus(1,3).entries, nucleus(2,1).
        entries, nucleus(2,2).entries, nucleus(2,3).
        entries, nucleus(3,1).entries, nucleus(3,2).
        entries, nucleus(3,3).entries, nucleus(4,1).
        entries, nucleus(4,2).entries, nucleus(4,3).
        entries);
25 printf("\nis %5.3f MeV", E_th);
26
27 // Result
28 // The threshold energy required to initiate the
        reaction
29 // P(15,31) + n(0,1) --> Si(14,31) + p(1,1)
30 // is 0.721 MeV

```

**Scilab code Exa 17.8** Kinetic energy of emitted protons and threshold energy

```

1 // Scilab code Ex17.8 : Pg:892 (2011)
2 clc;clear;
3 amu = 931.5; // Energy equivalent of 1 amu, MeV
4 nucleus = cell(4,3);
5 nucleus(1,1).entries = 'F';
6 nucleus(1,2).entries = 9;
7 nucleus(1,3).entries = 19;
8 M_P = 19.0457; // Mass of product nucleus, amu
9 nucleus(2,1).entries = 'n';
10 nucleus(2,2).entries = 0;
11 nucleus(2,3).entries = 1;
12 m_i = 1.0087; // Mass of incident particle, amu
13 nucleus(3,1).entries = 'O';
14 nucleus(3,2).entries = 8;
15 nucleus(3,3).entries = 19;
16 nucleus(4,1).entries = 'H';
17 nucleus(4,2).entries = 1;
18 nucleus(4,3).entries = 1;
19 m_e = 1.00728; // Mass of emitted particle, amu
20 K_i = 15; // Kinetic energy of incident neutrons,
    MeV
21 Q = -7.6342; // Q-value of the reaction, MeV
22 K_e = (Q*M_P-(m_i-M_P)*K_i)/(m_e+M_P); // Kinetic
    energy of emitted photon, MeV
23 E_th = -1*Q*(M_P+m_i)/M_P; // Threshold energy
    required to initiate the reaction, MeV
24 printf("\nThe kinetic energy of emitted photon = %5
    .3f MeV", K_e);
25 printf("\nThe threshold energy required to initiate
    the reaction");
26 printf("\n\t%s(%d,%d) + %s(%d,%d) --> %s(%d,%d) + %s

```

```

    (%d,%d)", nucleus(1,1).entries, nucleus(1,2).
    entries, nucleus(1,3).entries, nucleus(2,1).
    entries, nucleus(2,2).entries, nucleus(2,3).
    entries, nucleus(3,1).entries, nucleus(3,2).
    entries, nucleus(3,3).entries, nucleus(4,1).
    entries, nucleus(4,2).entries, nucleus(4,3).
    entries);
27 printf("\nis %5.3f MeV", E_th);
28
29 // Result
30 // The kinetic energy of emitted photon = 6.241 MeV
31 // The threshold energy required to initiate the
    reaction
32 //  $F(9,19) + n(0,1) \rightarrow O(8,19) + H(1,1)$ 
33 // is 8.039 MeV

```

---

**Scilab code Exa 17.9** Energy released by the fission of 1 kg of U235

```

1 // Scilab code Ex17.9 : Pg:893 (2011)
2 clc;clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 N_A = 6.023e+023; // Avogadro's number
5 E_f = 200*1e+06*e; // Energy released per fission
    , J
6 E_mol = E_f*N_A; // Energy released by one mole
    of U235, J
7 E = E_mol*1000/235; // Energy released by the
    fission of 1 kg of U235, J
8 printf("\nThe Energy released by the fission of 1 kg
    of U235 = %4.2e kWh", E/(1000*3600));
9
10 // Result
11 // The Energy released by the fission of 1 kg of
    U235 = 2.28e+007 kWh

```

---

**Scilab code Exa 17.10** Rate of fission of U235

```
1 // Scilab code Ex17.10 : Pg:894 (2011)
2 clc;clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 E = 3.2e+07; // Energy released per second by the
   reactor, J
5 E_f = 200*1e+06*e; // Energy released per fission
   , J
6 N = E/E_f; // Number of fissions per second of
   U235, per second
7 printf("\nThe number of U235 atoms undergoing
   fissions per second = %1.0e", N);
8
9 // Result
10 // The number of U235 atoms undergoing fissions per
   second = 1e+018
```

---

**Scilab code Exa 17.11** Rate of fission and energy released in the complete fissioning of U235

```
1 // Scilab code Ex17.11 : Pg:894 (2011)
2 clc;clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 N_A = 6.023e+026; // Avogadro's number, per kmol
5 P = 2; // Power produced by the fission of U235,
   watt
6 E_f = 200*1e+06*e; // Energy released per fission
   , J
7 FR = P/E_f; // Fission rate of U235, fission/sec
8 N = 0.5/235*N_A; // Number of U235 nuclei in 0.5
   kg of U235
```

```

9 E = 200*N; // Energy released in the complete
    fissioning of 0.5 kg of U235, MeV
10 printf("\nThe fission rate of U235 = %4.2e fissions/
    sec", FR);
11 printf("\nThe energy released in the complete
    fissioning of 0.5 kg of U235 = %1.0e kcal", E*1e
    +06*e/(1000*4.186));
12
13 // Result
14 // The fission rate of U235 = 6.25e+010 fissions/sec
15 // The energy released in the complete fissioning of
    0.5 kg of U235 = 1e+010 kcal

```

---

**Scilab code Exa 17.12** Energy and oscillator frequency of some positively charged particles accelerating in a cyclotron

```

1 // Scilab code Ex17.12 : Pg:894 (2011)
2 clc;clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 R_max = 0.6; // Radius of two dees of the
    cyclotron, m
5 B = 1.6; // Strength of pole pieces of the
    cyclotron, tesla
6 // For proton
7 m = 1.67e-027; // Mass of the proton, kg
8 q = 1.6e-019; // Charge on a proton, C
9 E = 1/2*q^2*R_max^2*B^2/(m*e*1e+06); // Energy of
    the proton, MeV
10 f_proton = q*B/(2*pi*m*1e+06); // Cyclotron
    oscillator frequency for the proton, MHz
11 printf("\nEnergy of the proton = %5.2f MeV", E);
12 printf("\nCyclotron frequency for proton = %5.2f MHz
    ", f_proton);
13 // For deuteron
14 m = 2*1.67e-027; // Mass of the deuteron, kg

```

```

15 q = 1.6e-019;    // Charge on a deuteron , C
16 E = 1/2*q^2*R_max^2*B^2/(m*e*1e+06);    // Energy of
    the deuteron , MeV
17 f_deuteron = q*B/(2*pi*m*1e+06);    // Cyclotron
    oscillator frequency for the deuteron , MHz
18 printf("\nEnergy of the deuteron = %5.2f MeV", E);
19 printf("\nCyclotron frequency for deuteron = %5.2f
    MHz", f_deuteron);
20 // For alpha-particle
21 m = 4*1.67e-027;    // Mass of the alpha-particle ,
    kg
22 q = 2*1.6e-019;    // Charge on a alpha-particle , C
23 E = 1/2*q^2*R_max^2*B^2/(m*e*1e+06);    // Energy of
    the deuteron , MeV
24 f_alpha = q*B/(2*pi*m*1e+06);    // Cyclotron
    oscillator frequency for the alpha-particle , MHz
25 printf("\nEnergy of the alpha-particle = %5.2f MeV",
    E);
26 printf("\nCyclotron frequency for alpha-particle =
    %5.2f MHz", f_alpha);
27
28 // Result
29 // Energy of the proton = 44.15 MeV
30 // Cyclotron frequency for proton = 24.40 MHz
31 // Energy of the deuteron = 22.07 MeV
32 // Cyclotron frequency for deuteron = 12.20 MHz
33 // Energy of the alpha-particle = 44.15 MeV
34 // Cyclotron frequency for alpha-particle = 12.20
    MHz

```

---

**Scilab code Exa 17.13** Energy of the protons issuing out of the cyclotron

```

1 // Scilab code Ex17.13 : Pg:895 (2011)
2 clc;clear;
3 e = 1.6e-019;    // Energy equivalent of 1 eV, J/eV

```

```

4 R_max = 0.75;    // Radius of two dees of the
    cyclotron , m
5 f = 15e+06;    // Frequency of alternating potential
    , Hz
6 m = 1.67e-027;    // Mass of the proton , kg
7 // As  $E = 1/2 * q^2 * R_{max}^2 * B^2 / (m * e)$  and  $f = q * B / (2 *$ 
     $\%pi * m)$ , solving for E
8 E = 2 * %pi^2 * m * f^2 * R_max^2 / (e * 1e+06);
9 disp(E)
10 printf("\nEnergy of the protons issuing out of the
    cyclotron = %6.4f MeV", E);
11
12 // Result
13 // Energy of the protons issuing out of the
    cyclotron = 26.0754 MeV

```

---

**Scilab code Exa 17.14** Energy gained per turn and maximum energy of the electron in a betatron

```

1 // Scilab code Ex17.14 : Pg:896 (2011)
2 clc;clear;
3 e = 1.6e-019;    // Charge on an electron , C
4 c = 3e+08;    // Speed of light , m/s
5 B_orbit = 0.5;    // Magnetic field at the orbit of
    the betatron , T
6 f = 60;    // Operating frequency of the betatron ,
    Hz
7 omega = 2 * %pi * f;    // Angular frequency of
    operation , rad/s
8 r = 1.6/2;    // Radius of stable orbit , m
9 K_av = 4 * omega * e * r^2 * B_orbit / 1.6e-019;    // Average
    energy gained by the electron per turn , eV
10 K_max = c * e * r * B_orbit / 1.6e-019;    // Maximum energy
    gained by the electron , eV
11 printf("\nThe average energy gained by the electron

```



```

    per turn = %5.1f eV", K_av);
12 printf("\nThe maximum energy gained by the electron
    = %5.1e eV", K_max);
13
14 // Result
15 // The average energy gained by the electron per
    turn = 482.5 eV
16 // The maximum energy gained by the electron = 1.2e
    +008 eV

```

---

**Scilab code Exa 17.15** Maximum frequency of Dee voltage and gain in energy of the deuteron in a synchrocyclotron

```

1 // Scilab code Ex17.15 : Pg:896 (2011)
2 clc;clear;
3 q = 1.6e-019; // Charge on a deuteron , C
4 amu = 931.5; // Energy equivalent of 1 amu, MeV
5 m0 = 2.0141; // Rest mass of a deuteron , kg
6 B0 = 1.5; // Magnetic field at the centre of the
    synchrocyclotron , T
7 B = 1.431; // Magnetic field at the periphery of
    the synchrocyclotron , T
8 f0 = q*B0/(2*3.14*m0*1.67e-027*1e+06); // Maximum
    frequency of Dee voltage of synhrocyclotron , MHz
9 f = 1e+07; // Minimum frequency of Dee voltage ,
    Hz
10 m = q*B/(2*3.14*f*1.67e-027); // Mass of deuteron
    at the periphery of the Dee, amu
11 K = (m-m0)*amu; // Gain in energy of the deuteron
    , MeV
12 printf("\nThe maximum frequency of Dee voltage = %5
    .2f MHz", f0);
13 printf("\nThe gain in energy of the deuteron = %6.2 f
    MeV", K);
14

```

```

15 // Result
16 // The maximum frequency of Dee voltage = 11.36 MHz
17 // The gain in energy of the deuteron = 157.47 MeV

```

---

**Scilab code Exa 17.16** Maximum radial field and the life of a GM counter

```

1 // Scilab code Ex17.16 : Pg:897 (2011)
2 clc;clear;
3 V = 1000; // Operating voltage of the GM counter ,
    volt
4 a = 1e-04; // Radius of GM counter wire , m
5 b = 2e-02; // Radius of cathode , m
6 E = V/(2.3026*a*log10(b/a)); // Maximum radial
    field at the surface of central wire of GM tube ,
    V/m
7 tau = 1e+09; // Life time of GM tube , counts
8 N = tau/(50*60*60*2000); // Life of the GM
    counter , years
9 printf("\nThe maximum radial field at the surface of
    central wire of GM tube = %4.2e V/m", E);
10 printf("\nThe life of the GM counter = %4.2f years",
    N);
11
12 // Result
13 // The maximum radial field at the surface of
    central wire of GM tube = 1.89e+006 V/m
14 // The life of the GM counter = 2.78 years

```

---

**Scilab code Exa 17.17** Avalanche voltage in a GM tube

```

1 // Scilab code Ex17.17 : Pg:898 (2011)
2 clc;clear;

```

```

3 I = 15.7;    // Ionization potential of argon in GM
  counter, volt
4 a = 0.012/2*1e-02;    // Radius of GM counter wire,
  m
5 b = 5/2*1e-02;    // Radius of cathode, m
6 lambda = 7.8e-006;    // Mean free path of argon in
  GM counter, m
7 // As  $E \cdot \lambda = I = V \cdot \lambda / (2.3026 \cdot a \cdot \log_{10}(b/a))$ ,
  solving for V
8 V = 2.3026*a*I*log10(b/a)/lambda;    // Voltage that
  must be applied to produce an avalanche in GM
  tube, volt
9 printf("\nThe voltage that must be applied to
  produce an avalanche in GM tube = %6.2f volt", V)
  ;
10
11 // Result
12 // The voltage that must be applied to produce an
  avalanche in GM tube = 728.52 volt

```

---

**Scilab code Exa 17.18** Maximum permissible voltage fluctuation in a GM counter

```

1 // Scilab code Ex17.18 : Pg:898 (2011)
2 clc;clear;
3 count_err = 1e-03;    // Fractional error in
  counting
4 m = 3;    // Plateau slope
5 delta_V = count_err*100/m*100;    // Maximum
  permissible voltage fluctuation in a GM counter,
  volt
6 printf("\nThe maximum permissible voltage
  fluctuation in a GM counter = %3.1f volts",
  delta_V);
7

```

```
8 // Result
9 // The maximum permissible voltage fluctuation in a
  GM counter = 3.3 volts
```

---

# Chapter 19

## Superconductivity

Scilab code Exa 19.1 Critical field for lead at 4 K

```
1 // Scilab Code Ex19.1: Page-959 (2011)
2 clc; clear;
3 T_c = 6.2; // Critical temperature of lead in
  superconducting state, K
4 T = 4; // Temperature at which critical field
  of lead is to be found out, K
5 H_c0 = 0.064; // Critical field for lead at 0 K,
  MA/m
6 H_cT = H_c0*(1-(T/T_c)^2); // Critical field for
  lead at 4 K, MA/m
7 printf("\\nThe critical field for lead at 4 K = %5.3f
  MA/m", H_cT);
8 // Result
9 // The critical field for lead at 4 K = 0.037 MA/m
```

---

Scilab code Exa 19.2 Isotopic effect in mercury

```
1 // Scilab Code Ex19.2: Page-959 (2011)
```

```

2 clc; clear;
3 T_c1 = 4.153;    // Critical temperature of mercury
   for its one isotope, K
4 M1 = 200.59;    // Mass of first isotope of mercury,
   amu
5 M2 = 204;      // Mass of second isotope of mercury
   , amu
6 // From isotopic effect of superconductivity,
7 // T_c2/T_c1 = sqrt(M1/M2), solving for T_c2
8 T_c2 = T_c1*sqrt(M1/M2);    // Critical temperature
   of mercury for second isotope, K
9 printf("\\nThe critical temperature of mercury for
   its isotope of mass 204 amu = %5.3f K", T_c2);
10
11 // Result
12 // The critical temperature of mercury for its
   isotope of mass 204 amu = 4.118 K

```

---

**Scilab code Exa 19.3** Critical current through superconducting aluminium wire

```

1 // Scilab Code Ex19.3: Page-960 (2011)
2 clc; clear;
3 d = 1e-003;    // Diameter of aluminium wire, m
4 r = d/2;      // Radius of aluminium wire, m
5 H_c = 7.9e+003; // Critical magnetic field for Al
   , A/m
6 I_c = 2*3.14*r*H_c; // Critical current through
   superconducting aluminium wire, A
7 printf("\\nThe critical current through
   superconducting aluminium wire = %6.3f A", I_c);
8
9 // Result
10 // The critical current through superconducting
   aluminium wire = 24.806 A

```

---

**Scilab code Exa 19.4** Critical current density for superconducting wire of lead at different temperatures

```
1 // Scilab Code Ex19.4: Page-960 (2011)
2 clc; clear;
3 T_c = 7.18; // Critical temperature of lead in
  superconducting state, K
4 H_c0 = 6.5e+004; // Critical field for lead at 0
  K, A/m
5 // At T = 4.2 K
6 T = 4.2; // Temperature at which critical
  field of lead is to be found out, K
7 H_cT = H_c0*(1-(T/T_c)^2); // Critical field for
  lead at 4 K, A/m
8 d = 1e-003; // Diameter of lead wire, m
9 r = d/2; // Radius of lead wire, m
10 I_c = 2*3.14*r*H_cT; // Critical current through
  superconducting lead wire, A
11 J_c = I_c/(3.14*r^2); // Critical current density
  for superconducting lead wire, A/Sq. meter
12 printf("\nThe critical current density at %3.1f K =
  %5.3e A/Sq.m", T, J_c);
13 // At T = 7 K
14 T = 7; // Temperature at which critical field
  of lead is to be found out, K
15 H_cT = H_c0*(1-(T/T_c)^2); // Critical field for
  lead at 4 K, A/m
16 d = 1e-003; // Diameter of lead wire, m
17 r = d/2; // Radius of lead wire, m
18 I_c = 2*3.14*r*H_cT; // Critical current through
  superconducting lead wire, A
19 J_c = I_c/(3.14*r^2); // Critical current density
  for superconducting lead wire, A/Sq. meter
20 printf("\nThe critical current density at %3.1f K =
```

```

    %4.2 e A/Sq.m", T, J_c);
21
22 // Result
23 // The critical current density at 4.2 K = 1.710e
    +008 A/Sq.m
24 // The critical current density at 7.0 K = 1.29e+007
    A/Sq.m

```

---

**Scilab code Exa 19.5** Critical temperature of lead from London penetration depth

```

1 // Scilab Code Ex19.5:Page-961 (2011)
2 clc; clear;
3 T1 = 3; // Initial temperature of lead wire, K
4 T2 = 7.1; // Final temperature of lead wire, K
5 lambda1 = 39.6; // Initial London penetration
    depth for lead, mm
6 lambda2 = 173; // Final London penetration depth
    for lead, mm
7 // As  $\lambda_T = \lambda_0 * (1 - (T/T_c)^4)^{-1/2}$  so
8 //  $(\lambda_1/\lambda_2)^2 = (T_c^4 - T_2^4)/(T_c^4 - T_1^4)$ 
9 // Solving for T_c
10 T_c = ((T2^4 - T1^4 * (lambda1/lambda2)^2) / (1 - (lambda1/
    lambda2)^2))^(1/4);
11 printf("\\nThe critical temperature of lead = %5.3f K
    ", T_c);
12
13 // Result
14 // The critical temperature of lead = 7.193 K

```

---

**Scilab code Exa 19.6** Field strength required by lead to lose its superconducting state at 0 K



```

1 // Scilab Code Ex19.6: Page-962 (2011)
2 clc; clear;
3 T_c = 7.2; // Critical temperature of lead in
    superconducting state, K
4 T = 5; // Temperature at which lead loses its
    superconducting state, K
5 H_cT = 3.3e+004; // Critical magnetic field for
    superconducting lead at 5 K, A/m
6 // As  $H_{cT} = H_{c0} * (1 - (T/T_c)^2)$ , solving for  $H_{c0}$ 
7 H_c0 = H_cT / (1 - (T/T_c)^2); // Critical field for
    lead at 0 K, A/m
8 printf("\nThe critical magnetic field for lead at 0
    K = %4.2e A/m", H_c0);
9
10 // Result
11 // The critical magnetic field for lead at 0 K =
    6.37e+004 A/m

```

---

### Scilab code Exa 19.7 Critical temperature for niobium

```

1 // Scilab Code Ex19.7: Page-962 (2011)
2 clc; clear;
3 H_c0 = 2e+005; // Critical field for niobium at 0
    K, A/m
4 H_cT = 1e+005; // Critical magnetic field for
    superconducting niobium at 5 K, A/m
5 T = 8; // Temperature at which lead loses its
    superconducting state, K
6 // As  $H_{cT} = H_{c0} * (1 - (T/T_c)^2)$ , solving for  $T_c$ 
7 T_c = T / (1 - H_cT/H_c0)^(1/2);
8 printf("\nThe critical temperature for niobium = %6
    .3f K", T_c);
9
10 // Result
11 // The critical temperature for niobium = 11.314 K

```

