

Scilab Textbook Companion for  
Engineering Physics  
by V. Yadav<sup>1</sup>

Created by  
Ujval Gupta  
Modern Physics  
Physics  
Shri Mata Vaishno Devi University  
College Teacher  
Dr. Yugal Khajuria  
Cross-Checked by  
Lavitha Pereira

April 11, 2014

<sup>1</sup>Funded by a grant from the National Mission on Education through ICT,  
<http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab  
codes written in it can be downloaded from the "Textbook Companion Project"  
section at the website <http://scilab.in>

# **Book Description**

**Title:** Engineering Physics

**Author:** V. Yadav

**Publisher:** Tata McGraw-Hill

**Edition:** 1

**Year:** 2009

**ISBN:** 978-0-07-008569-5

Scilab numbering policy used in this document and the relation to the above book.

**Exa** Example (Solved example)

**Eqn** Equation (Particular equation of the above book)

**AP** Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

# Contents

<b>List of Scilab Codes</b>	<b>4</b>
<b>1 Quantum Mechanics</b>	<b>12</b>
<b>2 Interference</b>	<b>41</b>
<b>3 Diffraction</b>	<b>90</b>
<b>4 Polarization</b>	<b>125</b>
<b>5 Nuclear Physics</b>	<b>141</b>
<b>6 Semiconductors and Nano Physics</b>	<b>146</b>
<b>7 Fiber Optics</b>	<b>148</b>
<b>8 Laser</b>	<b>156</b>

# List of Scilab Codes

Exa 1.1	Energy of the particle from de Broglie wavelength . . . . .	12
Exa 1.2	de Broglie wavelength of the particle . . . . .	12
Exa 1.3	de Broglie wavelength of an accelerated electron . . . . .	13
Exa 1.4	Energy of the electron from de Broglie wavelength . . . . .	13
Exa 1.5	Velocity and de Broglie wavelength of a neutron . . . . .	14
Exa 1.6	Wavelength of thermal neutron at room temperature . . . . .	15
Exa 1.7	Angle of deviation for first order diffraction maxima . . . . .	15
Exa 1.8	de Broglie wavelength of a moving electron . . . . .	16
Exa 1.9	de Broglie wavelength of a neutron of given kinetic energy . . . . .	17
Exa 1.10	de Broglie wavelength associated with moving proton . . . . .	17
Exa 1.11	Wavelength of matter wave associated with moving proton . . . . .	18
Exa 1.12	de Broglie wavelength of an electron accelerated through a given potential . . . . .	18
Exa 1.13	Interplanar spacing of the crystal . . . . .	19
Exa 1.14	Interplanar spacing using Bragg law . . . . .	20
Exa 1.15	de Broglie wavelength of electron accelerated at V volts . . . . .	20
Exa 1.16	de Broglie wavelength of electron accelerated from rest . . . . .	21
Exa 1.17	The wavelength associated with moving mass . . . . .	22
Exa 1.18	Wavelength of an electron from its kinetic energy . . . . .	22
Exa 1.19	Kinetic energy of electron . . . . .	22
Exa 1.20	Speed of proton for an equivalent wavelength of that of electron . . . . .	23
Exa 1.21	de Broglie wavelength of the electron . . . . .	24
Exa 1.23	Minimum accuracy to locate the position of an electron . . . . .	24
Exa 1.24	Uncertainty in energy of an emitted photon . . . . .	25
Exa 1.25	Minimum uncertainty in velocity of electron . . . . .	26

Exa 1.26	Minimum uncertainty in momentum and minimum kinetic energy of proton . . . . .	26
Exa 1.27	Uncertainty in momentum of electron . . . . .	27
Exa 1.28	Uncertainty in position of the particle . . . . .	28
Exa 1.29	Uncertainty in position of the moving electron . . . . .	28
Exa 1.30	Smallest possible uncertainty in position of the electron . . . . .	29
Exa 1.31	Difference in the energy between the neighboring levels of Na at the highest state . . . . .	30
Exa 1.32	Energy of the neutron confined in a nucleus . . . . .	31
Exa 1.33	Energy of an electron moving in one dimensional infinitely high potential box . . . . .	31
Exa 1.34	Lowest energy of an electron in a one dimensional force free region . . . . .	32
Exa 1.35	First three energy levels of an electron in one dimensional box . . . . .	33
Exa 1.36	Lowest two permitted energy values of the electron in a 1D box . . . . .	34
Exa 1.37	Lowest energy of the neutron confined to the nucleus .	35
Exa 1.38	X ray scattering . . . . .	36
Exa 1.39	Wavelength of scattered Xray . . . . .	36
Exa 1.40	Wavelength of scattered radiation with changed angle of view . . . . .	37
Exa 1.41	Compton scattering through aluminium foil . . . . .	38
Exa 1.42	Energetic electrons in the Xray tube . . . . .	39
Exa 2.1	Slit separation in Double Slit experiment . . . . .	41
Exa 2.2	Wavelength of light in Young Double Slit experiment .	41
Exa 2.3	Ratio of maximum intensity to minimum intensity of interference fringes . . . . .	42
Exa 2.4	Wavelength of light from monochromatic coherent sources	43
Exa 2.5	Separation between fourth order dark fringes . . . . .	43
Exa 2.6	Distance between two coherent sources . . . . .	44
Exa 2.7	Fringe width of the interference pattern due to biprism	45
Exa 2.8	Angle of vertex of the biprism . . . . .	45
Exa 2.9	Number of interference fringes for changed wavelength	46
Exa 2.10	Wavelength of light in a biprism experiment . . . . .	47
Exa 2.11	Fringe width at a certain distance from biprism . . . . .	47
Exa 2.12	Distance between coherent sources in biprism experiment	48
Exa 2.13	Effect of slit separation on fringe width . . . . .	49

Exa 2.14	Effect of slit biprism separation on fringe width . . . . .	49
Exa 2.15	Distance between interference bands . . . . .	50
Exa 2.16	Angle of vertex of Fresnel biprism . . . . .	51
Exa 2.17	Wavelength of light used in biprism experiment to illuminate slits . . . . .	52
Exa 2.18	Vertex angle of Fresnel biprism . . . . .	52
Exa 2.19	Order of visible fringe for changed wavelength of light	53
Exa 2.20	Angle of vertex of biprism . . . . .	54
Exa 2.21	Separation between two coherent sources . . . . .	55
Exa 2.22	Refractive index of the glass sheet . . . . .	55
Exa 2.23	Refractive index of material of sheet . . . . .	56
Exa 2.24	Wavelength of light used in biprism arrangement . . . . .	56
Exa 2.25	Thickness of the transparent sheet . . . . .	57
Exa 2.26	Thickness of the transparent sheet from fringe shift . .	57
Exa 2.27	Refractive index of thin mica sheet . . . . .	58
Exa 2.28	Wavelength of light used in double slit experiment . .	59
Exa 2.29	Thickness of mica sheet from central fringe shift . . .	59
Exa 2.30	Refractive index of material from shifting fringe pattern	60
Exa 2.31	Fringe width and optical path change during interference of waves through mica and glass . . . . .	60
Exa 2.32	Thickness of mica sheet from Fresnel biprism experiment	61
Exa 2.33	Smallest thickness of glass plate for a fringe of minimum intensity . . . . .	62
Exa 2.34	The wavelength reflected strongly from the soap film .	63
Exa 2.35	Order of interference of the dark band . . . . .	63
Exa 2.36	Absent wavelength of reflected light in the visible spectrum . . . . .	64
Exa 2.37	Minimum thickness of the plate that will appear dark in the reflection pattern . . . . .	65
Exa 2.38	Thickness of the thin soap film . . . . .	66
Exa 2.39	Order of interference for which light is strongly reflected	67
Exa 2.40	Minimum thickness of the film for which light is strongly reflected . . . . .	68
Exa 2.41	Thickness of the soap film for dark fringe in reflected pattern . . . . .	68
Exa 2.42	Wavelength in the visible range which is intensified in the reflected beam . . . . .	69
Exa 2.43	Thickness of the film with incident white light . . . . .	70

Exa 2.44	Thickness of the film with parallel beam of yellow light	71
Exa 2.46	Refractive index of oil . . . . .	71
Exa 2.47	Thickness of the soap film to produce constructive interference during reflection . . . . .	72
Exa 2.48	Wavelength of light falling on wedge shaped film . . . . .	73
Exa 2.49	Difference between the thicknesses of the films . . . . .	73
Exa 2.50	Angle of thin wedge shaped film . . . . .	74
Exa 2.51	Wavelength of light used to illuminate a wedge shaped film . . . . .	74
Exa 2.52	Thickness of the wire separating two glass surfaces . .	75
Exa 2.53	Angle of the wedge shaped air film . . . . .	76
Exa 2.54	Acute angle of the wedge shaped film . . . . .	76
Exa 2.55	Diameter of nth dark ring due to first wavelength . .	77
Exa 2.56	Diameter of fifteenth dark ring . . . . .	78
Exa 2.57	Order of a dark ring having thrice the diameter of the thirtieth ring . . . . .	78
Exa 2.58	Radius of curvature of lens and thickness of air film . .	79
Exa 2.59	Refractive index of the liquid . . . . .	80
Exa 2.60	Wavelength of light used in Newton rings experiment .	80
Exa 2.61	Diameter of fifteenth bright ring . . . . .	81
Exa 2.62	Wavelength of light used in Newton rings experiment .	82
Exa 2.63	Diameter of fifteenth dark Newton ring . . . . .	82
Exa 2.64	Diameter of fifteenth dark ring due to first wavelength	83
Exa 2.65	Refractive index of the liquid filled into container . .	84
Exa 2.67	Refractive index of the liquid . . . . .	85
Exa 2.68	Diameter of eighteenth dark ring . . . . .	85
Exa 2.69	Wavelength of light used to illuminate plano convex lens in Newton rings experiment . . . . .	86
Exa 2.70	Wavelength of monochromatic light used in Michelson Interferometer . . . . .	86
Exa 2.71	Number of fringes that passes across the cross wire of telescope . . . . .	87
Exa 2.72	Distance between two successive positions of movable mirror . . . . .	88
Exa 2.73	Thickness of the transparent glass film . . . . .	88
Exa 3.1	Position of the screen so that light is focused on the brightest spot . . . . .	90
Exa 3.2	Zone plate with a point source of light on the axis . .	90

Exa 3.3	Position of the first image in a zone plate . . . . .	91
Exa 3.4	Principal focal length of zone plate . . . . .	92
Exa 3.5	Half angular width at central maximum in Fraunhoffer diffraction . . . . .	93
Exa 3.6	Width of the slit . . . . .	93
Exa 3.7	Angular width of central maximum . . . . .	94
Exa 3.8	Distance between first minima and the next minima from the axis . . . . .	95
Exa 3.9	Width of central maxima in diffraction pattern . . . . .	96
Exa 3.10	Slit width in Fraunhoffer single slit experiment . . . . .	96
Exa 3.11	Half angular width of central maxima . . . . .	97
Exa 3.12	Wavelength of light used in Fraunhoffer diffraction due to single slit . . . . .	97
Exa 3.13	Width of central maxima from position of first secondary minima . . . . .	98
Exa 3.14	Wavelength of monochromatic light used in illuminating a slit . . . . .	99
Exa 3.15	Distance between second dark and next bright fringe on the axes . . . . .	99
Exa 3.16	Width of the slit from first order diffraction . . . . .	100
Exa 3.18	Fraunhoffer diffraction due to double slits . . . . .	101
Exa 3.19	Fringe separation in Fraunhoffer double slit diffraction pattern . . . . .	101
Exa 3.20	Positions of first secondary maxima and minima in double slit diffraction . . . . .	102
Exa 3.21	Missing orders of spectra in Fraunhoffer double slit diffraction . . . . .	103
Exa 3.22	Angles of diffraction for the principal maxima for two lines of sodium . . . . .	104
Exa 3.23	Highest order spectrum which can be seen in monochromatic light . . . . .	105
Exa 3.24	Angle of separation in second order of diffraction spectrum . . . . .	105
Exa 3.25	Separation of two lines in first order spectrum . . . . .	106
Exa 3.26	Difference in the deviation in the first and third order spectra . . . . .	107
Exa 3.27	Order of diffraction for the given grating element and wavelength of light . . . . .	108

Exa 3.28	Number of lines ruled on the grating surface . . . . .	109
Exa 3.29	Angles at which first and second order maxima are observed . . . . .	110
Exa 3.30	Least width of plane transmission grating . . . . .	110
Exa 3.31	Minimum grating width required to resolve two wavelengths . . . . .	111
Exa 3.32	Angle of diffraction for maxima in first order . . . . .	113
Exa 3.33	Wavelength of light used in obtaining second order diffraction maximum . . . . .	113
Exa 3.34	Number of visible orders using diffraction grating . . .	114
Exa 3.35	Distance between two wavelengths seen as separate . .	115
Exa 3.36	Number of lines per cm on grating surface . . . . .	115
Exa 3.37	Total number of lines on grating surface . . . . .	116
Exa 3.38	Angular separation between the sodium D1 and D2 lines	117
Exa 3.39	Minimum number of lines in a grating . . . . .	118
Exa 3.40	Linear separation of two points on the moon . . . . .	119
Exa 3.41	Minimum required number of lines on the plane transmission grating . . . . .	119
Exa 3.42	Number of lines on the plane transmission grating to just resolve the sodium lines . . . . .	120
Exa 3.43	Minimum width of the grating to resolve the sodium lines in third order . . . . .	121
Exa 3.44	Dispersive power and diffraction angle for grating . .	121
Exa 3.45	Distance between centres of images of the two stars .	123
Exa 3.46	Aperture of the objective of the microscope . . . . .	123
Exa 4.1	Refractive index of the material . . . . .	125
Exa 4.2	Polarization by reflection . . . . .	125
Exa 4.3	Angle of refraction of the ray . . . . .	126
Exa 4.4	Angle of minimum deviation for green light . . . . .	126
Exa 4.5	Polarizing angles of the materials for given refractive indices . . . . .	127
Exa 4.6	Angle of rotation of analyser . . . . .	128
Exa 4.7	Angles of rotation of analyser for given transmitted light intensities . . . . .	128
Exa 4.8	Angle of minimum deviation for green light . . . . .	129
Exa 4.9	Ratio of ordinary to extraordinary ray intensities . . .	130
Exa 4.10	Thickness of quarter wave plate . . . . .	130

Exa 4.11	Least thickness of plate for which emergent beam is plane polarised . . . . .	131
Exa 4.12	Difference in refractive indices of rays . . . . .	131
Exa 4.13	The thickness of a half wave plate . . . . .	132
Exa 4.14	The thickness of a quarter wave plate . . . . .	132
Exa 4.15	The thickness of a half wave plate quartz . . . . .	133
Exa 4.16	Difference between refractive indices . . . . .	134
Exa 4.17	Specific rotation of superposition . . . . .	134
Exa 4.18	Strength of sugar solution . . . . .	135
Exa 4.19	Quantity of sugar contained in the tube in the form of solution . . . . .	135
Exa 4.20	Specific rotation of sugar solution from the given data . . . . .	136
Exa 4.21	Angle of rotation of the plane of polarization . . . . .	137
Exa 4.22	Angle of rotation of the optically active solution . . . . .	137
Exa 4.23	Angle of rotation in a tube of new length . . . . .	138
Exa 4.24	Mass of sugar in the solution contained in the tube . . . . .	138
Exa 4.25	Percentage purity of the sugar sample . . . . .	139
Exa 4.26	Angle of rotation produced by the polarimeter plate . . . . .	140
Exa 5.1	Mass defect of He . . . . .	141
Exa 5.3	Maximum energy of proton in a cyclotron . . . . .	141
Exa 5.4	Energy of an electron in a betatron . . . . .	142
Exa 5.5	Final energy gained by electrons in a betatron . . . . .	142
Exa 5.6	Energy produced in fission of U235 . . . . .	143
Exa 5.7	Power output of nuclear reactor . . . . .	144
Exa 5.8	Average current in the GM counter circuit . . . . .	144
Exa 5.9	Energy needed to remove a neutron from Ca nucleus . . . . .	145
Exa 6.1	Resistivity of intrinsic semiconductor at 300 K . . . . .	146
Exa 6.2	Velocity of electron at Fermi level . . . . .	147
Exa 7.1	Critical angle and acceptance angle in an optical fibre . . . . .	148
Exa 7.2	Critical angle acceptance angle and numerical aperture in an optical fibre . . . . .	149
Exa 7.3	Parameters of an optical fibre using relative refractive index difference . . . . .	149
Exa 7.4	Refractive index of cladding . . . . .	150
Exa 7.5	Numerical aperture for an optical fibre . . . . .	151
Exa 7.6	Refractive index of the cladding . . . . .	151
Exa 7.7	Comparison of the acceptance angle for meridional rays with that for the skew rays . . . . .	152

Exa 7.8	Normalized frequency for V number for the fibre . . . . .	153
Exa 7.9	Number of modes in the step index fibre . . . . .	153
Exa 7.10	Radius of core for single mode operation in step index fibre . . . . .	154
Exa 7.11	Signal attenuation in optical fibre . . . . .	155
Exa 8.1	Difference between upper and lower energy levels for the most prominent wavelength . . . . .	156
Exa 8.2	Frequency and wavelength of carbon dioxide laser . . . . .	156
Exa 8.3	Energy of one emitted photon and total energy available per laser pulse . . . . .	157
Exa 8.4	Relative population of levels in Ruby laser . . . . .	158
Exa 8.5	Population of two states in He Ne laser . . . . .	158

# Chapter 1

## Quantum Mechanics

**Scilab code Exa 1.1** Energy of the particle from de Broglie wavelength

```
1 // Scilab Code Ex1.1: Page -1.5 (2009)
2 clc; clear;
3 lambda = 2.1e-010; // de Broglie wavelength of the
4 m = 1.67e-027; // Mass of the particle , kg
5 h = 6.626e-034; // Planck's constant , Js
6 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
7 // From de Broglie relation , lambda = h/sqrt(2*m*E) ,
8 // solving for E
9 E = h^2/(2*m*lambda^2*e); // Energy of the
10 // particle , eV
11 printf("\nThe energy of the particle from de Broglie
12 // wavelength = %5.3e eV", E);
13
14 // Result
15 // The energy of the particle from de Broglie
16 // wavelength = 1.863e-002 eV
```

---

**Scilab code Exa 1.2** de Broglie wavelength of the particle

```

1 // Scilab Code Ex1.2: Page-1.5 (2009)
2 clc; clear;
3 m = 1.67e-027;           // Mass of the particle , kg
4 h = 6.626e-034;          // Planck's constant , Js
5 e = 1.6e-019;            // Energy equivalent of 1 eV, J/eV
6 E = 1e+011*e;            // Energy of the particle , J
7 lambda = h/sqrt(2*m*E);   // de Broglie wavelength
                                of the particle , m
8 printf("\nThe de Broglie wavelength of the particle
         = %4.2e m", lambda);
9
10 // Result
11 // The de Broglie wavelength of the particle = 9.06e
         -017 m

```

---

**Scilab code Exa 1.3** de Broglie wavelength of an accelerated electron

```

1 // Scilab Code Ex1.3: Page-1.5 (2009)
2 clc; clear;
3 V = 20e+03;              // Accelerating voltage of electron ,
                           V
4 lambda = 12.25/sqrt(V);    // de Broglie wavelength
                                of the accelerated electron , m
5 printf("\nThe de Broglie wavelength of the electron
         = %6.4f angstrom", lambda);
6
7 // Result
8 // The de Broglie wavelength of the electron =
         0.0866 angstrom

```

---

**Scilab code Exa 1.4** Energy of the electron from de Broglie wavelength

```
1 // Scilab Code Ex1.4: Page-1.6 (2009)
```

```

2 clc; clear;
3 lambda = 5.2e-03; // de Broglie wavelength of the
        electron , m
4 m = 9.1e-031;      // Mass of the electron , kg
5 h = 6.626e-034;    // Planck's constant , Js
6 e = 1.6e-019;     // Energy equivalent of 1 eV, J/eV
7 // From de Broglie relation , lambda = h/sqrt(2*m*E) ,
        solving for E
8 E = h^2/(2*m*lambda^2*e); // Energy of the
        electron , eV
9 printf("\nThe energy of the electron from de Broglie
        wavelength = %5.3e eV", E);
10
11 // Result
12 // The energy of the electron from de Broglie
        wavelength = 5.576e-014 eV

```

---

### Scilab code Exa 1.5 Velocity and de Broglie wavelength of a neutron

```

1 // Scilab Code Ex1.5: Page-1.6 (2009)
2 clc; clear;
3 m = 1.67e-027;      // Mass of the neutron , kg
4 h = 6.626e-034;    // Planck's constant , Js
5 e = 1.6e-019;     // Energy equivalent of 1 eV, J/eV
6 E = 1e+04*e;      // Energy of the neutron , J
7 // As E = 1/2*m*v^2, solving for v
8 v = sqrt(2*E/m);   // Velocity of the neutron , m/s
9 lambda = h/(m*v); // de Broglie wavelength of the
        neutron , m
10 printf("\nThe velocity of the neutron = %4.2e m/s",
        v);
11 printf("\nThe de Broglie wavelength of the neutron =
        %4.2e m", lambda);
12
13 // Result

```

```
14 // The velocity of the neutron = 1.38e+006 m/s  
15 // The de Broglie wavelength of the neutron = 2.87e  
    -013 m
```

---

**Scilab code Exa 1.6** Wavelength of thermal neutron at room temperature

```
1 // Scilab Code Ex1.6: Page-1.6 (2009)  
2 clc; clear;  
3 m = 1.67e-027;           // Mass of the neutron , kg  
4 k = 1.38e-023;           // Boltzmann constant , J/mol/K  
5 T = 27+273;              // Room temperature , K  
6 h = 6.626e-034;          // Planck's constant , Js  
7 e = 1.6e-019;             // Energy equivalent of 1 eV, J/eV  
8 v = sqrt(3*k*T/m);       // Velocity of the neutron ,  
    m/s  
9 lambda = h/(m*v);        // de Broglie wavelength of the  
    neutron , m  
10 printf("\nThe de Broglie wavelength of the thermal  
    neutrons = %4.2f angstrom", lambda/1e-010);  
11  
12 // Result  
13 // The de Broglie wavelength of the thermal neutrons  
    = 1.45 angstrom
```

---

**Scilab code Exa 1.7** Angle of deviation for first order diffraction maxima

```
1 // Scilab Code Ex1.7: Page-1.6 (2009)  
2 clc; clear;  
3 m = 9.1e-031;             // Mass of the electron , kg  
4 e = 1.6e-019;              // Energy equivalent of 1 eV, J/eV  
5 h = 6.626e-034;            // Planck's constant , Js  
6 E = 20e+03*e;              // Energy of the electron , J  
7 // As 1/2*m*v^2 = E, solving for v
```

```

8 v = sqrt(2*E/m);           // Velocity of the electron ,
m/s
9 lambda = h/(m*v); // de Broglie wavelength of the
electron , m
10 n = 1; // First order diffraction
11 d = 9.8e-011; // Atomic spacing for thin gold
foil , m
12 // Using Bragg's equation , 2*d*sin(theta) = n*lambda
and solving for theta
13 theta = asind(n*lambda/(2*d)); // Angle of
deviation for first order diffraction maxima,
degree
14 printf("\nThe angle of deviation for first order
diffraction maxima = %4.2f degrees", theta);
15
16 // Result
17 // The angle of deviation for first order
diffraction maxima = 2.54 degrees

```

---

**Scilab code Exa 1.8** de Broglie wavelength of a moving electron

```

1 // Scilab Code Ex1.8: Page-1.7 (2009)
2 clc; clear;
3 m = 9.1e-031; // Mass of the electron , kg
4 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
5 h = 6.626e-034; // Planck's constant , Js
6 E = 5*e; // Energy of the electron , J
7 // As 1/2*m*v^2 = E, solving for v
8 v = sqrt(2*E/m); // Velocity of the electron ,
m/s
9 lambda = h/(m*v); // de Broglie wavelength of the
electron , m
10 printf("\nThe de Broglie wavelength of the electron
= %3.1f angstrom", lambda/1e-010);
11

```

```
12 // Result
13 // The de Broglie wavelength of the electron = 5.5
angstrom
```

---

**Scilab code Exa 1.9** de Broglie wavelength of a neutron of given kinetic energy

```
1 // Scilab Code Ex1.9: Page-1.7 (2009)
2 clc; clear;
3 m = 1.67e-027;           // Mass of the neutron , kg
4 e = 1.6e-019;            // Energy equivalent of 1 eV, J/eV
5 h = 6.626e-034;          // Planck's constant , Js
6 E = 1*e;                // Energy of the electron , J
7 lambda = h/sqrt(2*m*E); // de Broglie wavelength of
                           the neutron , m
8 printf("\nThe de Broglie wavelength of the neutron =
        %4.2f angstrom", lambda/1e-010);
9
10 // Result
11 // The de Broglie wavelength of the neutron = 0.29
angstrom
```

---

**Scilab code Exa 1.10** de Broglie wavelength associated with moving proton

```
1 // Scilab Code Ex1.10: Page-1.8 (2009)
2 clc; clear;
3 m = 1.67e-027;           // Mass of the proton , kg
4 c = 3e+08;                // Speed of light , m/s
5 v = 1/20*c;               // Velocity of the proton , m/s
6 h = 6.626e-034;          // Planck's constant , Js
7 lambda = h/(m*v);        // de Broglie wavelength of the
                           neutron , m
```

```

8 printf("\nThe de Broglie wavelength associated with
      moving proton = %5.3e m", lambda);
9
10 // Result
11 // The de Broglie wavelength associated with moving
      proton = 2.645e-14 m

```

---

**Scilab code Exa 1.11** Wavelength of matter wave associated with moving proton

```

1 // Scilab Code Ex1.11: Page-1.8 (2009)
2 clc; clear;
3 m = 1.67e-027;           // Mass of the proton , kg
4 v = 2e+08;                // Velocity of the proton , m/s
5 h = 6.626e-034;          // Planck's constant , Js
6 lambda = h/(m*v);        // de Broglie wavelength of the
      neutron , m
7 printf("\nThe wavelength of matter wave associated
      with moving proton = %5.3e m", lambda);
8
9 // Result
10 // The wavelength of matter wave associated with
      moving proton = 1.984e-15 m

```

---

**Scilab code Exa 1.12** de Broglie wavelength of an electron accelerated through a given potential

```

1 // Scilab Code Ex1.12: Page-1.17 (2009)
2 clc; clear;
3 m = 9.1e-031;             // Mass of the electron , kg
4 q = 1.6e-019;              // Charge on an electron , C
5 V = 50;                   // Accelerating potential , V
6 E = q*V;                  // Energy gained by the electron , J

```

```

7 h = 6.626e-034;      // Planck's constant , Js
8 lambda = h/sqrt(2*m*E); // de Broglie wavelength of
                           the electron , m
9 printf("\nThe de Broglie wavelength of the electron
        accelearted through a given potential = %5.3e m" ,
        lambda);
10
11 // Result
12 // The de Broglie wavelength of the electron
        accelearted through a given potential = 1.736e-10
        m

```

---

### Scilab code Exa 1.13 Interplanar spacing of the crystal

```

1 // Scilab Code Ex1.13: Page-1.17 (2009)
2 clc; clear;
3 theta = 45;      // Diffraction angle , degrees
4 h = 6.626e-034; // Planck's constant
5 m = 1.67e-027; // Mass of a neutron , kg
6 n = 1;          // Order of diffraction
7 k = 1.38e-023; // Boltzmann constant , J/mol/K
8 T = 27+273;    // Absolute room temperature , K
9 E = 3/2*k*T;   // Energy of the neutron , J
10 lambda = h/sqrt(2*m*E); // de-Broglie wavelength of
                           neutrons , m
11 // From Bragg's law , 2*d*sin(theta) = n*lambda ,
                           solving for d
12 d = n*lambda/(2*sind(theta));
13 printf("\nThe interplanar spacing of the crystal =
        %4.2f angstrom", d/1e-010);
14
15 // Result
16 // The interplanar spacing of the crystal = 1.03
        angstrom

```

---

**Scilab code Exa 1.14** Interplanar spacing using Bragg law

```
1 // Scilab Code Ex1.14: Page-1.18 (2009)
2 clc; clear;
3 theta = 70;      // Glancing angle at which
                   reflection occurs , degrees
4 h = 6.626e-034;    // Planck's constant
5 m = 9.1e-031;      // Mass of a electron , kg
6 e = 1.6e-019;      // Electronic charge , C
7 V = 1000;          // Accelerating potential , V
8 n = 1;             // Order of diffraction
9 E = e*V;           // Energy of the electron , J
10 lambda = h/sqrt(2*m*E); // de-Broglie wavelength of
                           electron , m
11 // From Bragg's law , 2*d*sin(theta) = n*lambda ,
   solving for d
12 d = n*lambda/(2*sind(theta)); // Interplanar
                           spacing , m
13 printf("\nThe interplanar spacing of the crystal =
       %6.4e m", d);
14
15 // Result
16 // The interplanar spacing of the crystal = 2.0660e
   -11 m
```

---

**Scilab code Exa 1.15** de Broglie wavelength of electron accelerated at V volts

```
1 // Scilab Code Ex1.15: Page-1.18 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant
4 m = 9.1e-031;      // Mass of a electron , kg
```

```

5 e = 1.6e-019;           // Electronic charge , C
6 V = 1;      // For simplicity the accelerating
    potential is assumed to be unity , V
7 E = e*V;       // Energy of the electron , J
8 lambda = h/sqrt(2*m*E); // de-Broglie wavelength of
    electron , m
9 printf("\nde-Broglie wavelength of electron
    accelerated at V volts = %5.2f/sqrt(V) angstrom",
    lambda/1e-010);
10
11 // Result
12 // de-Broglie wavelength of electron accelerated at
    V volts = 12.23/sqrt(V) angstrom

```

---

**Scilab code Exa 1.16** de Broglie wavelength of electron accelerated from rest

```

1 // Scilab Code Ex1.16: Page -1.18 (2009)
2 clc; clear;
3 h = 6.6e-034;           // Planck 's constant
4 m = 9.1e-031;          // Mass of a electron , kg
5 e = 1.6e-019;          // Electronic charge , C
6 V = 100;      // Accelerating potential for electron , V
7 E = e*V;       // Energy of the electron , J
8 lambda = h/sqrt(2*m*E); // de-Broglie wavelength of
    electron , m
9 printf("\nde-Broglie wavelength of electron
    accelerated at %d volts = %6.4e m", V, lambda);
10
11 // Result
12 // de-Broglie wavelength of electron accelerated at
    100 volts = 1.2231e-10 m

```

---

**Scilab code Exa 1.17** The wavelength associated with moving mass

```
1 // Scilab Code Ex1.17: Page-1.19 (2009)
2 clc; clear;
3 m = 10e-03;      // Mass of the body , kg
4 v = 110;         // Velocity of the mass , m/s
5 h = 6.6e-034;    // Planck's constant
6 lambda = h/(m*v); // de-Broglie wavelength of
                     electron , m
7 printf("\nThe wavelength associated with mass moving
        with velocity %d m/s = %1.0e m", v, lambda);
8
9 // Result
10 // The wavelength associated with mass moving with
    velocity 110 m/s = 6e-34 m
```

---

**Scilab code Exa 1.18** Wavelength of an electron from its kinetic energy

```
1 // Scilab Code Ex1.18: Page-1.19 (2009)
2 clc; clear;
3 m = 9.1e-031;     // Mass of the electron , kg
4 Ek = 1.27e-017;    // Kinetic energy of electron , J
5 h = 6.6e-034;      // Planck's constant
6 lambda = h/sqrt(2*m*Ek); // de-Broglie wavelength of
                           electron , m
7 printf("\nThe wavelength associated with moving
        electron = %4.2f angstrom", lambda/1e-010);
8
9 // Result
10 // The wavelength associated with moving electron =
    1.37 angstrom
```

---

**Scilab code Exa 1.19** Kinetic energy of electron

```

1 // Scilab Code Ex1.19: Page -1.19 (2009)
2 clc; clear;
3 m = 9.1e-031; // Mass of the electron , kg
4 h = 6.6e-034; // Planck 's constant
5 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
6 lambda = 9.1e-012; // de-Broglie wavelength of
    electron , m
7 // We have lambda = h/(m*v) , solving for v
8 v = h/(m*lambda); // Velocity of the electron , m/s
9 K = 1/2*m*v^2; // Kinetic energy of electron , J
10 printf("\nThe kinetic energy of electron having
    wavelength %3.1e m = %4.2e eV", lambda, K/e);
11
12 // Result
13 // The kinetic energy of electron having wavelength
    9.1e-12 m = 1.81e+04 eV

```

---

**Scilab code Exa 1.20** Speed of proton for an equivalent wavelength of that of electron

```

1 // Scilab Code Ex1.20: : Page -1.19 (2009)
2 clc; clear;
3 m_e = 9.1e-031; // Mass of the electron , kg
4 m_p = 1.67e-027; // Mass of the proton , kg
5 v_e = 1; // For simplicity assume velocity of
    electron to be unity , m/s
6 // From de-Broglie relation ,
7 // lambda_p = lambda_e = h(m*v_p) , solving for v_p
8 v_p = m_e*v_e/m_p; // Velocity of the proton , m/s
9 // As lambda_e = h/sqrt(2*m_e*K_e) and lambda_p = h/
    sqrt(2*m_p*K_p) , solving for K_e/K_p
10 K_ratio = m_p/m_e; // Ratio of kinetic energies
    of electron and proton
11
12 printf("\nThe speed of proton for an equivalent

```

```

    wavelength of that of electron = %3.1e ve", v_p);
13 printf("\nRatio of kinetic energies of electron and
      proton = %3.1e, therefore Ke > Kp", K_ratio);
14
15 // Result
16 // The speed of proton for an equivalent wavelength
      of that of electron = 5.4e-04 ve
17 // Ratio of kinetic energies of electron and proton
      = 1.8e+03, therefore Ke > Kp

```

---

### Scilab code Exa 1.21 de Broglie wavelength of the electron

```

1 // Scilab Code Ex1.21: Page-1.20 (2009)
2 clc; clear;
3 V = 50; // Potential difference , V
4 m = 9.1e-031; // Mass of the electron , kg
5 e = 1.6e-019; // Electronic charge , C
6 h = 6.6e-034; // Planck's constant , Js
7 lambda = h/sqrt(2*m*e*V); // From de-Broglie
      relation ,
8 printf("\nde-Broglie wavelength of the electron = %4
      .2 f angstrom", lambda/1e-010);
9
10 // Result
11 // de-Broglie wavelength of the electron = 1.73
      angstrom

```

---

### Scilab code Exa 1.23 Minimum accuracy to locate the position of an electron

```

1 // Scilab Code Ex1.23:: Page-1.31 (2009)
2 clc; clear;
3 v = 740; // Speed of the electron , m/s

```

```

4 m = 9.1e-031;           // Mass of the electron , kg
5 h = 6.6e-034;           // Planck's constant , Js
6 p = m*v;                // Momentum of the electron , kg-m/s
7 frac_v = 0.05/100;      // Correctness in the speed
8 delta_p = p*frac_v;     // Uncertainty in momentum,
                           kg-m/s
9 delta_x = h/(4*pi)*1/delta_p; // Uncertainty in
                                 position , m
10
11 printf("\nThe minimum accuracy to locate the
          position of an electron = %4.2e m",delta_x);
12
13 // Result
14 // The minimum accuracy to locate the position of an
   electron = 1.56e-04 m

```

---

### Scilab code Exa 1.24 Uncertainty in energy of an emitted photon

```

1 // Scilab Code Ex1.24: : Page-1.31 (2009)
2 clc; clear;
3 h = 6.6e-034;           // Planck's constant , Js
4 h_cross = h/(2*pi);    // Reduced Planck's constant
                           , Js
5 delta_t = 1e-010;       // Uncertainty in time , s
6 // From Energy-time uncertainty ,
7 // delta_E*delta_t = h_cross/2, solving for delta_E
8 delta_E = h_cross/(2*delta_t); // Uncertainty in
                               energy of an emitted photon , J
9
10 printf("\nThe uncertainty in energy of an emitted
         photon = %5.3e eV", delta_E/1.6e-019);
11
12 // Result
13 // The uncertainty in energy of an emitted photon =
   3.283e-06 eV

```

---

**Scilab code Exa 1.25** Minimum uncertainty in velocity of electron

```
1 // Scilab Code Ex1.25: : Page-1.31 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 delta_x_max = 1e-007; // Uncertainty in length , m
5 m = 9.1e-031;      // Mass of an electron , kg
6 // From Position-momentum uncertainty ,
7 // delta_p_min = m*delta_v_min = h/delta_x_max ,
8 // solving for delta_v_min
8 delta_v_min = h/(delta_x_max*m);      // Minimum
     uncertainty in velocity of electron , m/s
9
10 printf("\nThe minimum uncertainty in velocity of
         electron = %4.2e m/s" , delta_v_min);
11
12 // Result
13 // The minimum uncertainty in velocity of electron =
    7.25e+03 m/s
```

---

**Scilab code Exa 1.26** Minimum uncertainty in momentum and minimum kinetic energy of proton

```
1 // Scilab Code Ex1.26: Page-1.32 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 delta_x_max = 8.5e-014; // Uncertainty in length ,
     m
5 m = 1.67e-027;    // Mass of proton , kg
6 // From Position-momentum uncertainty ,
7 // delta_p_min*delta_x_max = h, solving for
     delta_p_min
```

```

8 delta_p_min = h/delta_x_max;      // Minimum
    uncertainty in momentum of electron , kg-m/s
9 p_min = delta_p_min;      // Minimum momentum of the
    proton , kg.m/s
10 delta_E = p_min^2/(2*m);
11
12 printf("\nThe minimum uncertainty in momemtum of
    proton = %4.2e kg-m/s", p_min);
13 printf("\nThe kinetic energy of proton = %6.3e eV",
    delta_E/1.6e-019);
14
15 // Result
16 // The minimum uncertainty in momemtum of proton =
    7.76e-21 kg-m/s
17 // The kinetic energy of proton = 1.128e+05 eV

```

---

### Scilab code Exa 1.27 Uncertainty in momentum of electron

```

1 // Scilab Code Ex1.27:: Page-1.32 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
5 E = 0.15*1e+03*e; // Energy of the electron , J
6 m = 9.1e-031;      // Mass of electron , kg
7 delta_x = 0.5e-010; // Position uncertainty of
    electron , m
8 p = (2*m*E)^(1/2); // Momentum of the electron , kg-
    m/s
9 // delta_x*delta_p = h/(4*pi) , solving for delta_p
10 delta_p = h/(4*pi*delta_x); // Uncertainty in
    momentum of electron , kg-m/s
11 frac_p = delta_p/p*100; // Percentage
    uncertainty in momentum of electron , kg-m/s
12
13 printf("\nThe percentage uncertainty in momentum of

```

```

electron = "%2d percent", frac_p);
14
15 // Result
16 // The percentage uncertainty in momentum of
   electron = 15 percent

```

---

### Scilab code Exa 1.28 Uncertainty in position of the particle

```

1 // Scilab Code Ex1.28:: Page-1.33 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
5 delta_v = 7.54e-015; // Uncertainty in velocity
   of the particle , m/s
6 m = 0.25e-06;      // Mass of particle , kg
7 // delta_x*delta_p = h/(4*pi), solving for delta_x
8 delta_x = h/(4*pi*m*delta_v); // Position
   uncertainty of particle , m
9
10 printf("\nThe position uncertainty of particle = %4
.2e m", delta_x);
11
12 // Result
13 // The position uncertainty of particle = 2.79e-14 m

```

---

### Scilab code Exa 1.29 Uncertainty in position of the moving electron

```

1 // Scilab Code Ex1.29:: Page-1.33 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
5 v = 450;           // Velocity of the electron , m/s

```

```

6 delta_v = v*0.05/100;      // Uncertainty in velocity
    of the particle , m/s
7 m = 9.1e-031;   // Mass of electron , kg
8 // delta_x*delta_p = h/(4*pi) , solving for delta_x
9 delta_x = h/(4*pi*m*delta_v); // Position
    uncertainty of particle , m
10
11 printf("\nThe position uncertainty of moving
    electron = %4.2e m", delta_x);
12
13 // Result
14 // The position uncertainty of moving electron =
    2.57e-04 m

```

---

**Scilab code Exa 1.30** Smallest possible uncertainty in position of the electron

```

1 // Scilab Code Ex1.30:: Page-1.33 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 e = 1.6e-019;      // Energy equivalent of 1 eV , J/eV
5 c = 3e+08;         // Speed of light , m/s
6 v = 3e+07;         // Velocity of the electron , m/s
7 m0 = 9.1e-031;     // Rest mass of electron , kg
8 m = m0/sqrt(1-v^2/c^2); // Mass of moving electron ,
    kg
9 delta_p_max = m*v;      // Maximum uncertainty in
    momentum of the particle , m/s
10 // delta_x_min*delta_p_max = h/(4*pi) , solving for
    delta_x_min
11 delta_x_min = h/(4*pi*delta_p_max); // Minimum
    position uncertainty of particle , m
12
13 printf("\nThe smallest possible uncertainty in
    position of the electron = %5.3f angstrom",

```

```

    delta_x_min/1e-010);
14
15 // Result
16 // The smallest possible uncertainty in position of
   the electron = 0.019 angstrom

```

---

**Scilab code Exa 1.31** Difference in the energy between the neighboring levels of Na at the highest state

```

1 // Scilab Code Ex1.31: : Page-1.44 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m = 9.1e-031;      // Electronic mass , kg
5 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
6 l = 2e-002;         // Length of the side of the cube , m
7 E_F = 9*e;          // Fermi energy , J
8 // As E_F = h^2/(8*m*l^2)*(nx^2 + ny^2 + nz^2) and
   nx = ny = nz for a cube , solving for nx
9 nx = sqrt(E_F*(8*m*l^2)/(3*h^2));      // Value of
   integer for a cube
10 E = h^2/(8*m*l^2)*3*nx^2; // Fermi energy , J
11 E1 = h^2/(8*m*l^2)*((nx-1)^2 + nx^2 + nx^2); // 
   Energy of the level just below the fermi level , J
12 delta_E = E - E1;        // Difference in the energy
   between the neighbouring levels of Na at the
   highest state , J
13
14 printf("\nThe energy difference between the
   neighbouring levels of Na at the highest state =
   %4.2e eV", delta_E/e);
15
16 // Result
17 // The energy difference between the neighbouring
   levels of Na at the highest state = 1.06e-07 eV

```

---

**Scilab code Exa 1.32** Energy of the neutron confined in a nucleus

```
1 // Scilab Code Ex1.32:: Page-1.45 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m = 1.67e-027;    // Electronic mass , kg
5 e = 1.6e-019;     // Energy equivalent of 1 eV, J/eV
6 nx = 1, ny = 1, nz = 1; // Principle quantum numbers
                           in 3D corresponding to the longest energy state
7 lx = 1e-014, ly = 1e-014, lz = 1e-014;      //
                           Dimensions of the box to which the neutron is
                           confined , m
8 E = h^2/(8*m)*(nx^2/lx^2+ny^2/ly^2+nz^2/lz^2);   //
                           Energy of the neutron confined in the nucleus , J
9
10 printf("\nThe energy of the neutron confined in a
           nucleus = %4.2e eV", E/e);
11
12 // Result
13 // The energy of the neutron confined in a nucleus =
           6.11e+06 eV
```

---

**Scilab code Exa 1.33** Energy of an electron moving in one dimensional infinitely high potential box

```
1 // Scilab Code Ex1.33:: Page-1.46 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m = 9.1e-031;     // Electronic mass , kg
5 e = 1.6e-019;     // Energy equivalent of 1 eV, J/eV
6 n = 1;             // For simplicity assume principle
                     quantum number to be unity
```

```

7 l = 2.1e-010; // Length of one dimensional
    potential box, m
8 E = h^2*n^2/(8*m*l^2); // Energy of the electron ,
    J
9
10 printf("\nThe energy of the electron moving in one
    dimensional infinitely high potential box = %4.2f
    n^2 eV", E/e);
11
12 // Result
13 // The energy of the electron moving in one
    dimensional infinitely high potential box = 8.48
    n^2 eV

```

---

**Scilab code Exa 1.34** Lowest energy of an electron in a one dimensional force free region

```

1 // Scilab Code Ex1.34:: Page-1.46 (2009)
2 clc; clear;
3 h = 6.6e-034; // Planck's constant , Js
4 m = 9.1e-031; // Electronic mass , kg
5 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
6 n = 1; // The lowest energy state of electron
7 l = 3.5e-010; // Length of one dimensional
    potential box, m
8 E = h^2*n^2/(8*m*l^2); // Energy of the electron
    in the lowest state , J
9
10 printf("\nThe lowest energy of the electron in a one
    dimensional force free region = %1d eV", E/e);
11
12 // Result
13 // The lowest energy of an electron in a one
    dimensional force free region = 3 eV

```

---

**Scilab code Exa 1.35** First three energy levels of an electron in one dimensional box

```
1 // Scilab Code Ex1.35:: Page-1.46 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m = 9.1e-031;      // Electronic mass , kg
5 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
6 l = 9.5e-010;      // Length of one dimensional
                      potential box , m
7
8 // First energy level
9 n = 1;              // The first energy state of electron
10 E1 = h^2*n^2/(8*m*l^2);    // Energy of the electron
                               in first state , J
11
12 // Second energy level
13 n = 2;              // The second energy state of electron
14 E2 = h^2*n^2/(8*m*l^2);    // Energy of the electron
                               in second state , J
15
16 // Third energy level
17 n = 3;              // The third energy state of electron
18 E3 = h^2*n^2/(8*m*l^2);    // Energy of the electron
                               in third state , J
19
20 printf("\nThe energy of the electron in first state
          = %4.1e J", E1);
21 printf("\nThe energy of the electron in second state
          = %4.1e J", E2);
22 printf("\nThe energy of the electron in third state
          = %4.1e J", E3);
23
24 // Result
```

```

25 // The energy of the electron in first state = 6.6e
   -20 J
26 // The energy of the electron in second state = 2.7
   e-19 J
27 // The energy of the electron in third state = 6.0e
   -19 J

```

---

**Scilab code Exa 1.36** Lowest two permitted energy values of the electron in a 1D box

```

1 // Scilab Code Ex1.36:: Page-1.47 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m = 9.1e-031;      // Electronic mass , kg
5 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
6 l = 2.5e-010;      // Length of one dimensional
                      potential box , m
7
8 // First energy level
9 n = 1;              // The lowest energy state of electron
10 E1 = h^2*n^2/(8*m*l^2);    // Energy of the electron
                                in first state , J
11
12 // Second energy level
13 n = 2;              // The second energy state of electron
14 E2 = h^2*n^2/(8*m*l^2);    // Energy of the electron
                                in second state , J
15
16 printf("\nThe energy of the electron in lowest state
          = %5.2f eV", E1/e);
17 printf("\nThe energy of the electron in second state
          = %5.2f eV", E2/e);
18
19
20 // Result

```

```
21 // The energy of the electron in lowest state =  
    5.98 eV  
22 // The energy of the electron in second state =  
    23.93 eV
```

---

**Scilab code Exa 1.37** Lowest energy of the neutron confined to the nucleus

```
1 // Scilab Code Ex1.37:: Page -1.47 (2009)  
2 clc; clear;  
3 h = 6.6e-034;      // Planck's constant , Js  
4 m = 1.67e-027;    // Electronic mass , kg  
5 e = 1.6e-019;     // Energy equivalent of 1 eV, J/eV  
6 l = 2.5e-010;     // Length of one dimensional  
                     potential box , m  
7 delta_x = 1e-014;   // Uncertainty in position of  
                     neutron , m  
8 // From uncertainty principle ,  
9 // delta_x*delta_p = h/(4*pi) , solving for delta_p  
10 delta_p = h/(4*pi*delta_x); // Uncertainty in  
                     momentum of neutron , kg-m/s  
11 p = delta_p;        // Momemtum of neutron in the box ,  
                     kg-m/s  
12 KE = p^2/(2*m);   // Kinetic energy of neutron in the  
                     box , J  
13  
14 printf("\nThe lowest energy of the neutron confined  
        to the nucleus = %4.2f MeV", KE/(e*1e+06));  
15  
16 // Result  
17 // The lowest energy of the neutron confined to the  
    nucleus = 0.05 MeV
```

---

### Scilab code Exa 1.38 X ray scattering

```
1 // Scilab Code Ex1.38: : Page-1.56 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m0 = 9.1e-031;     // Electronic mass , kg
5 c = 3e+08;         // Speed of light , m/s
6 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
7 phi = 45;          // Scattering angle of X-rays , degrees
8 E = 75;            // Incident energy of X-rays , keV
9 // As from Compton shift formula
10 //  $1/E_{\text{prime}} - 1/E = 1/(m_0 \cdot c^2) \cdot (1 - \cos(\phi))$ 
11 // Solving for E_prime
12 E_prime = 1/((1/(m0*c^2/(e*1e+03)))*(1-cosd(phi))+1/
   E); // Energy of scattered photon , keV
13 E_recoil = E - E_prime;      // Energy of recoil
   electron , keV
14
15 printf("\nThe energy of scattered X-ray = %4.1f keV"
   , E_prime);
16 printf("\nThe energy of recoil electron = %3.1f keV"
   , E_recoil);
17
18 // Result
19 // The energy of scattered X-ray = 71.9 keV
20 // The energy of recoil electron = 3.1 keV
```

---

### Scilab code Exa 1.39 Wavelength of scattered Xray

```
1 // Scilab Code Ex1.39: : Page-1.57 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m0 = 9.1e-031;     // Electronic mass , kg
5 c = 3e+08;         // Speed of light , m/s
6 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
```

```

7 phi = 60;      // Scattering angle of X-rays , degrees
8 E = 75;        // Incident energy of X-rays , keV
9 // As from Compton shift formula
10 delta_L = h/(m0*c)*(1-cosd(phi));    // Change in
    photon wavelength , m
11 lambda = 0.198e-010;      // Wavelength of incident
    photon , m
12 lambda_prime = (lambda+delta_L)/1e-010; // 
    Wavelength of scattered X-ray , angstrom
13
14 printf("\nThe wavelength of scattered X-ray = %6.4f
    angstrom" , lambda_prime);
15
16 // Result
17 // The wavelength of scattered X-ray = 0.2101
    angstrom

```

---

**Scilab code Exa 1.40** Wavelength of scattered radiation with changed angle of view

```

1 // Scilab Code Ex1.40:: Page-1.57 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m0 = 9.1e-031;    // Electronic mass, kg
5 c = 3e+08;         // Speed of light , m/s
6 e = 1.6e-019;     // Energy equivalent of 1 eV, J/eV
7 phi = 180;         // Scattering angle of X-rays ,
    degrees
8 lambda = 1.78;     // Wavelength of incident photon ,
    m
9 lambda_prime = 1.798; // Wavelength of scattered X-
    ray , angstrom
10 // As from Compton shift formula
11 // lambda_prime - lambda = h/(m0*c)*(1-cosd(phi)) ,
    Change in photon wavelength , m

```

```

12 // Or we may write , lambda_prime - lambda = k*(1-
   cosd(phi))
13 // solving for k
14 k = (lambda_prime - lambda)/(1-cosd(phi)); // k = h
   /(m0*c) value , angstrom
15
16 // For phi = 60
17 phi = 60;           // New angle of scattering , degrees
18 lambda_prime = lambda + k*(1-cosd(phi)); //
   Wavelength of scattered radiation at 60 degree
   angle , angstrom
19 printf("\nThe wavelength of scattered X-ray at %d
   degrees view = %6.4f angstrom", phi, lambda_prime
);
20 // Recoil energy of electron
21 E = h*c*(1/lambda - 1/lambda_prime)*1e+010;      //
   Recoil energy of electron , joule
22 printf("\nThe recoil energy of electron scattered
   through %d degrees = %4.1f eV", phi, E/e);
23
24 // Result
25 // The wavelength of scattered X-ray at 60 degrees
   view = 1.7845 angstrom
26 // The recoil energy of electron scattered through
   60 degrees = 17.5 eV

```

---

### Scilab code Exa 1.41 Compton scattering through aluminium foil

```

1 // Scilab Code Ex1.41:: Page-1.58 (2009)
2 clc; clear;
3 h = 6.6e-034;      // Planck's constant , Js
4 m0 = 9.1e-031;     // Electronic mass , kg
5 c = 3e+08;         // Speed of light , m/s
6 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
7 phi = 90;          // Scattering angle of X-rays , degrees

```

```

8 E = 510*1e+03*e;      // Energy of incident photon , J
9 // As E = h*c/lambda , solving for lambda
10 lambda = h*c/E;      // Wavelength of incident photon ,
   m
11 // As from Compton shift formula
12 // lambda_prime - lambda = h/(m0*c)*(1-cosd(phi)) ,
   solving for lambda_prime
13 lambda_prime = lambda + h/(m0*c)*(1-cosd(phi)); // 
   Wavelength of scattered X-ray , m
14 printf("\nThe wavelength of scattered X-ray as
   viewed at %d degrees = %4.2e m", phi,
   lambda_prime);
15
16 // Recoil energy of electron
17 E = h*c*(1/lambda - 1/lambda_prime);      // Recoil
   energy of electron , joule
18 printf("\nThe recoil energy of electron scattered
   through %d degrees = %4.2e eV", phi, E/e);
19
20 // Direction of recoil electron
21 theta = atan(lambda*sind(phi)/(lambda_prime-lambda*
   cosd(phi))); // Direction of recoil electron ,
   degrees
22 printf("\nThe direction of emission of recoil
   electron = %5.2f degrees", theta);
23
24
25 // Result
26 // The wavelength of scattered X-ray as viewed at 90
   degrees = 4.84e-12 m
27 // The recoil energy of electron scattered through
   90 degrees = 2.55e+05 eV
28 // The direction of emission of recoil electron =
   26.61 degrees

```

---

### Scilab code Exa 1.42 Energetic electrons in the Xray tube

```
1 // Scilab Code Ex1.42: : Page-1.59 (2009)
2 clc; clear;
3 m = 9.1e-031;      // Electronic mass , kg
4 c = 3e+08;         // Speed of light , m/s
5 e = 1.6e-019;      // Charge on the electron , C
6 V = 12.4e+03;      // Potential difference applied
                      across the X-ray tube , V
7 i = 2e-03;          // Current through the X-ray tube , A
8 t = 1;              // Time for which the electrons strike the
                      target material , s
9 N = i*t/e;          // Number of electrons striking the
                      target per sec , per sec
10 v_max = sqrt(2*e*V/m); // Maximum speed of the
                           electrons , m/s
11
12 printf("\nThe number of electrons striking the
           target per sec = %4.2e electrons/sec", N);
13 printf("\nThe maximum speed of the electrons when
           they strike = %3.1e m/s", v_max);
14
15
16 // Result
17 // The number of electrons striking the target per
       sec = 1.25e+16 electrons/sec
18 // The maximum speed of the electrons when they
       strike = 6.6e+07 m/s
```

---

# Chapter 2

## Interference

**Scilab code Exa 2.1** Slit separation in Double Slit experiment

```
1 // Scilab Code Ex2.1:: Page-2.9 (2009)
2 clc; clear;
3 lambda = 5893e-008; // Wavelength of light used, m
4 D = 200; // Distance of the source from the
           screen , m
5 b = 0.2; // Fringe separation , cm
6 d = lambda*D/b; // Separation between the slits , cm
7
8 printf("\nThe separation between the slits = %3.1e
         cm", d);
9
10 // Result
11 // The separation between the slits = 5.9e-002 cm
```

---

**Scilab code Exa 2.2** Wavelength of light in Young Double Slit experiment

```
1 // Scilab Code Ex2.2:: Page-2.10 (2009)
2 clc; clear;
```

```

3 d = 0.2; // Separation between the slits , cm
4 D = 100; // Distance of the source from the
    screen , m
5 b = 0.35e-01; // Fringe separation , cm
6 lambda = b*d/D; // Wavelength of light used , m
7 printf("\nThe wavelength of the light = %3.1e cm" ,
    lambda);
8
9 // Result
10 // The wavelength of the light = 7.0e-005 cm

```

---

**Scilab code Exa 2.3** Ratio of maximum intensity to minimum intensity of interference fringes

```

1 // Scilab Code Ex2 .3:: Page -2.10 (2009)
2 clc; clear;
3 I2 = 1; // For simplicity assume intensity from
    slit 2 to be unity , W/sq-m
4 I1 = I2*25; // Intensity from slit 1, W/sq-m
5 I_ratio = I1/I2; // Intensity ratio
6 a_ratio = sqrt(I_ratio); // Amplitude ratio
7 a2 = 1; // For simplicity assume amplitude from
    slit 2 to be unity , m
8 a1 = a_ratio*a2; // Amplitude from slit 1, m
9 I_max = (a1 + a2)^2; // Maximum intensity of wave
    during interference , W/sq-m
10 I_min = (a1 - a2)^2; // Minimum intensity of wave
    during interference , W/sq-m
11 cf = 4; // Common factor
12 printf("\nThe ratio of maximum intentisy to minimum
    intensity of interference fringes = %d/%d" , I_max
    /cf , I_min/cf);
13
14 // Result
15 // The ratio of maximum intentisy to minimum

```

intensity of interference fringes = 9/4

---

**Scilab code Exa 2.4** Wavelength of light from monochromatic coherent sources

```
1 // Scilab Code Ex2.4:: Page-2.10 (2009)
2 clc; clear;
3 d = 0.02; // Separation between the slits , cm
4 D = 100; // Distance of the source from the
            screen , m
5 n = 6; // No. of bright fringe from the centre
6 x = 1.22; // Position of 6th bright fringe , cm
7 lambda = x*d/(n*D); // Wavelength of light used , m
8 printf("\nThe wavelength of the light from coherent
        sources = %5.3e cm", lambda);
9
10 // Result
11 // The wavelength of the light from coherent sources
    = 4.067e-005 cm
```

---

**Scilab code Exa 2.5** Separation between fourth order dark fringes

```
1 // Scilab Code Ex2.5:: Page-2.10 (2009)
2 clc; clear;
3 lambda1 = 5890e-008; // Wavelength of D1 line of
                      sodium , cm
4 lambda2 = 5896e-008; // Wavelength of D2 line of
                      sodium , cm
5 D = 120; // Distance between source and the
            screen , cm
6 d = 0.025; // Separation between the slits , cm
7 n = 4; // Order of dark fringe
```

```

8 x1 = (2*n+1)*lambda1*D/(2*d); // Position of 4th
      dark fringe due to D1 line , cm
9 x2 = (2*n+1)*lambda2*D/(2*d); // Position of 4th
      dark fringe due to D2 line , cm
10 delta_x = x2-x1; // Fringe separation , cm
11
12 printf("\nThe separation between fourth order dark
      fringes = %4.2e cm", x2-x1);
13
14 // Result
15 // The separation between fourth order dark fringes
      = 1.30e-03 cm

```

---

### Scilab code Exa 2.6 Distance between two coherent sources

```

1 // Scilab Code Ex2.6:: Page -2.11 (2009)
2 clc; clear;
3 lambda = 5500e-008; // Wavelength of light used ,
      cm
4 Y1 = 10; // Distance of biprism from the source ,
      cm
5 Y2 = 90; // Distance of biprism from the screen ,
      cm
6 D = Y1 + Y2; // Distance between slits and the
      screen , cm
7 b = 8.526e-02; // Fringe width , cm
8 d = lambda*D/b; // Separation between the slits , cm
9
10 printf("\nThe distance between two coherent sources
      = %4.2e cm", d);
11
12 // Result
13 // The distance between two coherent sources = 6.45e
      -02 cm

```

---

**Scilab code Exa 2.7** Fringe width of the interference pattern due to biprism

```
1 // Scilab Code Ex2.7:: Page-2.11 (2009)
2 clc; clear;
3 alpha = %pi/180;      // Acute angle of biprism ,
radian
4 mu = 1.5;      // Refractive index of biprism
5 lambda = 5500e-008;    // Wavelength of light used ,
cm
6 y1 = 5;        // Distance of biprism from the source ,
cm
7 y2 = 75;       // Distance of biprism from the screen ,
cm
8 D = y1 + y2;    // Distance between slits and the
screen , cm
9 d = 2*(mu-1)*alpha*y1; // Separation between the
slits , cm
10 b = lambda*D/d;   // Fringe width of the interference
pattern due to biprism , cm
11
12 printf("\nThe fringe width of the interference
pattern due to biprism = %4.2e cm", b);
13
14 // Result
15 // The fringe width of the interference pattern due
to biprism = 5.04e-02 cm
```

---

**Scilab code Exa 2.8** Angle of vertex of the biprism

```
1 // Scilab Code Ex2.8:: Page-2.11 (2009)
2 clc; clear;
3 mu = 1.5;      // Refractive index of biprism
```

```

4 lambda = 5500e-008;      // Wavelength of light used ,
    cm
5 y1 = 5;      // Distance of biprism from the source ,
    cm
6 y2 = 95;      // Distance of biprism from the screen ,
    cm
7 D = y1 + y2;      // Distance between slits and the
    screen , cm
8 b = 0.025;      // Fringe width of the interference
    pattern due to biprism , cm
9 // As d = 2*(mu-1)*alpha*y1, solving for alpha
10 alpha = lambda*D/(b*2*(mu-1)*y1)      // Angle of
    vertex of the biprism , radian
11
12 printf("\nThe angle of vertex of the biprism = %3.1e
    rad", alpha);
13
14 // Result
15 // The angle of vertex of the biprism = 4.4e-02 rad

```

---

**Scilab code Exa 2.9** Number of interference fringes for changed wavelength

```

1 // Scilab Code Ex2.9:: Page-2.12 (2009)
2 clc; clear;
3 n1 = 69;      // Number of interference fringes
    obtained with yellow wavelength
4 lambda1 = 5893e-008;      // Wavelength of yellow
    light used , cm
5 lambda2 = 5461e-008;      // Wavelength of green light
    used , cm
6 // As n*lambda = l*D = constant , therefore
7 n2 = n1*lambda1/lambda2;      // Number of
    interference fringes for green wavelength
8
9 printf("\nThe number of interference fringes for

```

```
changed wavelength = %2d" , ceil(n2));  
10  
11 // Result  
12 // The number of interference fringes for changed  
wavelength = 75
```

---

**Scilab code Exa 2.10** Wavelength of light in a biprism experiment

```
1 // Scilab Code Ex2.10:: Page-2.12 (2009)  
2 clc; clear;  
3 D = 100;      // Distance between slits and the screen  
               , cm  
4 d = 0.08;    // Separation between the slits , cm  
5 b = 2.121/25; // Fringe width of the interference  
                  pattern due to biprism , cm  
6 lambda = b*d/D; // Wavelength of light in a  
                   biprism experiment , cm  
7  
8 printf("\nThe wavelength of light in a biprism  
       experiment = %5.0f angstrom" , lambda/1e-008);  
9  
10 // Result  
11 // The wavelength of light in a biprism experiment =  
   6787 angstrom
```

---

**Scilab code Exa 2.11** Fringe width at a certain distance from biprism

```
1 // Scilab Code Ex2.11:: Page-2.13 (2009)  
2 clc; clear;  
3 alpha = %pi/180;      // Acute angle of biprism ,  
                     radian  
4 mu = 1.5;      // Refractive index of biprism
```

```

5 lambda = 5900e-008;      // Wavelength of light used ,
   cm
6 y1 = 10;      // Distance of biprism from the source ,
   cm
7 y2 = 100;     // Distance of biprism from the screen ,
   cm
8 D = y1 + y2;    // Distance between slits and the
   screen , cm
9 d = 2*(mu-1)*alpha*y1; // Separation between the
   slits , cm
10 b = lambda*D/d; // Fringe width of the interference
   pattern due to biprism , cm
11
12 printf("\nThe fringe width at a distance of %d cm
   from biprism = %4.2e cm", y2, b);
13
14 // Result
15 // The fringe width at a distance of 100 cm from
   biprism = 3.72e-02 cm

```

---

**Scilab code Exa 2.12** Distance between coherent sources in biprism experiment

```

1 // Scilab Code Ex2.12:: Page -2.13 (2009)
2 clc; clear;
3 lambda = 5893e-008;      // Wavelength of light used ,
   cm
4 y1 = 10;      // Distance of biprism from the source ,
   cm
5 y2 = 100;     // Distance of biprism from the screen ,
   cm
6 D = y1 + y2;    // Distance between slits and the
   screen , cm
7 b = 3.5e-02; // Fringe width of the interference
   pattern due to biprism , cm

```

```

8 d = lambda*D/b; // Distance between coherent
sources , cm
9
10 printf("\nThe distance between coherent sources = %5
.3f cm", d);
11
12 // Result
13 // The distance between coherent sources = 0.185 cm

```

---

**Scilab code Exa 2.13** Effect of slit separation on fringe width

```

1 // Scilab Code Ex2.13:: Page-2.13 (2009)
2 clc; clear;
3 b = 0.125; // Fringe width of the interference
pattern due to biprism , cm
4 d = 1; // For simplicity assume distance between
sources to be unity , cm
5 d_prime = 3/4*d; // New distance between sources ,
cm
6 // As b is proportional to 1/d, so
7 b_prime = b*d/d_prime; // New fringe width of the
interference pattern due to biprism , cm
8
9 printf("\nThe new value of fringe width due to
reduced slit separation = %5.3f cm", b_prime);
10
11 // Result
12 // The new value of fringe width due to reduced slit
separation = 0.167 cm

```

---

**Scilab code Exa 2.14** Effect of slit biprism separation on fringe width

```
1 // Scilab Code Ex2.14:: Page-2.13 (2009)
```

```

2 clc; clear;
3 b = 0.187; // Fringe width of the interference
    pattern due to biprism , cm
4 y1 = 1; // For simplicity assume distance between
    slit and biprism to be unity , cm
5 y1_prime = 1.25*y1; // New distance between slit
    and biprism , cm
6 // As d is directly proportional to y1 and b is
    directly proportional to d, so
7 // b is inversely proportional to y1
8 b_prime = b*y1/y1_prime; // New fringe width of the
    interference pattern due to biprism , cm
9
10 printf("\nThe new value of fringe width due to
    increased slit - biprism separation = %5.3f cm",
    b_prime);
11
12 // Result
13 // The new value of fringe width due to increased
    slit - biprism separation = 0.150 cm

```

---

### Scilab code Exa 2.15 Distance between interference bands

```

1 // Scilab Code Ex2.15:: Page-2.14 (2009)
2 clc; clear;
3 d1 = 5e-01; // First distance between images of the
    slit , cm
4 d2 = 2.25e-01; // Second distance between images of
    the slit , cm
5 lambda = 5896e-008; // Wavelength of the light used ,
    cm
6 D = 120; // Distance between screen and the slits
    , cm
7 d = sqrt(d1*d2); // Geometric mean of distance
    between the two slits , cm

```

```

8 b = lambda*D/d;      // Distance between interference
bands , cm
9
10 printf("\nThe distance between interference bands =
%5.3e cm", b);
11
12 // Result
13 // The distance between interference bands = 2.109e
-02 cm

```

---

### Scilab code Exa 2.16 Angle of vertex of Fresnel biprism

```

1 // Scilab Code Ex2.16:: Page-2.14 (2009)
2 clc; clear;
3 mu = 1.5;    // Refractive index of biprism
4 lambda = 5500e-008;    // Wavelength of light used,
cm
5 y1 = 25;    // Distance of biprism from the source ,
cm
6 y2 = 150;    // Distance of biprism from the screen ,
cm
7 D = y1 + y2;    // Distance between slits and the
screen , cm
8 b = 0.05;    // Fringe width of the interference
pattern due to biprism , cm
9 // As d = 2*(mu-1)*alpha*y1, solving for alpha
10 alpha = lambda*D/(b*2*(mu-1)*y1)    // Angle of
vertex of the biprism , radian
11
12 printf("\nThe angle of vertex of the biprism = %6.4f
rad", alpha);
13
14 // Result
15 // The angle of vertex of the biprism = 0.0077 rad

```

---

**Scilab code Exa 2.17** Wavelength of light used in biprism experiment to illuminate slits

```
1 // Scilab Code Ex2.17:: Page-2.15 (2009)
2 clc; clear;
3 theta = 178;      // Vertex angle of biprism , degrees
4 alpha = (180-theta)/2*pi/180;      // Acute angle of
    biprism , radian
5 mu = 1.5;        // Refractive index of biprism
6 y1 = 20;         // Distance of biprism from the source ,
    cm
7 y2 = 125;        // Distance of biprism from the screen ,
    cm
8 D = y1 + y2;    // Distance between slits and the
    screen , cm
9 d = 2*(mu-1)*alpha*y1;    // Separation between the
    slits , cm
10 b = 0.025;      // Fringe width of the interference
    pattern due to biprism , cm
11 lambda = b*d/D;    // Wavelength of light used , cm
12
13 printf("\nThe wavelength of light used to illuminate
    slits = %4d angstrom", lambda/1e-08);
14
15 // Result
16 // The wavelength of light used to illuminate slits
    = 6018 angstrom
```

---

**Scilab code Exa 2.18** Vertex angle of Fresnel biprism

```
1 // Scilab Code Ex2.18:: Page-2.15 (2009)
2 clc; clear;
```

```

3 mu = 1.5;      // Refractive index of biprism
4 lambda = 6600e-008;    // Wavelength of light used ,
cm
5 y1 = 40;       // Distance of biprism from the source ,
cm
6 y2 = 175;      // Distance of biprism from the screen ,
cm
7 D = y1 + y2;   // Distance between slits and the
screen , cm
8 b = 0.04;      // Fringe width of the interference
pattern due to biprism , cm
9 // As d = 2*(mu-1)*alpha*y1, solving for alpha
10 alpha = lambda*D/(b*2*(mu-1)*y1)    // Acute angle
of the biprism , radian
11 theta = (%pi-2*alpha);   // Vertex angle of the
biprism , radian
12
13 printf("\nThe vertex angle of the biprism = %6.2 f
degrees", theta*180/%pi);
14
15 // Result
16 // The vertex angle of the biprism = 178.98 degrees

```

---

**Scilab code Exa 2.19** Order of visible fringe for changed wavelength of light

```

1 // Scilab Code Ex2.19: : Page - 2.16 (2009)
2 clc; clear;
3 lambda1 = 7000e-008;    // Original wavelength of
light , cm
4 lambda2 = 5000e-008;    // New wavelength of light ,
cm
5 n1 = 10;      // Order of the fringes with original
wavelength
6 // As x = n*lambda*D/d, so n*lambda = constant

```

```

7 // n1*lambda1 = n2*lambda2, solving for n2
8 n2 = n1*lambda1/lambda2;      // Order of visible
      fringe for changed wavelength of light
9
10 printf("\nThe order of visible fringe for changed
      wavelength of light = %2d", ceil(n2));
11
12 // Result
13 // The order of visible fringe for changed
      wavelength of light = 14

```

---

### Scilab code Exa 2.20 Angle of vertex of biprism

```

1 // Scilab Code Ex1.20:: Page-2.16 (2009)
2 clc; clear;
3 y1 = 40;      // Distance between biprism from the
               slit , cm
4 D = 160;      // Distance between slit and the screen ,
               cm
5 mu = 1.52;    // Refractive index of material of the
               prism
6 lambda = 5893e-008; // Wavelength of light used , cm
7 b = 0.01;      // Fringe width , cm
8 // As b = lambda*D/d, solving for d
9 d = lambda*D/b;      // Distance between virtual
               sources , cm
10 // But d = 2*y1*(mu-1)*alpha, solving for alpha
11 alpha = d/(2*y1*(mu-1))*180/%pi;      // Angle of
               biprism , degrees
12 theta = 180-2*alpha;      // Angle of vertex of
               biprism , degrees
13
14 printf("\nThe angle of vertex of biprism = %5.1f
      degree", theta);
15

```

```
16 // Result  
17 // The angle of vertex of biprism = 177.4 degree
```

---

**Scilab code Exa 2.21** Separation between two coherent sources

```
1 // Scilab Code Ex2.21: : Page-2.16 (2009)  
2 clc; clear;  
3 lambda = 6000e-008;      // Wavelength of light used ,  
                           cm  
4 D = 100;      // Distance between slits and the screen  
                           , cm  
5 b = 0.05;     // Fringe width of the interference  
                           pattern due to biprism , cm  
6 d = lambda*D/b; // Distance between coherent  
                           sources , cm  
7  
8 printf("\nThe distance between coherent sources = %3  
       .1f mm" , d/1e-01);  
9  
10 // Result  
11 // The distance between coherent sources = 1.2 mm
```

---

**Scilab code Exa 2.22** Refractive index of the glass sheet

```
1 // Scilab Code Ex2.22:: Page-2.19 (2009)  
2 clc; clear;  
3 t = 3.2e-04;      // Thickness of the glass sheet , cm  
4 lambda = 5500e-008; // Wavelength of light used ,  
                           cm  
5 n = 5;           // Order of interference fringes  
6 // As path difference (mu - 1)*t = n*lambda  
7 mu = n*lambda/t + 1; // Refractive indexof the  
                           glass sheet
```

```
8
9 printf("\nThe refractive index of the glass sheet=
    %4.2f", mu);
10
11 // Result
12 // The refractive index of the glass sheet= 1.86
```

---

### Scilab code Exa 2.23 Refractive index of material of sheet

```
1 // Scilab Code Ex2.23:: Page-2.19 (2009)
2 clc; clear;
3 t = 2.1e-03;      // Thickness of the glass sheet , cm
4 lambda = 5400e-008;      // Wavelength of light used ,
    cm
5 n = 11;          // Order of interference fringes
6 // As path difference , (mu - 1)*t = n*lambda
7 mu = n*lambda/t + 1;      // Refractive index of the
    glass sheet
8
9 printf("\nThe refractive index of the glass sheet =
    %4.2f", mu);
10
11 // Result
12 // The refractive index of the glass sheet= 1.28
```

---

### Scilab code Exa 2.24 Wavelength of light used in biprism arrangement

```
1 // Scilab Code Ex2.24:: Page-2.19 (2009)
2 clc; clear;
3 t = 9.21e-05;      // Thickness of the mica sheet , cm
4 mu = 1.5;          // Refractive index of material of sheet
5 n = 1;            // Order of interference fringes
```

```

6 // As path difference , (mu - 1)*t = n*lambda ,
   solving for lambda
7 lambda = (mu - 1)*t/n;      // Wavelength of light
   used , cm
8
9 printf("\nThe wavelength of light used = %5.3e cm" ,
   lambda);
10
11 // Result
12 // The wavelength of light used = 4.605e-005 cm

```

---

**Scilab code Exa 2.25** Thickness of the transparent sheet

```

1 // Scilab Code Ex2.25:: Page-2.19 (2009)
2 clc; clear;
3 lambda = 5890e-008;      // Wavelength of light used ,
   cm
4 mu = 1.5;    // Refractive index of material sheet
5 // As shift = 9*lambda*D/d = D/d*(mu - 1)*t , solving
   for t
6 t = 9*lambda/(mu - 1);    // Thickness of the glass
   sheet , cm
7 printf("\nThe thickness of the glass sheet = %4.2e
   cm" , t);
8
9 // Result
10 // The thickness of the glass sheet = 1.06e-003 cm

```

---

**Scilab code Exa 2.26** Thickness of the transparent sheet from fringe shift

```

1 // Scilab Code Ex2.26:: Page-2.20 (2009)
2 clc; clear;

```

```

3 lambda = 5400e-008;      // Wavelength of light used ,
   cm
4 mu = 1.7;    // Refractive index of material sheet
   convering the first slit
5 mu_prime = 1.5;    // Refractive index of material
   sheet convering the seecond slit
6 // As shift , S = D/d*(mu - mu_prime)*t = b/lambda*(mu - mu_prime)*t , solving for t
7 t = 8*lambda/(mu-mu_prime)    // Thickness of the
   glass sheet , cm
8
9 printf("\nThe thickness of the glass sheet = %4.2e
   cm" , t);
10
11 // Result
12 // The thickness of the glass sheet = 2.16e-003 cm

```

---

### Scilab code Exa 2.27 Refractive index of thin mica sheet

```

1 // Scilab Code Ex2.27:: Page-2.20 (2009)
2 clc; clear;
3 t = 21.5e-05;      // Thickness of the glass sheet , cm
4 lambda = 5890e-008;    // Wavelength of light used ,
   cm
5 n = 1;      // Order of interference fringes
6 // As path difference , (mu - 1)*t = n*lambda
7 mu = n*lambda/t + 1;    // Refractive indexof the
   glass sheet
8
9 printf("\nThe refractive index of the glass sheet =
   %5.3f" , mu);
10
11 // Result
12 // The refractive index of the glass sheet = 1.274

```

---

**Scilab code Exa 2.28** Wavelength of light used in double slit experiment

```
1 // Scilab Code Ex2.28:: Page-2.20 (2009)
2 clc; clear;
3 D = 1; // For simplicity assume distance between
        source and slits to be unity , unit
4 d = 1; // For simplicity assume slit separation to
        be unity , unit
5 t = 2.964e-06; // Thickness of the mica sheet , cm
6 mu = 1.5; // Refractive index of material of shee
7 L = poly(0, 'L');
8 // As b = b_prime or 2.25*D*L/d = D/d*(mu-1)*t , or
    we may write
9 L = roots(2.25*D*L/d-D/d*(mu-1)*t);      //
    Wavelength of the light used , m
10
11 printf("\nThe wavelength of the light used = %4.0f
        angstrom", L/1e-010);
12
13 // Result
14 // The wavelength of the light used = 6587 angstrom
```

---

**Scilab code Exa 2.29** Thickness of mica sheet from central fringe shift

```
1 // Scilab Code Ex2.29:: Page-2.21 (2009)
2 clc; clear;
3 lambda = 5890e-008; // Wavelength of light used ,
        cm
4 n = 5; // Order of interference fringes
5 mu = 1.5; // Refractive index of the mica sheet
6 // As path difference , (mu - 1)*t = n*lambda ,
    solving for t
```

```

7 t = n*lambda/(mu-1);      // Thickness of the mica
    sheet , cm
8
9 printf("\nThe thickness of the mica sheet = %4.2e cm
    ", t);
10
11 // Result
12 // The thickness of the mica sheet = 5.89e-004 cm

```

---

**Scilab code Exa 2.30** Refractive index of material from shifting fringe pattern

```

1 // Scilab Code Ex2.30:: Page -2.21 (2009)
2 clc; clear;
3 b = 1;          // For simplicity assume fringe width to
    be unity , cm
4 S = 30*b;      // Fringe shift , cm
5 lambda = 6600e-008;      // Wavelength of light used ,
    cm
6 t = 4.9e-003;      // Thickness of the film , cm
7 // As S = b/lambda*(mu-1)*t, solving for mu
8 mu = S*lambda/t + 1;      // Refractive index of
    material from shifting fringe pattern
9
10 printf("\nThe refractive index of material from
    shifting fringe pattern = %3.1f", mu);
11
12 // Result
13 // The refractive index of material from shifting
    fringe pattern = 1.4

```

---

**Scilab code Exa 2.31** Fringe width and optical path change during interference of waves through mica and glass

```

1 // Scilab Code Ex2.31:: Page -2.22 (2009)
2 clc; clear;
3 mu1 = 1.55;      // Refractive index of mica
4 mu2 = 1.52;      // Refractive index of glass
5 t = 0.75e-003;   // Thickness of the sheets , m
6 d = 0.25e-02;    // Separation between the slits , m
7 lambda = 5896e-010; // Wavelength of light used , m
8 D = 1.5;         // Distance between the source ans the
                   slits , m
9 // Fringe width
10 b = lambda*D/d; // Fringe width , m
11 // Optical path difference
12 delta_x = (mu1-1)*t-(mu2-1)*t; // Optical path
                   change , m
13
14 printf("\nThe fringe width = %3.1e m" , b);
15 printf("\nThe optical path change = %5.3e m" ,
       delta_x);
16
17 // Result
18 // The fringe width = 3.5e-004 m
19 // The optical path change = 2.250e-005 m

```

---

**Scilab code Exa 2.32** Thickness of mica sheet from Fresnel biprism experiment

```

1 // Scilab Code Ex2.32:: Page -2.22 (2009)
2 clc; clear;
3 b = 1;          // For simplicity assume fringe width to
                  be unity , cm
4 S = 3*b;        // Fringe shift , cm
5 lambda = 5890e-008; // Wavelength of light used ,
                  cm
6 mu = 1.6;        // Refractive index of the mica sheet
7 // As S = b/lambda*(mu-1)*t , solving for t

```

```

8 t = S*lambda/(mu-1);           // Thickness of the mica
                                sheet , cm
9
10 printf("\nThe thickness of the mica sheet = %3.1e m"
        , t/1e+02);
11
12 // Result
13 // The thickness of the mica sheet = 2.9e-006 m

```

---

**Scilab code Exa 2.33** Smallest thickness of glass plate for a fringe of minimum intensity

```

1 // Scilab Code Ex2.33: : Page-2.26 (2009)
2 clc; clear;
3 mu = 1.5;          // Refractive index of glass
4 lambda = 5100e-008; // Wavelength of light used , cm
5 i = 30;            // Angle of incidence , degrees
6 n = 1;             // Order of interference fringes
7 // From Snell's law , mu = sind(i)/sind(r) , solving
    for r
8 r = asind(sind(i)/mu); // Angle of refraction ,
    degrees
9 // For a dark fringe in reflection , 2*mu*t*cosd(r) =
    n*lambda, solving for t
10 t = n*lambda/(2*mu*cosd(r));      // Smallest
    thickness of glass plate for a fringe of minimum
    intensity , cm
11 printf("\nThe smallest thickness of glass plate for
        a fringe of minimum intensity = %4.2e cm" , t);
12
13 // Result
14 // The smallest thickness of glass plate for a
    fringe of minimum intensity = 1.80e-005 cm

```

---

**Scilab code Exa 2.34** The wavelength reflected strongly from the soap film

```
1 // Scilab Code Ex2.34:: Page-2.26 (2009)
2 clc; clear;
3 t = 3.1e-05;      // Thickness of the soap film , cm
4 mu = 1.33;        // Refractive index of the soap film
5 r = 0;            // Angle of refraction of the light ray
                     on the soap film , degrees
6 // For bright fringe in reflected pattern ,
7 // 2*mu*t*cosd(r) = (2*n+1)*lambda/2
8 lambda = zeros(3);
9 for n = 1:1:3
10    lambda(n) = 4*mu*t*cosd(r)/(2*(n-1)+1);      //
                     Wavelengths for n = 1, 2 and 3
11    if lambda(n) > 4000e-008 & lambda(n) < 7500e-008
12       then
13          lambda_reflected = lambda(n);
14    end
15
16 printf("\nThe wavelength reflected strongly from the
           soap film = %5.3e cm", lambda_reflected);
17
18 // Result
19 // The wavelength reflected strongly from the soap
           film = 5.497e-05 cm
```

---

**Scilab code Exa 2.35** Order of interference of the dark band

```
1 // Scilab Code Ex2.35:: Page-2.27 (2009)
2 clc; clear;
```

```

3 t = 3.8e-05;      // Thickness of the transparent film
, cm
4 mu = 1.5;         // Refractive index of the
transparent film
5 i = 45;           // Angle of incidence of the light ray
on the transparent film , degrees
6 lambda = 5700e-008; // Wavelength of light , cm
7 // As mu = sind(i)/sind(r) , solving for r
8 r = asind(sind(i)/mu);
9 // For dark fringe in reflected pattern ,
10 // 2*mu*t*cosd(r) = 2*n*lambda , solving for n
11 n = 2*mu*t*cosd(r)/lambda;        // Order of
interference of dark band
12
13 printf("\nThe order of interference of dark band =
%d", ceil(n));
14
15 // Result
16 // The order of interference of dark band = 2
velength reflected strongly from the soap film =
5.497e-05 cm

```

---

**Scilab code Exa 2.36** Absent wavelength of reflected light in the visible spectrum

```

1 // Scilab Code Ex2.36:: Page -2.27 (2009)
2 clc; clear;
3 t = 4.5e-05;      // Thickness of the soap film , cm
4 mu = 1.33;         // Refractive index of the soap film
5 i = 45;           // Angle of incidence of the light ray
on the soap film , degrees
6 // As mu = sind(i)/sind(r) , solving for r
7 r = asind(sind(i)/mu);
8 // For dark fringe in reflected pattern ,
9 // 2*mu*t*cosd(r) = n*lambda , solving for lambda for

```

```

        different n's
10 lambda = zeros(4);
11 for n = 1:1:4
12     lambda(n) = 2*mu*t*cosd(r)/n;      // Wavelengths
        for n = 1, 2, 3 and 4
13     if lambda(n) > 4000e-008 & lambda(n) < 7500e-008
        then
14         lambda_absent = lambda(n);
15     end
16 end
17 printf("\nThe absent wavelength of reflected light
        in the visible spectrum = %4.2e", lambda_absent);
18
19 // Result
20 // The absent wavelength of reflected light in the
        visible spectrum = 5.07e-05

```

---

**Scilab code Exa 2.37** Minimum thickness of the plate that will appear dark in the reflection pattern

```

1 // Scilab Code Ex2.37:: Page-2.28 (2009)
2 clc; clear;
3 mu = 1.6;          // Refractive index of the mica plate
4 r = 60;           // Angle of refraction of the light ray
        on the mica plate , degrees
5 lambda = 5500e-008;    // Wavelength of light used ,
        cm
6 n = 1;   // Order of interference for minimum
        thickness
7 // For dark fringe in reflected pattern ,
8 // 2*mu*t*cosd(r) = 2*n*lambda , solving for t
9 t = n*lambda/(2*mu*cosd(r));    // Minimum thickness
        of the plate that will appear dark in the
        reflection pattern

```

10

```

11 printf("\nThe minimum thickness of the plate that
      will appear dark in the reflection pattern = %4.2
      e cm", t);
12
13 // Result
14 // The minimum thickness of the plate that will
      appear dark in the reflection pattern = 3.44e-05
      cm

```

---

### Scilab code Exa 2.38 Thickness of the thin soap film

```

1 // Scilab Code Ex2.38:: Page-2.28 (2009)
2 clc; clear;
3 mu = 1.33;          // Refractive index of the thin soap
      film
4 lambda1 = 5500e-008;    // Wavelength of the first
      dark fringe , cm
5 lambda2 = 5400e-008;    // Wavelength of the
      consecutive dark fringe , cm
6 i = 30;           // Angle of incidence of the light ray
      on the soap film , degrees
7 // For overlapping fringes ,
8 // n*lambda1 = (n+1)*lambda2 , solving for n
9 n = lambda2/(lambda1-lambda2); // Order of
      interference fringes
10 // As mu = sind(i)/sind(r) , solving for r
11 r = asind(sind(i)/mu);
12 // For dark fringe in reflected pattern ,
13 // 2*mu*t*cosd(r) = 2*n*lambda1 , solving for t
14 t = n*lambda1/(2*mu*cosd(r)); // Thickness of the
      thin soap film
15
16 printf("\nThe thickness of the thin soap film = %5.3
      e cm", t);
17

```

```
18 // Result
19 // The thickness of the thin soap film = 1.205e-03
cm
```

---

**Scilab code Exa 2.39** Order of interference for which light is strongly reflected

```
1 // Scilab Code Ex2.39:: Page-2.29 (2009)
2 clc; clear;
3 t = 0.75e-06;      // Thickness of the glass plate , m
4 mu = 1.5;          // Refractive index of the glass
plate
5 lambda1 = 4000e-010;    // First wavelength of
visible range , cm
6 lambda2 = 7000e-010;    // Last wavelength of
visible range , cm
7 r = 0;              // Angle of refraction for normal
incidence , degrees
8 n = zeros(2);
9 // For bright fringe in reflected pattern ,
10 //  $2*\mu*t*\cos(d(r)) = (2*n+1)*\lambda/2$ , solving for n
11 // For lambda1
12 n(1) = (4*mu*t*cosd(r)/lambda1-1)/2;
13 // For lambda2
14 n(2) = (4*mu*t*cosd(r)/lambda2-1)/2;
15
16 printf("\nFor n = %d and n = %d the light is
strongly reflected .", n(1), ceil(n(2)));
17
18 // Result
19 // For n = 5 and n = 3 the light is strongly
reflected .
```

---

**Scilab code Exa 2.40** Minimum thickness of the film for which light is strongly reflected

```
1 // Scilab Code Ex2.40:: Page-2.30 (2009)
2 clc; clear;
3 mu = 1.45;          // Refractive index of the film
4 lambda = 5500e-010; // First wavelength of
                      visible range , cm
5 r = 0;              // Angle of refraction for normal
                      incidence , degrees
6 n = 0;              // Order of interference is zero for
                      minimum thickness
7 // For bright fringe in reflected pattern ,
8 //  $2*\mu*t*\cos(r) = (2*n+1)*\lambda/2$ , solving for t
9 t = (2*n+1)*lambda/(4*mu*cosd(r)); // Minimum
                      thickness of the film for which light is strongly
                      reflected
10
11 printf("\nThe minimum thickness of the film for
           which light is strongly reflected = %4.2e cm", t)
12 ;
13 // Result
14 // The minimum thickness of the film for which light
   is strongly reflected = 9.48e-08 cm
```

---

**Scilab code Exa 2.41** Thickness of the soap film for dark fringe in reflected pattern

```
1 // Scilab Code Ex2.41:: Page-2.30 (2009)
2 clc; clear;
3 mu = 5/4;          // Refractive index of the film
4 lambda = 5890e-010; // Wavelength of visible
                      light , cm
5 i = 45;            // Angle of incidence , degrees
```

```

6 n = 1;           // Order of interference is unity for
                  minimum thickness in dark reflected pattern
7 // As mu = sind(i)/sind(r), solving for r
8 r = asind(sind(i)/mu);
9 // For dark fringe in reflected pattern ,
10 // 2*mu*t*cosd(r) = n*lambda , solving for t
11 t = n*lambda/(2*mu*cosd(r));      // Thickness of the
                  soap film for dark fringe in reflected pattern
12
13 printf("\nThe thickness of the soap film for dark
      fringe in reflected pattern = %5.3e cm", t);
14
15 // Result
16 // The thickness of the soap film for dark fringe in
      reflected pattern = 2.857e-07 cm

```

---

**Scilab code Exa 2.42** Wavelength in the visible range which is intensified in the reflected beam

```

1 // Scilab Code Ex2.42:: Page -2.30 (2009)
2 clc; clear;
3 mu = 1.5;           // Refractive index of the plate
4 t = 0.5e-006;       // Thickness of the plate , m
5 r = 0;              // Angle of refraction for normal
                  incidence , degrees
6 // For bright fringe in reflected pattern ,
7 // 2*mu*t*cosd(r) = (2*n+1)*lambda/2, solving for
      lambda for different n's
8 lambda = zeros(4);
9 for n = 0:1:3
10     lambda(n+1) = 4*mu*t*cosd(r)/(2*n+1);          //
                  Wavelengths for n = 0, 1, 2 and 3
11     lambda_strong = lambda(n+1);
12     if lambda(n+1) >= 4000e-010 & lambda(n+1) <=
      7500e-010 then

```

```

13         if lambda_strong > lambda(n+1) then // 
14             Search for the stronger wavelength
15             lambda_strong = lambda(n+1);
16         end
17     end
18
19 printf("\nFor n = %d, %4.0f angstrom will be
20 reflected strongly", n, lambda_strong/1e-010);
21 // Result
22 // For n = 3, 4286 angstrom will be reflected
23 // strongly

```

---

**Scilab code Exa 2.43** Thickness of the film with incident white light

```

1 // Scilab Code Ex2.43:: Page -2.31(2009)
2 clc; clear;
3 mu = 1.33;      // Refractive index of the film
4 i = asind(0.8); // Angle of refraction for
5 // normal incidence , degrees
6 r = asind(sind(i)/sind(r)); solving for r
7 lambda1 = 6100e-010; // First wavelength of dark
8 // band , m
9 lambda2 = 6000e-010; // Second wavelength of dark
10 // band , m
11 // For consecutive overlapping wavelenghts
12 // n*lambda1 = (n+1)*lambda2 , solving for n
13 n = lambda2/(lambda1-lambda2);
14 // For dark fringe in reflected pattern ,
15 // 2*mu*t*cosd(r) = n*lambda1, solving for t
16 t = n*lambda1/(2*mu*cosd(r)); // Thickness of the
17 // film with incident white light . m
18 printf("\nThickness of the film with incident white
19

```

```

    light = %3.1e m" , t);
16
17 // Result
18 // Thickness of the film with incident white light =
    1.7e-05 m

```

---

**Scilab code Exa 2.44** Thickness of the film with parallel beam of yellow light

```

1 // Scilab Code Ex2.44:: Page -2.31(2009)
2 clc; clear;
3 mu = 1.5;          // Refractive index of the film
4 i = 45;           // Angle of incidence , degrees
5 // As mu = sind(i)/sind(r) , solving for r
6 r = asind(sind(i)/mu);
7 lambda = 5500e-010; // Wavelength of parallel
    beam of light , m
8 n = 15;           // Order of dark band
9 // For dark fringe in reflected pattern ,
10 // 2*mu*t*cosd(r) = n*lambda , solving for t
11 t = n*lambda/(2*mu*cosd(r)); // Thickness of the
    film with incident parallel beam of light . m
12
13 printf("\nThe thickness of the film with paralle
    beam of yellow light = %4.2e m" , t);
14
15 // Result
16 // The thickness of the film with paralle beam of
    yellow light = 3.12e-06 m

```

---

**Scilab code Exa 2.46** Refractive index of oil

```

1 // Scilab Code Ex2.46:: Page -2.33(2009)

```

```

2 clc; clear;
3 V = 0.58e-006; // Volume of oil , metre cube
4 A = 2.5; // Area of water surface , metre
            square
5 t = V/A; // Thickness of film , m
6 r = 0; // Angle of refraction for normal
          incidence , degrees
7 n = 1; // Order of interference for minimum
          thickness
8 lambda = 4700e-010; // Wavelength of light used ,
           m
9 // For dark fringe in reflected pattern ,
10 //  $2*\mu*t*\cosd(r) = n*\lambda$  , solving for mu
11 mu = n*lambda/(2*t*cosd(r)); // Refractive index
           of oil
12
13 printf("\nThe refractive index of oil = %5.3f", mu);
14
15 // Result
16 // The refractive index of oil = 1.013

```

---

**Scilab code Exa 2.47** Thickness of the soap film to produce constructive interference during reflection

```

1 // Scilab Code Ex2.47:: Page - 2.33(2009)
2 clc; clear;
3 mu = 1.46; // Refractive index of the soap film
4 lambda = 6000e-010; // Wavelength of light used ,
           m
5 r = 0; // Angle of refraction for normal
          incidence , degrees
6 n = 0; // Order of interference for minimum
          thickness
7 // For bright fringe in reflected pattern ,
8 //  $2*\mu*t*\cosd(r) = (2*n+1)*\lambda/2$  , solving for mu

```

```

9 t = (2*n+1)*lambda/(4*mu*cosd(r)); // Thickness of
   soap film , m
10
11 printf("\nThe thickness of soap film = %5.3e m" , t);
12
13 // Result
14 // The thickness of soap film = 1.027e-07 m

```

---

**Scilab code Exa 2.48** Wavelength of light falling on wedge shaped film

```

1 // Scilab Code Ex2.48: : Page - 2.35(2009)
2 clc; clear;
3 mu = 1.4;          // Refractive index of the film
4 alpha = 1.07e-004; // Acute angle of the wedge,
   radian
5 b = 0.2;           // Fringe width , cm
6 // As b = lambda/(2*mu*alpha) , solving for lambda
7 lambda = 2*mu*alpha*b; // Wavelength of light
   falling on wedge shaped film , m
8
9 printf("\nThe wavelength of light falling on wedge
   shaped film = %4d ansgtrom" , lambda/1e-008);
10
11 // Result
12 // The wavelength of light falling on wedge shaped
   film = 5991 ansgtrom

```

---

**Scilab code Exa 2.49** Difference between the thicknesses of the films

```

1 // Scilab Code Ex2.49:: Page - 2.35(2009)
2 clc; clear;
3 mu = 1.4;          // Refractive index of the film
4 lambda = 5500e-008; // Wavelength of the light , cm

```

```

5 // As alpha = (delta_t)/x and x = 10*b; b = lambda
   /(2*mu*alpha), solving for dt
6 delta_t = 10*lambda/(2*mu);      // Difference
   between the thicknesses of the films , cm
7
8 printf("\nDifference between the thicknesses of the
   films = %4.2e cm", delta_t);
9
10 // Result
11 // Difference between the thicknesses of the films =
   1.96e-04 cm

```

---

**Scilab code Exa 2.50** Angle of thin wedge shaped film

```

1 // Scilab Code Ex2.50:: Page -2.36(2009)
2 clc; clear;
3 mu = 1.6;          // Refractive index of the film
4 lambda = 5500e-008; // Wavelength of the light , cm
5 b = 0.1;           // Fringe width , cm
6 // As b = lambda/(2*mu*alpha), solving for alpha
7 alpha = lambda/(2*mu*b);      // Angle of thin wedge
   shaped film , radian
8 printf("\nAngle of thin wedge shaped film = %3.1e
   radian", alpha);
9
10 // Result
11 // Angle of thin wedge shaped film = 1.7e-04 radian

```

---

**Scilab code Exa 2.51** Wavelength of light used to illuminate a wedge shaped film

```

1 // Scilab Code Ex2.51:: Page -2.36(2009)
2 clc; clear;

```

```

3 mu = 1.5;           // Refractive index of the film
4 b = 0.20;            // Fringe width, cm
5 theta = 25/(60*60)*%pi/180;      // Angle of the
       wedge, radian
6 // As b = lambda/(2*mu*theta), solving for lambda
7 lambda = 2*mu*b*theta;        // Wavelength of light
       used to illuminate a wedge shaped film, cm
8
9 printf("\nThe wavelength of light used to illuminate
       a wedge shaped film = %4d angstrom", lambda/1e
       -008);
10
11 // Result
12 // The wavelength of light used to illuminate a
       wedge shaped film = 7272 angstrom
13 // The answer is given wrong in the textbook

```

---

**Scilab code Exa 2.52** Thickness of the wire separating two glass surfaces

```

1 // Scilab Code Ex2.52:: Page – 2.36(2009)
2 clc; clear;
3 lambda = 5893e-010;        // Wavelength of light used ,
       m
4 mu = 1;           // Refractive index of the glass
5 b = 1;            // Assume fringe width to be unity , cm
6 // As b = 1/20, solving for l
7 l = b*20;          // Length of the film , m
8 // As b = lambda/(2*mu*theta) and theta = t/l ,
       solving for t
9 t = lambda*l/(2*mu);    // Thickness of the wire
       separating two glass surfaces , m
10
11 printf("\nThe thickness of the wire separating two
       glass surfaces = %4.2e m" , t);
12

```

```
13 // Result  
14 // The thickness of the wire separating two glass  
    surfaces = 5.89e-06 m
```

---

### Scilab code Exa 2.53 Angle of the wedge shaped air film

```
1 // Scilab Code Ex2.53:: Page - 2.37(2009)  
2 clc; clear;  
3 mu = 1;          // Refractive index of the air film  
4 b = 1.5/25;      // Fringe width , cm  
5 lambda = 5893e-008; // Wavelength of light used  
                      to illuminate a wedge shaped film , cm  
6 // As b = lambda/(2*mu*theta) , solving for theta  
7 theta = lambda/(2*mu*b); // Angle of the wedge ,  
                           radian  
8  
9 printf("\nThe angle of the wedge shaped air film =  
      %5.3f degrees", theta*180/%pi);  
10  
11 // Result  
12 // The angle of the wedge shaped air film = 0.028  
    degrees
```

---

### Scilab code Exa 2.54 Acute angle of the wedge shaped film

```
1 // Scilab Code Ex2.54:: Page - 2.37(2009)  
2 clc; clear;  
3 mu = 1.45;        // Refractive index of the film  
4 b = 1/10;         // Fringe width , cm  
5 lambda = 6600e-008; // Wavelength of light used  
                      to illuminate a wedge shaped film , cm  
6 // As b = lambda/(2*mu*theta) , solving for theta
```

```

7 theta = lambda/(2*mu*b);      // Angle of the wedge ,
radian
8
9 printf("\nThe acute angle of the wedge shaped film =
%6.4f degrees", theta*180/pi);
10
11 // Result
12 // The acute angle of the wedge shaped film = 0.0130
degrees

```

---

**Scilab code Exa 2.55** Diameter of nth dark ring due to first wavelength

```

1 // Scilab Code Ex2.55:: Page - 2.46(2009)
2 clc; clear;
3 lambda1 = 6000e-008;      // First visible wavelength ,
cm
4 lambda2 = 4500e-008;      // Second visible wavelength
, cm
5 R = 100;                  // Radius of curvature of the lens ,
cm
6 // As diameter of nth dark ring due to lambda1 is
7 // D_n^2 = 4*n*R*lambda1 and D_nplus1^2 = 4*(n+1)*R*
lambda2 , so that D_n^2 = D_nplus1^2 gives
8 n = lambda2/(lambda1-lambda2);      // Order of
interference for dark fringes
9 D_n = sqrt(4*n*R*lambda1);      // Diameter of nth
dark ring due to lambda1
10
11 printf("\nThe diameter of nth dark ring due to
wavelength of %4d angstrom = %4.2f cm", lambda1/1
e-008, D_n);
12
13 // Result
14 // The diameter of nth dark ring due to wavelength
of 6000 angstrom = 0.27 cm

```

---

**Scilab code Exa 2.56** Diameter of fifteenth dark ring

```
1 // Scilab Code Ex2.56:: Page -2.46(2009)
2 clc; clear;
3 R = 1;          // For simplicity assume radius of
                  curvature of the lens to be unity , cm
4 D_n = 0.251;    // Diameter of 3rd dark ring , cm
5 D_nplusp = 0.548;   // Diameter of 9th dark ring ,
                  cm
6 n = 3;          // Order of 3rd Newton ring
7 p = 9 - n;      // Order of 6th Newton ring from 3rd
                  ring
8 // As D_nplusp^2 - D_n^2 = 4*p*R*lambda , solving for
                  lambda
9 lambda = (D_nplusp^2 - D_n^2)/(4*p*R);           //
                  Wavelength of light used
10 D_15 = sqrt(D_n^2+4*(15-n)*lambda*R);           //
                  Diameter of 15th dark ring , cm
11
12 printf("\nThe diameter of 15th dark ring = %5.3f cm"
       , D_15);
13
14 // Result
15 // The diameter of 15th dark ring = 0.733 cm
```

---

**Scilab code Exa 2.57** Order of a dark ring having thrice the diameter of the thirtieth ring

```
1 // Scilab Code Ex2.57: : Page -2.47(2009)
2 clc; clear;
3 R = 1;          // For simplicity assume radius of
                  curvature of the lens to be unity , cm
```

```

4 n = 30;           // Order of 3rd Newton ring
5 D_30 = 1;         // Assume diameter of thirtieth ring to
                   be unity , cm
6 // As D_30^2 = 4*n*R*lambda , solving for lambda
7 lambda = D_30^2/(4*n*R);      // Wavelength of light
                   used , cm
8 D_n = 3*D_30;          // Diameter of nth dark ring
                   having thrice the diameter of the thirtieth ring ,
                   cm
9 n = D_n^2/(4*R*lambda);    // Order of a dark ring
                   having thrice the diameter of the thirtieth ring
10
11 printf("\nThe order of the dark ring having thrice
       the diameter of the thirtieth ring = %3d", n);
12
13 // Result
14 // The order of the dark ring having thrice the
       diameter of the thirtieth ring = 270

```

---

**Scilab code Exa 2.58** Radius of curvature of lens and thickness of air film

```

1 // Scilab Code Ex2.58:: Page - 2.47(2009)
2 clc; clear;
3 n = 15;           // Order of 15rd Newton ring
4 D_15 = 0.75;      // Diameter of fifteenth dark ring ,
                   cm
5 lambda = 5890e-008;     // Wavelength of light used ,
                   cm
6 // As D_15^2 = 4*15*R*lambda , solving for R
7 R = D_15^2/(4*15*lambda);    // Radius of curvature
                   of lens , cm
8 // For dark ring , 2*t = n*lambda , solving for t
9 t = n*lambda/2;        // Thickness of air film , cm
10
11 printf("\nThe radius of curvature of lens = %5.1f cm

```

```

    " , R);
12 printf("\nThe thickness of air film = %3.1e cm" , t);
13
14 // Result
15 // The radius of curvature of lens = 159.2 cm
16 // The thickness of air film = 4.4e-004 cm

```

---

### Scilab code Exa 2.59 Refractive index of the liquid

```

1 // Scilab Code Ex2.59:: Page - 2.47(2009)
2 clc; clear;
3 D_15 = 1.62;           // Diameter of 15th dark ring
                        // with air film , cm
4 D_15_prime = 1.47;      // Diameter of 15th dark
                        // ring with liquid , cm
5 R = 1;                 // For simplicity assume radius of
                        // curvature to be unity , cm
6 n = 15;                // Order of 15rd Newton ring
7 // As for ring with air film , D_15^2 = 4*15*R*lambda
                        // , solving for lambda
8 lambda = D_15^2/(4*15*R);     // Wavelength of light
                                // used , cm
9 // As for ring with liquid , D_15_prime^2 = 4*15*R*
                                // lambda/mu, solving for mu
10 mu = 4*15*R*lambda/D_15_prime^2;      //
                                // Refractive index of the liquid
11 printf("\nThe refractive index of the liquid = %4.2f
        " , mu)
12
13 // Result
14 // The refractive index of the liquid = 1.21

```

---

**Scilab code Exa 2.60** Wavelength of light used in Newton rings experiment

```
1 // Scilab Code Ex2.60:: Page - 2.48(2009)
2 clc; clear;
3 D_10 = 0.48;           // Diameter of 10th dark ring
                        with air film , cm
4 D_3 = 0.291;          // Diameter of 3rd dark ring
                        with air film , cm
5 p = 7;                // Order of the 10th ring next to the 3
                        rd ring
6 R = 90;               // Radius of curvature of the lens , cm
7 lambda = (D_10^2-D_3^2)/(4*p*R);      // Wavelength
                        of light used in Newton rings experiment
8
9 printf("\nThe wavelength of light used in Newton
        rings experiment = %4d angstrom", lambda/1e-008);
10
11 // Result
12 // The wavelength of light used in Newton rings
        experiment = 5782 angstrom
```

---

**Scilab code Exa 2.61** Diameter of fifteenth bright ring

```
1 // Scilab Code Ex2.61:: Page - 2.48(2009)
2 clc; clear;
3 R1 = 200;              // Radius of curvature of the convex
                        surface , cm
4 R2 = 250;              // Radius of curvature of the
                        concave surface , cm
5 lambda = 5500e-008;    // Wavelength of light used , cm
6 n = 15;                // Order of interfernce Newton ring
7 // As r_n^2*(1/R1-1/R2) = (2*n-1)*lambda/2, solving
        for r_n
8 r_n = sqrt((2*n-1)*lambda/(2*(1/R1-1/R2)));      //
```

```

    Radius of nth ring , cm
9 D_15 = 2*r_n;           // Daimeter of 15th bright ring ,
                           cm
10
11 printf("\nThe daimeter of 15th bright ring = %4.2f
          cm" , D_15);
12
13 // Result
14 // The daimeter of 15th bright ring = 1.79 cm

```

---

**Scilab code Exa 2.62** Wavelength of light used in Newton rings experiment

```

1 // Scilab Code Ex2.62:: Page - 2.49(2009)
2 clc; clear;
3 R = 80;           // Radius of curvature of the convex
                   surface , cm
4 D5 = 0.192;      // Diameter of 5th dark ring , cm
5 D25 = 0.555;     // Diameter of 25th dark ring , cm
6 n = 5;           // Order of interfearence Newton ring
7 P = 25 - n;
8 lambda = (D25^2 - D5^2)/(4*P*R);      // Wavelength of
                                           light used , cm
9 printf("\nThe wavelength of light used = %5.3e cm" ,
        lambda);
10
11 // Result
12 // The wavelength of light used = 4.237e-005 cm
13 // The expression for lambda is given wrong in the
   textbook but solved correctly

```

---

**Scilab code Exa 2.63** Diameter of fifteenth dark Newton ring

```

1 // Scilab Code Ex2.63:: Page -2.49(2009)
2 clc; clear;
3 R1 = 4;           // Radius of curvature of the convex
                   surface , m
4 R2 = 5;           // Radius of curvature of the concave
                   surface , m
5 lambda = 6600e-010; // Wavelength of light used , cm
6 n = 15;          // Order of Newton ring
7 // As D_n^2*(1/R1-1/R2) = 4*n*lambda , solving for
                   D_n
8 D_15 = sqrt(4*n*lambda/(1/R1-1/R2));      // Diameter
                   of 15th dark ring , cm
9
10 printf("\nThe diameter of %dth dark ring = %4.2e m" ,
        n, D_15);
11
12 // Result
13 // The diameter of 15th dark ring = 2.81e-002 m
14 // The answer is given wrong in the textbook (the
                   square root is not solved)

```

---

**Scilab code Exa 2.64** Diameter of fifteenth dark ring due to first wavelength

```

1 // Scilab Code Ex2.64:: Page -2.49(2009)
2 clc; clear;
3 lambda1 = 6000e-008;      // First visible wavelength ,
                           cm
4 lambda2 = 4500e-008;      // Second visible wavelength
                           , cm
5 R = 120;                  // Radius of curvature of the lens ,
                           cm
6 // As diameter of nth dark ring due to lambda1 is
7 // D_n^2 = 4*n*R*lambda1 and D_nplus1^2 = 4*(n+1)*R*
                           lambda2 , so that D_n^2 = D_nplus1^2 gives

```

```

8 n = lambda2/(lambda1-lambda2);           // Order of
    interference for dark fringes
9 printf("\nThe value of n = %d", n);
10 n = 15;        // Order of interference fringe
11 D_n = sqrt(4*n*R*lambda1);           // Diameter of nth
    dark ring due to lambda1
12 printf("\nThe diameter of 15th dark ring due to
    wavelength of %4d angstrom = %4.2f cm", lambda1/1
e-008, D_n);
13
14 // Result
15 // The value of n = 3
16 // The diameter of 15th dark ring due to wavelength
    of 6000 angstrom = 0.66 cm

```

---

**Scilab code Exa 2.65** Refractive index of the liquid filled into container

```

1 // Scilab Code Ex2.65:: Page - 2.49 (2009)
2 clc; clear;
3 lambda = 5896e-008;      // Wavelength of light used ,
    cm
4 R = 100;                // Radius of curvature of the lens ,
    cm
5 D10 = 0.4;              // Diametre of 10th dark ring , cm
6 n = 10;                 // Order of Newton ring
7 // As for a dark ring ,  $2*\mu*t = n*\lambda$  and  $2*t = (D_{10}/2)^2/R$ , solving for mu
8 mu = 4*n*lambda*R/D10^2; // Refractive index of
    the liquid filled into container
9
10 printf("\nThe refractive index of the liquid filled
    into container = %4.2f", mu);
11
12 // Result
13 // The refractive index of the liquid filled into

```

```
container = 1.47
```

---

### Scilab code Exa 2.67 Refractive index of the liquid

```
1 // Scilab Code Ex2.67:: Page -2.50(2009)
2 clc; clear;
3 Dn = 1.8;           // Diameter of 15th dark ring , cm
4 Dn_prime = 1.67;    // Diameter of 15th dark ring
                      // with liquid , cm
5 mu = (Dn/Dn_prime)^2; // Refractive index of the
                          // liquid
6
7 printf("\nThe refractive index of the liquid = %4.2f
          ", mu);
8
9 // Result
10 // The refractive index of the liquid = 1.16
```

---

### Scilab code Exa 2.68 Diameter of eighteenth dark ring

```
1 // Scilab Code Ex2.68:: Page -2.51(2009)
2 clc; clear;
3 R = 1;           // For simplicity assume radius of
                  // curvature to be unity , cm
4 D8 = 0.45;        // Diameter of 8th dark ring , cm
5 D15 = 0.81;        // Diameter of 15th dark ring , cm
6 n = 8;           // Order of 8th Newton ring
7 p = 7;           // Order of 7th Newton ring after 8
                  // th ring
8 lambda = (D15^2-D8^2)/(4*p*R);      // Wavelength of
                  // light used , cm
9 // As D18^2-D15^2 = 4*p*lambda*R
10 p = 3;          // For 18th and 15th rings
```

```

11 D18 = sqrt(D15^2+4*p*lambda*R);           // Diameter of
18th ring , cm
12
13 printf("\nThe diameter of 18th dark ring = %6.4f cm"
14 , D18);
15 // Result
16 // The diameter of 18th dark ring = 0.9222 cm

```

---

**Scilab code Exa 2.69** Wavelength of light used to illuminate plano convex lens in Newton rings experiment

```

1 // Scilab Code Ex2.69:: Page -2.51(2009)
2 clc; clear;
3 R = 100;          // Radius of curvature of plano-convex
lens , cm
4 D15 = 0.590;      // Diameter of 15th dark ring , cm
5 D5 = 0.336;       // Diameter of 5th dark ring , cm
6 p = 10;           // Order of 10th Newton ring after
5th ring
7 lambda = (D15^2-D5^2)/(4*p*R);           // Wavelength of
light used , cm
8
9 printf("\nThe wavelength of light used = %4.0f
angstrom", lambda/1e-008);
10
11 // Result
12 // The wavelength of light used = 5880 angstrom

```

---

**Scilab code Exa 2.70** Wavelength of monochromatic light used in Michelson Interferometer

```
1 // Scilab Code Ex2.70:: Page -2.57(2009)
```

```

2 clc; clear;
3 N = 250;           // Number of fringes crossing the
        field of view
4 delta_x = 0.0595e-01;    // Displacement in movable
        mirror , cm
5 // As N*lambda/2 = delta_x , solving for lambda
6 lambda = 2*delta_x/N;      // Wavelength of light
        used , cm
7
8 printf("\nThe wavelength of monochromatic light used
        = %4.0f ansgtrom", lambda/1e-008);
9
10 // Result
11 // The wavelength of monochromatic light used = 4760
        ansgtrom
12 // Answer is given wrong in the textbook

```

---

**Scilab code Exa 2.71** Number of fringes that passes across the cross wire of telescope

```

1 // Scilab Code Ex2.71:: Page -2.58(2009)
2 clc; clear;
3 delta_x = 0.02559e-01;    // Displacement in movable
        mirror , cm
4 lambda = 5890e-008;      // Wavelength of light used
        , cm
5 // As N*lambda/2 = delta_x , solving for N
6 N = 2*delta_x/lambda;      // Number of fringes
        crossing the field of view
7
8 printf("\nThe number of fringes that passes across
        the cross wire of telescope = %2d", ceil(N));
9
10 // Result
11 // The number of fringes that passes across the

```

cross wire of telescope = 87

---

**Scilab code Exa 2.72** Distance between two successive positions of movable mirror

```
1 // Scilab Code Ex2.72:: Page - 2.58(2009)
2 clc; clear;
3 lambda1 = 5890e-008;      // Wavelength corresponding
   to the D1 line , cm
4 lambda2 = 5896e-008;      // Wavelength corresponding
   to the D2 line , cm
5 delta_lambda = lambda2 - lambda1;    // Difference in
   the wavelengths , cm
6 // As delta_lambda = lambda1*lambda2/(2*x) , solving
   for x
7 x = lambda1*lambda2/(2*(lambda2-lambda1));    //
   Distance between two successive positions of
   movable mirror
8
9 printf("\nThe distance between two successive
   positions of movable mirror = %3.1e cm", x);
10
11 // Result
12 // The distance between two successive positions of
   movable mirror = 2.9e-002
```

---

**Scilab code Exa 2.73** Thickness of the transparent glass film

```
1 // Scilab Code Ex2.73:: Page - 2.58(2009)
2 clc; clear;
3 N = 550;           // Number of fringes crossing the
   field of view
```

```
4 lambda = 5500e-008;      // Wavelength of light used ,  
    cm  
5 mu = 1.5;              // Refractive index of the glass  
    slab  
6 // As 2*(mu-1)*t = N*lambda , solving for t  
7 t = N*lambda/(2*(mu-1));           // Thickness of the  
    transparent glass film  
8  
9 printf("\nThe distance between two successive  
    positions of movable mirror = %3.1e cm" , t);  
10  
11 // Result  
12 // The distance between two successive positions of  
    movable mirror = 3.0e-002 cm
```

---

# Chapter 3

## Diffraction

**Scilab code Exa 3.1** Position of the screen so that light is focused on the brightest spot

```
1 // Scilab Code Ex3.1:: Page-3.9 (2009)
2 clc; clear;
3 lambda = 5890e-008; // Wavelength of light used , cm
4 r1 = 0.2; // Radius of first ring of zone plate ,
cm
5 n = 1; // Order of zone plate
6 f1 = r1^2/(n*lambda); // Position of the screen so
that light is focused on the brightest spot , cm
7
8 printf("\nThe position of the screen so that light
is focused on the brightest spot = %3.1e cm",
lambda);
9
10 // Result
11 // The position of the screen so that light is
focused on the brightest spot = 5.9e-005 cm
```

---

**Scilab code Exa 3.2** Zone plate with a point source of light on the axis

```

1 // Scilab Code Ex3.2:: Page-3.9 (2009)
2 clc; clear;
3 v1 = 36;      // Position of the strongest image from
               the zone plate , cm
4 v2 = 9;       // Position of the next image from the
               zone plate , cm
5 lambda = 5890e-008; // Wavelength of light used , cm
6 r1 = 1;       // For simplicity assume radius of first
               ring of zone plate to be unity , cm
7 n = 1;        // Order of zone plate
8 // As 1/v1-1/u = n*lambda/r1^2 = 1/3*(1/v2-1/u) ,
               solving for u
9 u = 2/(3/36-1/9);      // Distance of the zone
               plate from source , cm
10 // As 1/v-1/u = n*lambda/r1^2, solving for r1
11 r1 = sqrt(lambda/(1/v1-1/abs(u)));           // Radius
               of first zone , cm
12 f1 = r1^2/(n*lambda);    // Principal focal length ,
               cm
13
14 printf("\nThe distance of the zone plate from source
         = %2d cm", u);
15 printf("\nThe radius of first zone = %3.1e cm", r1);
16 printf("\nThe principal focal length = %4.1f cm", f1
         );
17
18 // Result
19 // The distance of the zone plate from source = -72
               cm
20 // The radius of first zone = 6.5e-002 cm
21 // The principal focal length = 72.0 cm

```

---

**Scilab code Exa 3.3** Position of the first image in a zone plate

```
1 // Scilab Code Ex3.3:: Page-3.10 (2009)
```

```

2 clc; clear;
3 lambda = 5500e-010; // Wavelength of light used , cm
4 u = -4; // Distance of the zone plate from
           source , cm
5 D = 3.7e-003; // Diameter of central zone of zone
                  plate , cm
6 r = D/2; // Radius of central zone of zone plate ,
           cm
7 n = 1; // Order of zone plate
8 f1 = r^2/(n*lambda); // Principal focal
           length , cm
9 v1 = 36; // Position of the strongest image from
           the zone plate , cm
10 v2 = 9; // Position of the next image from the
           zone plate , cm
11 // As 1/v - 1/u = 1/f , solving for v
12 v = 1/(1/f1+1/u); // Position of the first
           image in a zone plate , cm
13
14 printf("\nThe position of the first image in a zone
           plate = %2d cm", floor(v));
15
16 // Result
17 // The position of the first image in a zone plate =
           -12 cm

```

---

### Scilab code Exa 3.4 Principal focal length of zone plate

```

1 // Scilab Code Ex3.4:: Page-3.11 (2009)
2 clc; clear;
3 lambda = 1; // For simplicity assume wavelength
              of light used to be unity , unit
4 R = 150; // Radius of curvature of the curved
           surface , cm
5 r1 = sqrt(lambda*R); // For Newton's ring , cm

```

```

6 f1 = r1^2/lambda;           // Principal focal length of
    zone plate , cm
7
8 printf("\nThe principal focal length of zone plate =
    %3d cm", f1);
9
10 // Result
11 // The principal focal length of zone plate = 150 cm

```

---

**Scilab code Exa 3.5** Half angular width at central maximum in Fraunhofer diffraction

```

1 // Scilab Code Ex3.5:: Page -3.22 (2009)
2 clc; clear;
3 lambda = 5000e-008;      // Wavelength of light used ,
    cm
4 a = 15e-005;      // Width of the slit , cm
5 n = 1;          // Order of diffraction
6 // For a single slit Fraunhofer diffraction , a*sin(
    theta) = n*lambda, solving for theta
7 theta = asin(n*lambda/a); // Half angular width at
    central maximum in Fraunhoffer diffraction ,
    radian
8
9 printf("\nThe half angular width at central maximum
    in Fraunhoffer diffraction = %5.3f rad", theta);
10
11 // Result
12 // The half angular width at central maximum in
    Fraunhoffer diffraction= 0.340 rad

```

---

**Scilab code Exa 3.6** Width of the slit

```

1 // Scilab Code Ex3.6:: Page -3.23 (2009)
2 clc; clear;
3 lambda = 5000e-010;      // Wavelength of light used ,
   cm
4 n = 1;      // Order of diffraction
5 x = 5e-003;      // Position of first minima on
   either sides of central maximum, m
6 D = 2.5;      // Distance of screen from the narrow
   slir , m
7 sin_theta = x/sqrt(x^2+D^2);      // Sine of angle
   theta , rad
8 // For a single slit Fraunhofer diffraction , a*sin(
   theta) = n*lambda , solving for a
9 a = n*lambda/sin_theta;      // Width of the slit , m
10
11 printf("\nThe Width of the slit = %3.1e m", a);
12
13 // Result
14 // The Width of the slit = 2.5e-004 m

```

---

### Scilab code Exa 3.7 Angular width of central maximum

```

1 // Scilab Code Ex3.7:: Page -3.23 (2009)
2 clc; clear;
3 lambda = 6000e-010;      // Wavelength of light used ,
   m
4 a = 15e-007;      // Width of the slit , m
5 // For a single slit Fraunhofer diffraction , a*sind(
   theta) = n*lambda , solving for theta
6 theta = asind(lambda/a); // Half angular width of
   central maximum, degrees
7
8 printf("\nThe angular width of central maximum = %2d
   degrees", 2*ceil(theta));
9

```

```
10 // Result  
11 // The angular width of central maximum = 48 degrees
```

---

**Scilab code Exa 3.8** Distance between first minima and the next minima from the axis

```
1 // Scilab Code Ex3.8:: Page-3.23 (2009)  
2 clc; clear;  
3 lambda = 5000e-010;      // Wavelength of light used ,  
                           m  
4 a = 0.7e-002;           // Width of the slit , m  
5 f = 0.5;                // Focal length of the lens , m  
6 n = 1;                  // Order of diffraction  
7 // For minima , a*sind(theta_n) = n*lambda  
8 // Also theta_n = n*lambda/a = x1/f , solving for x1  
9 x1 = f*n*lambda/a;      // Position of first  
                           minima , cm  
10 // For secondary maxima , a*sind(theta_n) = (2*n+1)*  
    lambda/2  
11 // Also theta_n = 3*lambda/(2*a) = x2/f , solving for  
    x2  
12 n = 1;                  // Order of diffraction for first  
                           secondary minima  
13 x2 = 3*f*lambda/(2*a); // Position of first  
                           secondary maxima , cm  
14  
15 printf("\nThe distance between first minima and the  
        next minima from the axis = %4.2e cm", x2-x1);  
16  
17 // Result  
18 // The distance between first minima and the next  
    minima from the axis = 1.79e-005 cm
```

---

**Scilab code Exa 3.9** Width of central maxima in diffraction pattern

```
1 // Scilab Code Ex3.9:: Page-3.24 (2009)
2 clc; clear;
3 lambda = 6600e-008;      // Wavelength of light used ,
   cm
4 a = 0.018;      // Width of the slit , cm
5 f = 200;        // Focal length of the lens , cm
6 n = 1;          // Order for first order diffraction
7 // As a*sin(theta) = n*lambda, a*theta = n*lambda
8 // As theta = lambda/a and theta = x/f, solving for
   x
9 x = lambda*f/a;    // Half angular width at central
   maximum , cm
10
11 printf("\nThe width of central maximum = %3.1f cm" ,
   2*x);
12
13 // Result
14 // The width of central maximum = 1.5 cm
```

---

**Scilab code Exa 3.10** Slit width in Fraunhoffer single slit experiment

```
1 // Scilab Code Ex3.10:: Page-3.24 (2009)
2 clc; clear;
3 f = 250;        // Focal length of the lens , cm
4 x = 0.8;        // Half width of central maxima, cm
5 lambda = 5500e-008;    // Wavelength of light used ,
   cm
6 // As x = f*lambda/a, solving for a
7 a = f*lambda/x;    // Slit width in Fraunhofer
   single slit experiment
8
9 printf("\nThe slit width = %5.3f cm" , a);
10
```

```
11 // Result  
12 // The slit width = 0.017 cm
```

---

### Scilab code Exa 3.11 Half angular width of central maxima

```
1 // Scilab Code Ex3.11:: Page-3.25 (2009)  
2 clc; clear;  
3 lambda = 5500e-008;      // Wavelength of light used ,  
                           cm  
4 a = 8.5e-005;      // Width of the slit , cm  
5 n = 1;          // Order of diffraction  
6 // For a single slit Fraunhofer diffraction , a*sind(  
      theta) = n*lambda, solving for theta  
7 theta = asind(n*lambda/a); // Half angular width at  
                           central maximum in Fraunhoffer diffraction ,  
                           degrees  
8  
9 printf("\nThe half angular width at central maximum  
       in Fraunhoffer diffraction = %4.1f degrees",  
       theta);  
10  
11 // Result  
12 // The half angular width at central maximum in  
   Fraunhoffer diffraction = 40.3 degrees
```

---

### Scilab code Exa 3.12 Wavelength of light used in Fraunhoffer diffraction due to single slit

```
1 // Scilab Code Ex3.12:: Page-3.25 (2009)  
2 clc; clear;  
3 a = 0.04;          // Slit width , cm  
4 x = 0.5;          // Half width of central maximum , cm  
5 f = 300;          // Focal length of the lens , cm
```

```

6 // As x = lambda*f/a, solving for lambda
7 lambda = a*x/f;      // Wavelength of light used in
   Fraunhoffer diffraction due to single slit , cm
8
9 printf("\nThe wavelength of light used in
   Fraunhoffer diffraction due to a single slit =
   %4d angstrom", lambda/1e-008);
10
11 // Result
12 // The wavelength of light used in Fraunhoffer
   diffraction due to a single slit = 6666 angstrom

```

---

**Scilab code Exa 3.13** Width of central maxima from position of first secondary minima

```

1 // Scilab Code Ex3.13:: Page -3.25 (2009)
2 clc; clear;
3 a = 0.045;           // Slit width , cm
4 lambda = 5500e-008;    // Wavelength of light used ,
   cm
5 f = 250;            // Focal length of the lens , cm
6 x = lambda*f/a;      // Position of central maxima ,
   cm
7
8 printf("\nThe position of central maxima = %5.3f cm"
   , x);
9 printf("\nThe width of central maxima from first
   minima = %5.3f cm" , 2*x);
10
11 // Result
12 // The position of central maxima = 0.306 cm
13 // The width of central maxima from first minima =
   0.611 cm

```

---

**Scilab code Exa 3.14** Wavelength of monochromatic light used in illuminating a slit

```
1 // Scilab Code Ex3.14:: Page-3.26 (2009)
2 clc; clear;
3 a = 0.025;           // Slit width , cm
4 n = 2;               // Order of diffraction
5 f = 400;              // Focal length of the lens , cm
6 x = 2.1;              // Position of central maxima , cm
7 // As theta = n*lambda/a and theta = x/f , solving
   for lambda
8 lambda = x*a/(n*f);    // Wavelength of light used ,
   cm
9 printf("\nThe wavelength of light used = %4d
   angstrom", lambda/1e-008);
10
11 // Result
12 // The wavelength of light used = 6562 angstrom
```

---

**Scilab code Exa 3.15** Distance between second dark and next bright fringe on the axes

```
1 // Scilab Code Ex3.15:: Page-3.26 (2009)
2 clc; clear;
3 a = 0.25;           // Slit width , cm
4 lambda = 5890e-008;    // Wavelength of light , cm
5 f = 80;               // Focal length of the lens , cm
6 n = 2;               // Order of diffraction
7 // As for minima, theta = n*lambda/a and theta = x/f
   , solving for x
8 x2 = 2*lambda*f/a;      // Position of 2nd dark
   fringe , cm
```

```

9 // As for maxima, theta = (2*n+1)*lambda/(2*a) and
   theta = x/f, solving for x
10 x2_prime = 5*lambda*f/(2*a);           // Position of 2
    nd bright fringe , cm
11 delta_x = x2_prime-x2;           // Distance between 2nd
    dark and next bright , cm
12 printf("\nThe distance between 2nd dark and next
    bright fringe = %4.2e cm", delta_x);
13
14 // Result
15 // The distance between 2nd dark and next bright
fringe = 9.42e-003 cm

```

---

**Scilab code Exa 3.16** Width of the slit from first order diffraction

```

1 // Scilab Code Ex3.16:: Page -3.27 (2009)
2 clc; clear;
3 lambda = 5500e-008;           // Wavelength of light used ,
   cm
4 x = 3.9e-001;           // Half width of central maximum
   , cm
5 f = 220;           // Focal length of the lens , cm
6 n = 1;           // Order for first order diffraction
7 // As a*sin(theta) = n*lambda, a*theta = n*lambda
8 // As theta = lambda/a and theta = x/f, solving for
   a
9 a = lambda*f/x;           // Half angular width at central
   maximum , cm
10
11 printf("\nThe width of the slit = %3.1e cm", a);
12
13 // Result
14 // The width of the slit = 3.1e-002 cm

```

---

### Scilab code Exa 3.18 Fraunhoffer diffraction due to double slits

```
1 // Scilab Code Ex3.18:: Page-3.30 (2009)
2 clc; clear;
3 a = 0.019e-003;           // Width of each slit , m
4 b = 2.0e-004;             // Width of opacity between two
                           slits , m
5 lambda = 5000e-010; // Wavelength of light used , m
6 D = 0.6;                  // Distance between slit and the
                           screen , m
7 // As angular separation , theta = x/D = lambda/(a+b)
   , solving for x
8 x = D*lambda/(a+b);      // Fringe spacing on the
                           screen , m
9 // As half angular separation , theta1 = x1/D =
   lambda/(2*(a+b)) , solving for x1
10 x1 = D*lambda/(2*(a+b)); // Distance between
                           central maxima and first minima , m
11
12 printf("\nThe fringe spacing on the screen = %4.2f
   mm" , x/1e-003);
13 printf("\nThe distance between central maxima and
   first minima = %4.2f mm" , x1/1e-003);
14
15 // Result
16 // The fringe spacing on the screen = 1.37 mm
17 // The distance between central maxima and first
   minima = 0.68 mm
```

---

### Scilab code Exa 3.19 Fringe separation in Fraunhoffer double slit diffraction pattern

```

1 // Scilab Code Ex3.19:: Page -3.31 (2009)
2 clc; clear;
3 f = 150;      // Distance between screen and slit , cm
4 a = 0.005;    // Slit width , cm
5 b = 0.06;     // Distance between slits , cm
6 lambda = 5500e-008;      // Wavelength of light used ,
                           cm
7 // As half angular separation , theta1 = x1/f =
   lambda/(2*(a+b)) , solving for x1
8 x1 = f*lambda/(2*(a+b));      // Distance between
                           central maxima and first minima , cm
9 delta_theta = lambda/(2*(a+b)); // Angular
                           separation between two consecutive minima ,
                           radians
10 printf("\nThe distance between central maxima and
           first minima = %4.2e cm", x1);
11 printf("\nThe angular separation between two
           consecutive minima = %3.1e radians", delta_theta)
;
12
13 // Result
14 // The distance between central maxima and first
   minima = 6.35e-002 cm
15 // The angular separation between two consecutive
   minima = 4.2e-004 radians

```

---

**Scilab code Exa 3.20** Positions of first secondary maxima and minima in double slit diffraction

```

1 // Scilab Code Ex3.20:: Page -3.32 (2009)
2 clc; clear;
3 f = 120;      // Distance between screen and slit , cm
4 a = 0.019;    // Slit width , cm
5 b = 0.041;    // Distance between slits , cm
6 lambda = 6500e-008;      // Wavelength of light used ,

```

```

        cm
7 // As theta1 = x1/f = lambda/(2*(a+b)) , solving for
   x1
8 x1 = f*lambda/(2*(a+b));           // Position of first
   secondary minima , cm
9 // As theta2 = x2/f = lambda/(a+b) , solving for x2
10 x2 = f*lambda/(a+b);            // Position of first
    secondary maxima , cm
11
12 printf("\nThe position of first secondary minima =
      %5.3f cm", x1);
13 printf("\nThe position of first secondary maxima =
      %4.2f cm", x2);
14
15 // Result
16 // The position of first secondary minima = 0.065 cm
17 // The position of first secondary maxima = 0.13 cm

```

---

**Scilab code Exa 3.21** Missing orders of spectra in Fraunhofer double slit diffraction

```

1 // Scilab Code Ex3.21:: Page -3.34 (2009)
2 clc; clear;
3 a = 0.2; // Slit width , mm
4 b = 0.8; // Distance between slits , mm
5 p = [1 2 3 4]; // Orders of pth diffraction
   maxima
6 // As diffraction of pth diffraction maxima , a*sin(
   theta)=p*lambda --- (i)
7 // and that of nth diffraction maxima , (a+b)*sin(
   theta)=n*lambda --- (ii)
8 // Dividing (ii) by (i), we have
9 // (a+b)/a = n/p, solving for n
10 n = (a+b)/a*p; // Orders of nth diffraction maxima
11

```

```

12 printf("\nThe missing orders of spectra in
        diffraction maxima, n = %d, %d, %d, %d,...", n(1)
        , n(2), n(3), n(4));
13
14
15 // Result
16 // The missing orders of spectra in diffraction
        maxima, n = 5, 10, 15, 20, ...

```

---

**Scilab code Exa 3.22** Angles of diffraction for the principal maxima for two lines of sodium

```

1 // Scilab Code Ex3.22:: Page-3.45 (2009)
2 clc; clear;
3 lambda1 = 5890e-008;      // Wavelength of D1 line of
    Na, cm
4 lambda2 = 5896e-008;      // Wavelength of D2 line of
    Na, cm
5 N = 3000/0.5;            // No. of lines per cm of
    grating, lines/cm
6 a_plus_b = 1/N;          // Grating element, cm
7 n = 1;                  // Order of diffraction for principal
    maxima
8 // As (a+b)*sin(theta1) = n*lambda, solving for
    theta1
9 theta1 = asind(n*lambda1/(a_plus_b)); // Angle of
    diffraction for the principal maxima of D1 line,
    degrees
10 theta2 = asind(n*lambda2/(a_plus_b)); // Angle of
    diffraction for the principal maxima of D2 line,
    degrees
11 printf("\nThe angle of diffraction for the principal
        maxima of D1 line = %5.2f degrees", theta1);
12 printf("\nThe angle of diffraction for the principal
        maxima of D2 line = %5.2f degrees", theta2);

```

```
13
14 // Result
15 // The angle of diffraction for the principal maxima
   of D1 line = 20.70 degrees
16 // The angle of diffraction for the principal maxima
   of D2 line = 20.72 degrees
```

---

**Scilab code Exa 3.23** Highest order spectrum which can be seen in monochromatic light

```
1 // Scilab Code Ex3.23:: Page -3.45 (2009)
2 clc; clear;
3 lambda = 5500e-008;      // Wavelength of light used ,
   cm
4 N = 15000;                // No. of lines per inch of
   grating , lines/inch
5 a_plus_b = 2.54/N;         // Grating element , cm
6 n = 1;                    // Order of diffraction for principal
   maxima
7 // As (a+b)*sin(theta_n) = n*lambda and for maximum
   possible order of spectra sin(theta_n) = 1
8 // So (a+b) = n*lambda , solving for n
9 n = (a_plus_b)/lambda;    // The highest order
   spectrum which can be seen in monochromatic light
10
11 printf("\nThe highest order spectrum which can be
   seen in monochromatic light = %d", n);
12
13 // Result
14 // The highest order spectrum which can be seen in
   monochromatic light = 3
```

---

**Scilab code Exa 3.24** Angle of separation in second order of diffraction spectrum

```
1 // Scilab Code Ex3.24: : Page-3.46 (2009)
2 clc; clear;
3 lambda1 = 5890e-008;      // Wavelength of D1 line , cm
4 lambda2 = 5896e-008;      // Wavelength of D2 line , cm
5 N = 15000;                // No. of lines per inch of
                           grating , lines/inch
6 a_plus_b = 2.54/N;         // Grating element , cm
7 n = 2;                   // Order of diffraction for secondary
                           maxima
8 // As (a+b)*sin(theta_n) = n*lambda , solving for
                           theta1 and theta2
9 theta1 = asind(n*lambda1/a_plus_b); // Direction
                           of secondary maxima with lambda1 , degrees
10 theta2 = asind(n*lambda2/a_plus_b); // Direction
                           of secondary maxima with lambda2 , degrees
11
12 printf("\nThe angle of separation in second order
                           diffraction spectrum = %3.1f degrees", theta2-
                           theta1);
13
14 // Result
15 // The angle of separation in second order
                           diffraction spectrum = 0.1 degrees
```

---

**Scilab code Exa 3.25** Separation of two lines in first order spectrum

```
1 // Scilab Code Ex3.25:: Page-3.46 (2009)
2 clc; clear;
3 lambda1 = 5500e-008;      // First wavelength , cm
4 lambda2 = 3700e-008;      // Second wavelength , cm
5 N = 15000;                // No. of lines per inch of
                           grating , lines/inch
```

```

6 a_plus_b = 2.54/N;           // Grating element , cm
7 f = 120;                   // Focal length of the lens , cm
8 n = 1;                     // Order of diffraction for principal
                             maxima
9 // As (a+b)*sin(theta_n) = n*lambda , solving for
   theta1 and theta2
10 theta1 = asind(n*lambda1/a_plus_b);    // Direction
   of principal maxima with lambda1 , degrees
11 theta2 = asind(n*lambda2/a_plus_b);    // Direction
   of principal maxima with lambda2 , degrees
12 // As tand(theta) = x/f, solving for x1 - x2 = dx
13 dx = f*(tand(theta1)-tand(theta2));    // Linear
   separation of two lines in first order spectrum ,
   cm
14
15 printf("\nThe linear separation of two lines in
   first order spectrum = %5.2f cm", dx);
16
17 // Result
18 // The linear separation of two lines in first order
   spectrum = 14.34 cm

```

---

**Scilab code Exa 3.26** Difference in the deviation in the first and third order spectra

```

1 // Scilab Code Ex3 .26:: Page –3.47 (2009)
2 clc; clear;
3 lambda = 5000e-008;      // Wavelength of light used ,
   cm
4 N = 5000;                // No. of lines per cm of
   grating , lines/cm
5 a_plus_b = 1/N;           // Grating element , cm
6 n = 1;                   // Order of diffraction for first order
   spectra
7 // As (a+b)*sin(theta_n) = n*lambda , solving for

```

```

    theta for first and third orders
8 theta1 = asind(n*lambda/a_plus_b);      // Direction
      of principal maxima with lambda1, degrees
9 n = 3;        // Order of diffraction for third order
               spectra
10 theta3 = asind(n*lambda/a_plus_b);      // Direction
      of principal maxima with lambda2, degrees
11 delta_theta = theta3 - theta1; // Angular
      separation in the first and third order spectra ,
12
13 printf("\nThe difference in the deviation in the
      first and third order spectra = %4.1f degrees",
      delta_theta);
14
15 // Result
16 // The difference in the deviation in the first and
      third order spectra = 34.1 degrees

```

---

**Scilab code Exa 3.27** Order of diffraction for the given grating element and wavelength of light

```

1 // Scilab Code Ex3.27:: Page-3.48 (2009)
2 clc; clear;
3 lambda = 6500e-008;      // Wavelength of light used ,
cm
4 N = 10000;                // No. of lines per cm of
grating , lines/cm
5 a_plus_b = 1/N;           // Grating element , cm
6 theta_n = 90;             // Direction for maximum possible
orders , degrees
7 // As (a+b)*sin(theta_n) = n*lambda , solving for
theta for n
8 n = a_plus_b*sind(theta_n)/lambda; // Order of
diffraction for
9

```

```

10 printf("\nThe order of diffraction for the given
           grating element and wavelength of light = %d", n)
           ;
11
12 // Result
13 // The order of diffraction for the given grating
           element and wavelength of light = 1

```

---

**Scilab code Exa 3.28** Number of lines ruled on the grating surface

```

1 // Scilab Code Ex3.28:: Page -3.48 (2009)
2 clc; clear;
3 lambda1 = 6500e-008;      // Wavelength of first line ,
           cm
4 lambda2 = 4500e-008;      // Wavelength of second
           line , cm
5 theta1 = 18;              // Direction of lower order ,
           degrees
6 theta2 = 18;              // Direction of higher order ,
           degrees
7 // As (a+b)*sin(theta1) = n*lambda1 and (a+b)*sin(
           theta2) = (n+1)*lambda2, solving for n
8 n = lambda2/(lambda1 - lambda2);    // Order of
           diffraction for first wavelength
9 // As a_plus_b = n*lambda1/sind(theta1), solving for
           a_plus_b
10 a_plus_b = ceil(n)*lambda1/sind(theta1); // Grating
           element , cm
11 N = 1/a_plus_b;      // No. of lines on the grating
           surface , lines/cm
12
13 printf("\nThe number of lines ruled on the grating
           surface = %4d lines/cm", N);
14
15 // Result

```

```
16 // The number of lines ruled on the grating surface  
= 1584 lines/cm
```

---

**Scilab code Exa 3.29** Angles at which first and second order maxima are observed

```
1 // Scilab Code Ex3.29:: Page-3.48 (2009)  
2 clc; clear;  
3 lambda = 6328e-008;      // Wavelength of He-Laser , cm  
4 a_plus_b = 1/6000;        // Grating element , cm  
5 n = 1;                  // First order of diffraction for given  
                           wavelength  
6 // As (a+b)*sin(theta1) = n*lambda , solving for  
   theta1  
7 theta1 = asind(n*lambda/a_plus_b);      // Angle at  
                           which first order maximum is observed , degrees  
8 n = 2;                  // second order of diffraction for given  
                           wavelength  
9 theta2 = asind(n*lambda/a_plus_b);      // Angle at  
                           which second order maximum is observed , degrees  
10  
11 printf("\nThe angle at which first order maximum is  
observed = %4.1f degrees", theta1);  
12 printf("\nThe angle at which second order maximum is  
observed = %4.1f degrees", theta2);  
13  
14 // Result  
15 // The angle at which first order maximum is  
   observed = 22.3 degrees  
16 // The angle at which second order maximum is  
   observed = 49.4 degrees
```

---

**Scilab code Exa 3.30** Least width of plane transmission grating

```

1 // Scilab Code Ex3.30:: Page -3.49 (2009)
2 clc; clear;
3 lambda1 = 5890e-008;      // Wavelength of D1 line of
   Na, cm
4 lambda2 = 5896e-008;      // Wavelength of D2 line of
   Na, cm
5 d_lambda = lambda2-lambda1; // Linear
   separation of two lines just seen as separate , cm
6 P = 500;                  // Number of lines per cm on grating
   , lines/cm
7 n = 2;                    // Order of diffraction
8 // As resolving power of grating , lambda/d_lambda =
   n*N, solving for N
9 N = lambda1/(d_lambda*n); // No. of lines
   required per cm on grating , lines/cm
10 w = N/P;                 // Least width of grating , cm
11
12 printf("\nThe least width of plane transmission
   grating = %5.3f cm", w);
13
14 // Result
15 // The least width of plane transmission grating =
   0.982 cm

```

---

**Scilab code Exa 3.31** Minimum grating width required to resolve two wavelengths

```

1 // Scilab Code Ex3.31:: Page -3.49 (2009)
2 clc; clear;
3 theta1 = 18;                // Direction at which first
   spectral line appears , degrees
4 theta2 = 18+5/(60*60);     // Direction at which second
   spectral line appears , degrees
5 d_theta = (theta2-theta1)*%pi/180; // Angular
   separation of two spectral lines , radians

```

```

6 d_lambda = 50e-010;           // Linear separation of two
      spectral lines just seen as separate , cm
7 DP = d_theta/d_lambda;        // Dispersive power of
      grating
8 n = 1;                      // Order of diffraction
9 // As dispersive power of grating d_theta/d_lambda =
      DP = n/((a_plus_b)*cosd(theta1)), solving for
      a_plus_b
10 a_plus_b = n/(DP*cosd(theta1)); // Grating
      element , cm
11 // But a_plus_b*sind(theta1)=n*lambda1, solving for
      lambda1
12 lambda1 = a_plus_b*sind(theta1)/n; // Wavelength of first spectral line , cm
13 lambda2 = lambda1+d_lambda/1e-002; // Wavelength of second spectral line , cm
14 // As resolving power of grating , lambda/d_lambda =
      n*N, solving for N
15 N = lambda1/(d_lambda*n); // No. of lines
      required per cm on grating
16 w = N*a_plus_b; // Minimum grating width
      required to resolve two wavelengths , cm
17
18 printf("\nThe wavelength of first spectral line = %4
      .0 f angstrom" , lambda1/1e-008);
19 printf("\nThe wavelength of second spectral line = %4.0 f angstrom" , lambda2/1e-008);
20 printf("\nThe minimum grating width required to
      resolve two wavelengths = %3.1 f cm" , w);
21
22 // Result
23 // The wavelength of first spectral line = 6702
      angstrom
24 // The wavelength of second spectral line = 6752
      angstrom
25 // The minimum grating width required to resolve two
      wavelengths = 2.9 cm

```

---

**Scilab code Exa 3.32** Angle of diffraction for maxima in first order

```
1 // Scilab Code Ex3.32:: Page -3.50 (2009)
2 clc; clear;
3 // Function to convert theta into degree-minute
4 function [degre, minute]=deg_2_degminsec(theta)
5     degre = floor(theta);
6     minute = (theta-floor(theta))*60;
7 endfunction
8
9 N = 15000;          // No. of lines on the grating per
10    inch, lines/inch
11 a_plus_b = 2.54/N;      // Grating element, cm
12 lambda = 6000e-008;      // Wavelength of light used,
13    cm
14 n = 1;          // Order of diffraction spectra
15 // But a_plus_b*sind(theta)=n*lambda, solving for
16 // theta
17 theta = asind(n*lambda/a_plus_b);           // Direction
18 // in which first order spectra is seen, degrees
19 [deg, mint] = deg_2_degminsec(theta);
20 printf("\nThe angle of diffraction for maxima in
21 first order = %2d degrees %2d min", deg, mint);
22
23 // Result
24 // The angle of diffraction for maxima in first
25 // order = 20 degrees 45 min
```

---

**Scilab code Exa 3.33** Wavelength of light used in obtaining second order diffraction maximum

```
1 // Scilab Code Ex3.33:: Page -3.50 (2009)
```

```

2 clc; clear;
3 N = 12000; // No. of lines on the grating per
   inch , lines/inch
4 a_plus_b = 2.54/N; // Grating element , cm
5 n = 2; // Order of diffraction spectra
6 theta = 39; // Angle of diffraction for maxima in
   second order , degrees
7 // But a_plus_b*sind(theta)=n*lambda , solving for
   lambda
8 lambda = a_plus_b*sind(theta)/n; // Wavelength
   of light used , cm
9
10 printf("\nThe wavelength of light used in obtaining
   second order diffraction maximum = %4d angstrom",
   lambda/1e-008);
11
12 // Result
13 // The wavelength of light used in obtaining second
   order diffraction maximum = 6660 angstrom

```

---

**Scilab code Exa 3.34** Number of visible orders using diffraction grating

```

1 // Scilab Code Ex3.34:: Page -3.51 (2009)
2 clc; clear;
3 lambda = 5890e-008; // Wavelength of light used ,
   cm
4 N = 6000; // No. of lines on the grating per
   inch , lines/inch
5 a_plus_b = 2.54/N; // Grating element , cm
6 theta_max = 90; // Direction of maxima for
   maximum possible orders
7 // But a_plus_b*sind(theta_max)=n*lambda , solving
   for n
8 n = a_plus_b*sind(theta_max)/lambda; // Number
   of visible orders

```

```

9
10 printf("\nThe number of visible orders using
11 diffraction grating = %d", n);
12 // Result
13 // The number of visible orders using diffraction
14 grating = 7

```

---

**Scilab code Exa 3.35** Distance between two wavelengths seen as separate

```

1 // Scilab Code Ex3.35:: Page -3.51 (2009)
2 clc; clear;
3 lambda = 5500e-008;           // Mean of two wavelengths ,
4 cm
5 theta = 35;                  // Angle of diffraction for
6 maxima in second order
7 d_theta = 0.15;              // Angular separation between
8 two neighbouring wavelengths , radians
9 d_lambda = lambda*cotd(theta)*d_theta; // Distance
10 between two wavelengths seen as separate , cm
11 printf("\nThe distance between two wavelengths seen
12 as separate = %d angstrom", d_lambda/1e-008);
13 // Result
14 // The distance between two wavelengths seen as
15 separate = 1178 angstrom

```

---

**Scilab code Exa 3.36** Number of lines per cm on grating surface

```

1 // Scilab Code Ex3.36:: Page -3.51 (2009)
2 clc; clear;

```

```

3 lambda1 = 5500e-008;      // First wavelength of
   light , cm
4 lambda2 = 4500e-008;      // Second wavelength of
   light , cm
5 theta = 45;      // Angle of diffraction for lower
   order , degrees
6 n = lambda2/(lambda1-lambda2); // Lower order of
   diffraction
7 // But a_plus_b*sind(theta)=n*lambda , solving for
   a_plus_b
8 a_plus_b = floor(n)*lambda1/sind(theta);           // 
   Grating element , cm
9 N = 1/a_plus_b;      // No. of lines per cm on
   grating surface , lines/cm
10
11 printf("\nThe number of lines per cm on grating
   surface = %4d lines/cm", ceil(N));
12
13 // Result
14 // The number of lines per cm on grating surface =
   3215 lines/cm

```

---

**Scilab code Exa 3.37** Total number of lines on grating surface

```

1 // Scilab Code Ex3.37:: Page-3.52 (2009)
2 clc; clear;
3 lambda = 6500e-008;      // Wavelength of light used ,
   cm
4 theta = 19.5;      // Angle of diffraction for maxima
   in first order , degrees
5 l = 3.5;      // Length of the grating , cm
6 n = 1;      // Order of diffraction
7 // But a_plus_b*sind(theta)=n*lambda , solving for
   a_plus_b
8 a_plus_b = n*lambda/sind(theta);           // Grating

```

```

    element , cm
9 N = 1/a_plus_b;      // No. of lines per cm on
                        grating surface , lines/cm
10 N_total = l*N;      // Total number of lines on
                        grating surface
11
12 printf("\nThe total number of lines on grating
           surface = %5d", N_total);
13
14 // Result
15 // The total number of lines on grating surface =
           17974

```

---

**Scilab code Exa 3.38** Angular separation between the sodium D1 and D2 lines

```

1 // Scilab Code EX3.38:: Page -3.52 (2009)
2 clc;clear;
3 function [mint, secnd]=degmin(theta)
4     mint = (theta-floor(theta))*60;
5     secnd = (mint-floor(mint))*60
6 endfunction
7 lambda_D1 = 5890e-008; // Wavelength of sodium D1
                           line , cm
8 lambda_D2 = 5896e-008; // Wavelength of sodium D2
                           line , cm
9 n = 2; // Order of diffraction
10 N = 6500; // Number of lines per cm on grating ,
               lines/cm
11 a_plus_b = 1/6500; // Grating element , cm
12 // As a_plus_b*sin(theta1)=n*lambda1 , solving for
               theta1
13 theta1 = asind(n*lambda_D1/a_plus_b);
14 // As a_plus_b*sin(theta2)=n*lambda2 , solving for
               theta1

```

```

15 theta2 = asind(n*lambda_D2/a_plus_b);
16 d_theta = theta2-theta1; // Angular separation
   between the sodium D1 and D2 lines , degrees
17 [mint, secnd] = degmin(d_theta); // Call
   deg_2_degmin function
18
19 printf("\nThe angular separation between the sodium
   D1 and D2 lines = %d minutes %d seconds", mint,
   secnd);
20
21 // Result
22 // The angular separation between the sodium D1 and
   D2 lines = 4 minutes 10 seconds
23 // Since theta1 and theta2 are rounded off in the
   textbook , therefore the answer is mismatching.

```

---

### Scilab code Exa 3.39 Minimum number of lines in a grating

```

1 // Scilab Code EX3.39:: Page-3.55 (2009)
2 clc;clear;
3 lambda1 = 5890e-008; // Wavelength of sodium D1
   line , cm
4 lambda2 = 5896e-008; // Wavelength of sodium D2
   line , cm
5 d_lambda = lambda2-lambda1; // Difference in the
   wavelength of two lines , cm
6 n = 2; // Order of diffraction
7 // As lambda/d_lambda = n*N, solving for N
8 N = lambda1/(d_lambda*n); // Minimum number of
   lines in a grating
9 printf("\nThe minimum number of lines in a grating =
   %3d lines", N);
10
11 // Result
12 // The minimum number of lines in a grating = 490

```

## lines

---

**Scilab code Exa 3.40** Linear separation of two points on the moon

```
1 // Scilab Code EX3.40:: Page-3.56 (2009)
2 clc;clear;
3 lambda = 5500e-008; // Wavelength of most sensitive
color to an eye , cm
4 a = 400; // Aperture of the telescope , cm
5 D = 3.8e+010; // Distance of the moon from the
earth , cm
6 d_theta = 1.22*lambda/a; // Limit of resolution
of telescope , radians
7 // As d_theta = x/D, solving for x
8 x = d_theta*D; // Linear separation of two points
on the moon , cm
9
10 printf("\nThe linear separation of two points on the
moon = %5.2f m", x/1e+002);
11
12 // Result
13 // The linear separation of two points on the moon =
63.74 m
```

---

**Scilab code Exa 3.41** Minimum required number of lines on the plane transmission grating

```
1 // Scilab Code EX3.41:: Page-3.56 (2009)
2 clc;clear;
3 lambda1 = 5890e-008; // Wavelength of sodium D1
line , cm
4 lambda2 = 5896e-008; // Wavelength of sodium D2
line , cm
```

```

5 d_lambda = lambda2-lambda1; // Wavelength difference
   , cm
6 n = 2;      // Order of diffraction
7 // As lambda/d_lambda = n*N, solving for N
8 N = 1/n*(lambda1+lambda2)/(2*d_lambda);    //
   Minimum required number of lines on the plane
   transmission grating
9
10 printf("\nThe minimum required number of lines on
   the plane transmission grating = %3d", N);
11
12 // Result
13 // The minimum required number of lines on the plane
   transmission grating = 491

```

---

**Scilab code Exa 3.42** Number of lines on the plane transmission grating to just resolve the sodium lines

```

1 // Scilab Code EX3.42:: Page-3.57 (2009)
2 clc;clear;
3 lambda1 = 5890e-008; // Wavelength of sodium D1
   line , cm
4 lambda2 = 5896e-008; // Wavelength of sodium D2
   line , cm
5 d_lambda = lambda2-lambda1; // Wavelength difference
   , cm
6 w = 2.5;      // Width of the grating , cm
7 n = 2;      // Order of diffraction
8 // As lambda/d_lambda = n*N, solving for N
9 N = 1/n*(lambda1+lambda2)/(2*d_lambda);    //
   Minimum required number of lines on the plane
   transmission grating
10
11 printf("\nThe number of lines on the plane
   transmission grating to just resolve the sodium

```

```

    lines = %3d" , N/w) ;
12
13 // Result
14 // The number of lines on the plane transmission
   grating to just resolve the sodium lines = 196

```

---

**Scilab code Exa 3.43** Minimum width of the grating to resolve the sodium lines in third order

```

1 // Scilab Code EX3.43:: Page-3.57 (2009)
2 clc;clear;
3 lambda1 = 5890e-008; // Wavelength of sodium D1
   line , cm
4 lambda2 = 5896e-008; // Wavelength of sodium D2
   line , cm
5 d_lambda = lambda2-lambda1; // Wavelength difference
   , cm
6 n = 3; // Order of diffraction
7 P = 2500; // Number of lines per unit length of
   grating
8 // As lambda/d_lambda = n*N, solving for N
9 N = 1/n*(lambda1+lambda2)/(2*d_lambda); // Total
   lines on the grating
10 w = N/P; // Minimum width of the grating , cm
11 printf("\nThe minimum width of the grating to
   resolve the sodium lines in third order = %5.3f
   cm" , w);
12
13 // Result
14 // The minimum width of the grating to resolve the
   sodium lines in third order = 0.131 cm

```

---

**Scilab code Exa 3.44** Dispersive power and diffraction angle for grating

```

1 // Scilab Code EX3.44:: Page -3.57 (2009)
2 clc;clear;
3 w = 2;           // Width of the grating , cm
4 P = 4500;        // Total number of lines on the grating
5 a_plus_b = w/P; // Grating element , cm
6 lambda1 = 5890e-008; // Wavelength of sodium D1
    line , cm
7 lambda2 = 5896e-008; // Wavelength of sodium D2
    line , cm
8 lambda = (lambda1+lambda2)/2; // Mean wavelength of
    light used , cm
9 d_lambda=lambda2-lambda1; // Difference in
    wavelengths of D-lines of sodium , cm
10 n = 2; // Order of diffraction
11 // As a_plus_b*sind(theta)=n*lambda, solving for
    theta
12 theta = asind(n*lambda/a_plus_b); // Angle of
    diffraction , degrees
13 DP = n/(a_plus_b*cosd(theta)); // Dispersive
    power of grating
14 d_theta = DP*d_lambda*180/%pi; // Angular
    separation between D-lines , degrees
15 RP = lambda/d_lambda; // Required resolving power
    of grating for sodium lines
16 N = 2.54/a_plus_b; // No. of lines per cm on
    grating , lines/cm
17 RP_cal = n*N; // Calculated resolving power of
    grating
18
19 printf("\nThe angle of diffraction for maxima in
    second order = %6.4f degrees", d_theta);
20 printf("\nAs %5.3e > %3d, D-lines can be resolved.", ,
    RP_cal, RP);
21
22 // Result
23 // The angle of diffraction for maxima in second
    order = 0.0160 degrees
24 // As 1.143e+04 > 982, D-lines can be resolved.

```

---

**Scilab code Exa 3.45** Distance between centres of the two stars

```
1 // Scilab Code EX3.45:: Page -3.58 (2009)
2 clc;clear;
3 lambda = 5500e-010; // Wavelength of light used , m
4 a = 0.01; // Diameter of objective of telescope , m
5 f = 3.0; // Focal length of telescope objective , m
6 // For telescope , the limit of resolution ,
7 // theta = x/f = 1.22*lambda/a, solving for x
8 x = 1.22*lambda/a*f; // Distance between centres
9 // of imgaes of the two stars
10 printf("\nThe distance between centres of imgaes of
11 the two stars = %4.2e m", x);
12 // Result
13 // The distance between centres of imgaes of the two
14 // stars = 2.01e-04 m
```

---

**Scilab code Exa 3.46** Aperture of the objective of the microscope

```
1 // Scilab Code EX3.46:: Page -3.59 (2009)
2 clc;clear;
3 lambda = 5461e-008; // Wavelength of light used , cm
4 d = 4e-005; // Separation distance between two
5 // self-luminous objects , cm
6 NA = 1.22*lambda/(2*d); // Numerical aperture of
7 // microscope , cm
8
9 printf("\nThe numerical aperture of the objective of
10 the microscopes = %6.4f cm", NA);
```

```
8
9 // Result
10 // The numerical aperture of the objective of the
   microscopes = 0.8328 cm
```

---

# Chapter 4

## Polarization

**Scilab code Exa 4.1** Refractive index of the material

```
1 // Scilab Code Ex4.1:: Page-4.5 (2009)
2 clc; clear;
3 ip = 60;      // Polarizing angle, degrees
4 mu = tand(ip); // Refractive index of the material
                  from Brewster's law
5 printf("\nThe refractive index of the material = %5
.3 f", mu);
6
7 // Result
8 // The refractive index of the material = 1.732
```

---

**Scilab code Exa 4.2** Polarization by reflection

```
1 // Scilab Code Ex4.2:: Page-4.6 (2009)
2 clc; clear;
3 ip = 57;      // Polarizing angle, degrees
4 mu = tand(ip); // Refractive index of the material
                  from Brewster's law
```

```
5 printf("\nThe refractive index of the material = %4
       .2f", mu);
6
7 // Result
8 // The refractive index of the material = 1.54
```

---

### Scilab code Exa 4.3 Angle of refraction of the ray

```
1 // Scilab Code Ex4.3:: Page-4.6 (2009)
2 clc; clear;
3 mu = 1.53; // Refractive index of the material from
             Brewster's law
4 // As mu = tand(ip), solving for ip
5 ip = atand(mu); // Polarizing angle, degrees
6 // But mu = sind(ip)/sind(r), solving for r
7 r = asind(sind(ip)/mu); // Angle of refraction,
                           degrees
8
9 printf("\nThe angle of refraction of the ray = %4.1f
       degrees", r);
10
11 // Result
12 // The angle of refraction of the ray = 33.2 degrees
```

---

### Scilab code Exa 4.4 Angle of minimum deviation for green light

```
1 // Scilab Code Ex4.4:: Page-4.6 (2009)
2 clc; clear;
3 ip = 60; // Polarizing angle, degrees
4 A = 60; // Angle of equilateral prism, degrees
5 mu = tand(ip); // Refractive index of the material
                   from Brewster's law
```

```

6 // For angle of minimum deviation in prism , delta_m ,
    refractive index
7 // mu = sind ((A+delta_m)/2)/sind (A/2) , solving for
    delta_m
8 delta_m = 2*asind(mu*sind(A/2))-A;           // Angle of
    minimum deviation , degrees
9
10 printf("\nThe angle of minimum deviation for green
    light = %2d degrees", ceil(delta_m));
11
12 // Result
13 // The angle of minimum deviation for green light =
    60 degrees

```

---

**Scilab code Exa 4.5** Polarizing angles of the materials for given refractive indices

```

1 // Scilab Code Ex4.5:: Page-4.7 (2009)
2 clc; clear;
3 mu = [1.33 1.65 1.55];           // Refractive indices of
    the material
4 // As mu = tand(ip) , solving for ip
5 ip = atand(mu);                 // Brewster's law gives
    polarizing angle , degrees
6 for i =1:1:3
7 printf("\nmu = %4.2f , ip = %4.1f degrees", mu(i), ip
    (i));
8 end
9
10 // Result
11 // mu = 1.33 , ip = 53.1 degrees
12 // mu = 1.65 , ip = 58.8 degrees
13 // mu = 1.55 , ip = 57.2 degrees

```

---

### **Scilab code Exa 4.6 Angle of rotation of analyser**

```
1 // Scilab Code Ex4.6:: Page-4.8 (2009)
2 clc; clear;
3 E0 = 1;      // For simplicity assume maximum
               intensity through polarizer and analyser to be
               unity , unit
4 E = 1/6*E0; // One-sixth of the maximum intensity ,
               unit
5 // From Malus law , E = E0*cosd(theta)^2, solving for
               theta
6 theta = acosd(sqrt(E));    // Angle through which
               analyser should be rotated , degrees
7 printf("\nThe angle of rotation of analyser = %4.1f
               degrees", theta);
8
9 // Result
10 // The angle of rotation of analyser = 65.9
```

---

### **Scilab code Exa 4.7 Angles of rotation of analyser for given transmitted light intensities**

```
1 // Scilab Code Ex4.7:: Page-4.8 (2009)
2 clc; clear;
3 E0 = 1;      // For simplicity assume maximum
               intensity through polarizer and analyser to be
               unity , unit
4 light_fraction = [0.25 0.45 0.65 0.75 0.0];
5 for i = 1:1:5
6 E = light_fraction(i)*E0; // Light fraction of the
               maximum intensity , unit
```

```

7 // From Malus law , E = E0*cosd(theta)^2, solving for
   theta
8 theta = acosd(sqrt(E)); // Angle through which
   analyser should be rotated , degrees
9 printf("\nE = %4.2fE0 , theta = %4.1f degrees" ,
   light_fraction(i), theta);
10 end
11
12 // Result
13 // E = 0.25E0 , theta = 60.0 degrees
14 // E = 0.45E0 , theta = 47.9 degrees
15 // E = 0.65E0 , theta = 36.3 degrees
16 // E = 0.75E0 , theta = 30.0 degrees
17 // E = 0.00E0 , theta = 90.0 degrees

```

---

### Scilab code Exa 4.8 Angle of minimum deviation for green light

```

1 // Scilab Code Ex4.8:: Page-4.9 (2009)
2 clc; clear;
3 ip = 60;           // Polarizing angle , degrees
4 mu = tand(ip);   // Brewster's law giving refractive
   index
5 A = 60;           // Angle of prism , degrees
6 d = (mu - 1)*A; // Angle of minimum deviation for
   green light , degrees
7
8 printf("\nThe angle of minimum deviation for green
   light = %5.2f degrees" , d);
9
10 // Result
11 // The angle of minimum deviation for green light =
   43.92 degrees

```

---

### Scilab code Exa 4.9 Ratio of ordinary to extraordinary ray intensities

```
1 // Scilab Code Ex4.9:: Page-4.9 (2009)
2 clc; clear;
3 theta = 30;      // Angle which the plane of
                  vibration makes with the incident beam, degrees
4 // As intensity of ordinary and extraordinary ray
      are
5 // E_E = A^2*cosd(theta)^2 and E_O = A^2*sind(theta)
      ^2, solving for E_E/E_O
6 EE_ratio_E0 = cotd(30)^2;      // Ratio of ordinary
      and extraordinary ray intensities
7
8 printf("\nThe ratio of ordinary to extraordinary ray
      intensities = %d", EE_ratio_E0);
9
10 // Result
11 // The ratio of ordinary to extraordinary ray
      intensities = 3
```

---

### Scilab code Exa 4.10 Thickness of quarter wave plate

```
1 // Scilab Code Ex4.10:: Page-4.23 (2009)
2 clc; clear;
3 mu_o = 1.658;      // Refractive index of ordinary wave
4 mu_e = 1.486;      // Refractive index of extraordinary
      wave
5 lambda = 5893e-008; // Wavelength of light used , m
6 // As (mu_o - mu_e)*t = lambda/4, solving for t
7 t = lambda/(4*(mu_o - mu_e));      // Thickness of
      quarter-wave plate , cm
8
9 printf("\nThe thickness of quarter-wave plate = %3.1
      e cm", t);
10
```

```
11 // Result  
12 // The thickness of quarter-wave plate = 8.6e-005 cm
```

---

**Scilab code Exa 4.11** Least thickness of plate for which emergent beam is plane polarised

```
1 // Scilab Code Ex4.11:: Page-4.23 (2009)  
2 clc; clear;  
3 mu_o = 1.5442; // Refractive index of ordinary  
wave  
4 mu_e = 1.5533; // Refractive index of  
extraordinary wave  
5 lambda = 5000e-008; // Wavelength of light used , m  
6 // As (mu_o - mu_e)*t = lambda/4, solving for t  
7 t = lambda/(4*(mu_e - mu_o)); // Least thickness  
of plate for which emergent beam is plane  
polarised , cm  
8  
9 printf("\nThe least thickness of plate for which  
emergent beam is plane polarised = %4.2e cm", t);  
10  
11 // Result  
12 // The least thickness of plate for which emergent  
beam is plane polarised = 1.37e-003 cm
```

---

**Scilab code Exa 4.12** Difference in refractive indices of rays

```
1 // Scilab Code Ex4.12:: Page-4.23 (2009)  
2 clc; clear;  
3 lambda = 5893e-008; // Wavelength of light used , m  
4 t = 0.005; // Thickness of the crystal , cm  
5 // As for quarter wave plate , mu_diff*t = (mu_o -  
mu_e)*t = lambda/4, solving for mu_diff
```

```

6 mu_diff = lambda/(4*t);      // The difference in
                                refractive indices of rays , cm
7 printf("\nThe least thickness of plate for which
          emergent beam is plane polarised = %4.2e cm",
          mu_diff);
8
9 // Result
10 // The least thickness of plate for which emergent
    beam is plane polarised = 2.95e-003 cm

```

---

**Scilab code Exa 4.13** The thickness of a half wave plate

```

1 // Scilab Code Ex4.13:: Page -4.24 (2009)
2 clc; clear;
3 mu_o = 1.54;      // Refractive index of ordinary wave
4 mu_e = 1.45;      // Refractive index of extraordinary
                    wave
5 lambda = 5500e-008; // Wavelength of light used , m
6 // As for a half wave plate , (mu_o - mu_e)*t =
                    lambda/4, solving for t
7 t = lambda/(2*(mu_o - mu_e));    // The thickness of
                    a half wave plate for wavelength , cm
8
9 printf("\nThe thickness of a half wave plate for
        wavelength = %4.2e cm", t);
10
11 // Result
12 // The thickness of a half wave plate for wavelength
    = 3.06e-004 cm

```

---

**Scilab code Exa 4.14** The thickness of a quarter wave plate

```
1 // Scilab Code Ex4.14:: Page -4.24 (2009)
```

```

2 clc; clear;
3 mu_o = 1.55; // Refractive index of ordinary wave
4 mu_e = 1.52; // Refractive index of extraordinary
    wave
5 lambda = 5500e-008; // Wavelength of light used , m
6 // As for a half wave plate , (mu_o - mu_e)*t =
    lambda/4, solving for t
7 t = lambda/(4*(mu_o - mu_e)); // The thickness of
    a quarter wave plate for wavelength , cm
8
9 printf("\nThe thickness of a quarter wave plate for
wavelength = %4.2e cm", t);
10
11 // Result
12 // The thickness of a quarter wave plate for
wavelength = 4.58e-004 cm

```

---

**Scilab code Exa 4.15** The thickness of a half wave plate quartz

```

1 // Scilab Code Ex4.15:: Page -4.24 (2009)
2 clc; clear;
3 mu_o = 1.51; // Refractive index of ordinary wave
4 mu_e = 1.55; // Refractive index of extraordinary
    wave
5 lambda = 6000e-008; // Wavelength of light used , m
6 // As for a half wave plate , (mu_o - mu_e)*t =
    lambda/4, solving for t
7 t = lambda/(2*(mu_e - mu_o)); // The thickness of
    a quarter wave plate for wavelength , cm
8
9 printf("\nThe thickness of a half wave plate quartz
= %4.2e cm", t);
10
11 // Result
12 // The thickness of a half wave plate quartz = 7.50e

```

-004 cm

---

### Scilab code Exa 4.16 Difference between refractive indices

```
1 // Scilab Code Ex4.16:: Page-4.24 (2009)
2 clc; clear;
3 lambda = 5890e-008; // Wavelength of light used , m
4 t = 7.5e-004; // Thickness of the crystal , cm
5 // As for quarter wave plate , mu_diff*t = (mu_e -
    mu_o)*t = lambda/4, solving for mu_diff
6 mu_diff = lambda/(4*t); // The difference in
    refractive indices of rays , cm
7 printf("\nThe difference between refractive indices
        = %6.4f cm", mu_diff);
8
9 // Result
10 // The difference between refractive indices =
    0.0196 cm
```

---

### Scilab code Exa 4.17 Specific rotation of superposition

```
1 // Scilab Code Ex4.17:: Page-4.34 (2009)
2 clc; clear;
3 theta = 15.2; // Angle through which plane of
    polarization is rotated , degrees
4 c = 0.2; // Concentration of sugar , g/cc
5 l = 25; // Length of sugar , cm
6 S = 10*theta/(l*c); // Specific rotation of
    superposition , degrees
7
8 printf("\nThe specific rotation of superposition =
        %4.1f cm", S);
9
```

```
10 // Result
11 // The specific rotation of superposition = 30.4 cm
```

---

**Scilab code Exa 4.18** Strength of sugar solution

```
1 // Scilab Code Ex4.18: : Page-4.34 (2009)
2 clc; clear;
3 theta = 15.2; // Angle through which plane of
    polarization is rotated , degrees
4 S = 65; // Specific rotation of sugar solution ,
    degrees
5 l = 15; // Length of sugar , cm
6 // As S = 10*theta/(l*c) , solving for c
7 c = 10*theta/(l*S); // Concentration of sugar , g
    /cc
8
9 printf("\nThe strength of sugar solution = %4.2f g/
    cc", c);
10
11 // Result
12 // The strength of sugar solution = 0.16 g/cc
```

---

**Scilab code Exa 4.19** Quantity of sugar contained in the tube in the form of solution

```
1 // Scilab Code Ex4.19:: Page-4.34 (2009)
2 clc; clear;
3 theta = 15; // Angle through which plane of
    polarization is rotated , degrees
4 S = 69; // Specific rotation of sugar solution ,
    degrees
5 l = 10; // Length of sugar , cm
6 V = 50; // Volume of the tube , cc
```

```

7 // As S = 10*theta/(l*c) , solving for c
8 c = 10*theta/(l*S);      // Concentration of sugar , g
   /cc
9 M = c*V;      // Mass of sugar in solution , g
10
11 printf("\nThe quantity of sugar contained in the
        tube in the form of solution = %5.2f g" , M);
12
13 // Result
14 // The quantity of sugar contained in the tube in
        the form of solution = 10.87 g

```

---

**Scilab code Exa 4.20** Specific rotation of sugar solution from the given data

```

1 // Scilab Code Ex4.20:: Page-4.35 (2009)
2 clc; clear;
3 theta = 8;      // Angle through which plane of
                  polarization is rotated , degrees
4 M = 10;         // Amount of sugar , g
5 l = 14;         // Length of the tube , cm
6 V = 44;         // Volume of sugar solution , cc
7 c = M/V;        // Concentration of sugar , g/cc
8 S = 10*theta/(l*c); // Specific rotation of sugar
                      solution from the given data , degrees
9
10 printf("\nThe specific rotation of sugar solution
          from the given data = %4.1f degrees" , S);
11
12 // Result
13 // The specific rotation of sugar solution from the
          given data = 25.1 degrees

```

---

**Scilab code Exa 4.21** Angle of rotation of the plane of polarization

```
1 // Scilab Code Ex4.21:: Page-4.35 (2009)
2 clc; clear;
3 m = 15;      // Amount of sugar , g
4 S = 66; // Specific rotation of sugar solution from
          the given data , degrees
5 l = 20;      // Length of the tube , cm
6 V = 100;     // Volume of sugar solution , cc
7 c = m/V;     // Concentration of sugar , g/cc
8 // As S = 10*theta/(l*c) , solving for theta
9 theta = S*l*c/10;      // Angle of rotation of the
                           plane of polarization , degrees
10
11 printf("\nThe angle of rotation of the plane of
          polarization = %4.1f degrees", theta);
12
13 // Result
14 // The angle of rotation of the plane of
          polarization = 19.8 degrees
```

---

**Scilab code Exa 4.22** Angle of rotation of the optically active solution

```
1 // Scilab Code Ex4.22: : Page-4.35 (2009)
2 clc; clear;
3 l = 5;      // Length of the tube , dm
4 m = 50;     // Amount of sugar , g
5 S = 50; // Specific rotation of sugar solution ,
          degrees
6 V = 150;     // Volume of sugar solution , cc
7 c = m/V;     // Concentration of sugar , g/cc
8 // As S = theta/(l*c) , solving for theta
9 theta = S*l*c;      // Angle of rotation of the
                           optically active solution
10
```

```
11 printf("\nThe angle of rotation of the optically
       active solution = %4.1f degrees", theta);
12
13 // Result
14 // The angle of rotation of the optically active
   solution = 83.3 degrees
```

---

**Scilab code Exa 4.23** Angle of rotation in a tube of new length

```
1 // Scilab Code Ex4.23:: Page-4.35 (2009)
2 clc; clear;
3 l = 3;      // Length of the tube, dm
4 theta = 17.0;      // Angle of rotation of the
   plane of polarization, degrees
5 c = 1.0;      // For simplicity assume concentration
   of solution to be unity, g/cc
6 l_prime = 2.5;      // New length of the tube, dm
7 c_prime = 1.25*c; // Concentration of solution with
   25 cm length of tube, g/cc
8 theta_prime = theta*l_prime*c_prime/(l*c); // Angle
   of rotation in a tube of new length
9
10
11 printf("\nThe angle of rotation in a tube of new
       length of %3.1f cm = %4.1f degrees", l_prime,
   theta_prime);
12
13 // Result
14 // The angle of rotation in a tube of new length of
   2.5 cm = 17.7 degrees
```

---

**Scilab code Exa 4.24** Mass of sugar in the solution contained in the tube

```

1 // Scilab Code Ex4.24:: Page -4.36 (2009)
2 clc; clear;
3 l = 17;      // Length of the tube , cm
4 V = 37;      // Volume of sugar solution , cc
5 theta = 15;    // Angle of rotation of the plane
                  of polarization , degrees
6 S = 68;       // Specific rotation of sugar
                  solution , degrees
7 // As S = 10*theta/(l*c) , solving for c
8 c = 10*theta/(l*S);   // Concentration of sugar
                  solution , g/cc
9 m = c*V;       // Mass of sugar in the solution
                  contained in the tube , g
10
11 printf("\nThe mass of sugar in the solution
           contained in the tube = %3.1f g", m);
12
13 // Result
14 // The mass of sugar in the solution contained in
           the tube = 4.8 g

```

---

### Scilab code Exa 4.25 Percentage purity of the sugar sample

```

1 // Scilab Code Ex4.25:: Page -4.36 (2009)
2 clc; clear;
3 m = 80;       // Mass of sugar in the solution , g
4 theta = 9.9;    // Angle of rotation of the plane
                  of polarization , degrees
5 l = 20;       // Length of the tube , cm
6 S_pure = 66;    // Specific rotation of pure
                  sugar solution , degrees per dm per (g/cc)
7 c = 0.08;      // Concentration of sugar solution , g/
                  cc
8 S = 10*theta/(l*c); // calculated specific rotation
                  of sugar solution , degrees per dm per (g/cc)

```

```

9 percent_purity = S/S_pure*100;           // Percentage
    purity of sugar sample , percent
10
11 printf("\nThe percentage purity of the sugar sample
    = %5.2f percent", percent_purity);
12
13 // Result
14 // The percentage purity of the sugar sample = 93.75
    percent

```

---

**Scilab code Exa 4.26** Angle of rotation produced by the polarimeter plate

```

1 // Scilab Code Ex4.26:: Page-4.42 (2009)
2 clc; clear;
3 lambda = 6600e-010;           // Wavelength of circularly
    polarized light , cm
4 mu_R = 1.53914;             // Refractive index of right
    -handed circularly polarized light
5 mu_L = 1.53920;             // Refractive index of left-
    handed circularly polarized light
6 t = 0.0005;                 // Thickness of polarimeter plate , m
7 theta = %pi/lambda*(mu_L-mu_R)*t;   // Angle of
    rotation produced by the polarimeter plate ,
    radian
8
9 printf("\nThe angle of rotation produced by the
    polarimeter plate = %4.2f degrees", theta*180/%pi
);
10
11 // Result
12 // The angle of rotation produced by the polarimeter
    plate = 8.18 degrees

```

---

# Chapter 5

## Nuclear Physics

**Scilab code Exa 5.1** Mass defect of He

```
1 // Scilab Code Ex5.1 :: Page-5.2 (2009)
2 clc;clear;
3 m_p = 1.007826;           // Mass of a proton , amu
4 m_n = 1.008665;           // Mass of a neutron , amu
5 M_He = 4.002604;          // Measured mass of He nucleuc ,
                                amu
6 delta_m = 2*m_p+2*m_n - M_He;    // Mass defect of He
                                , amu
7 printf("\nThe mass defect of He = %f amu", delta_m);
8
9 // Result
10 // The mass defect of He = 0.030378 amu
```

---

**Scilab code Exa 5.3** Maximum energy of proton in a cyclotron

```
1 // Scilab Code Ex5.3 :: Page-5.16 (2009)
2 clc;clear;
3 B = 0.70;                  // Magnetic field of cyclotron ,
                                weber/metre square
```

```

4 q = 1.6e-019; // Charge of the proton , C
5 R = 3; // Radius of Dee's , m
6 m = 1.67e-027; // Mass of the proton , kg
7 E_max = B^2*q^2*R^2/(2*m); // Maximum energy of the
     proton in the cyclotron , joule
8 printf("\nThe maximum energy of the proton in the
     cyclotron = %4.2e MeV" , E_max/1.6e-013);
9
10 // Result
11 // The maximum energy of the proton in the cyclotron
     = 2.11e+02 MeV
12 // The unit has been given wrong in the textbook. It
     should be MeV instead of eV

```

---

**Scilab code Exa 5.4** Energy of an electron in a betatron

```

1 // Scilab Code Ex5.4 :: Page -5.20 (2009)
2 clc;clear;
3 f = 1e+06; // Frequency of revolution of
     electron , Hz
4 rate_phi_B = 25; // Rate of change of magnetic
     flux , wb/s
5 E = f*rate_phi_B; // Energy of 'f' revolutions , eV
6 printf("\nThe energy of the electron in Betatron
     after %g revolutions = %3.1e eV" , f , E);
7
8 // Result
9 // The energy of the electron in Betatron after 1e
     +06 revolutions = 2.5e+07 eV

```

---

**Scilab code Exa 5.5** Final energy gained by electrons in a betatron

```

1 // Scilab Code Ex5.5 :: Page -5.20 (2009)

```

```

2 clc;clear;
3 e = 1.6e-019; // Charge on an electron , C
4 D = 2.0; // Diameter of the stable orbit in
            betatron , m
5 R = D/2; // Radius of the stable orbit in
            betatron , m
6 B = 0.5; // Magnetic field of betatron , wb/metre
            square
7 c = 3e+08; // final speed of electron in betatron ,
            m/s
8 E = B*e*R*c; // Final energy gained by electrons
            in a betatron , eV
9 printf("\nThe final energy gained by electrons in
            the betatron = %3.1e eV" , E/e);
10
11 // Result
12 // The final energy gained by electrons in the
            betatron = 1.5e+08 eV

```

---

### Scilab code Exa 5.6 Energy produced in fission of U235

```

1 // Scilab Code Ex5.6 :: Page-5.27 (2009)
2 clc;clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 A = 235; // Atomic weight of uranium , gm/mol
5 N_A = 6.023e+026; // No. of atoms present in 235
            kg of uranium
6 N = N_A/(A*1000); // No. of nuclci of uranium per
            gram
7 E = N*200; // Energy produced by 1 g of U-235, MeV
8 printf("\nThe energy produced by 1 g of U-235 = %3.1
            e joule" , E*e*1e+06);
9
10 // Result
11 // The energy produced by 1 g of U-235 = 8.2e+10

```

joule

---

### Scilab code Exa 5.7 Power output of nuclear reactor

```
1 // Scilab Code Ex5.7 :: Page-5.32 (2009)
2 clc;clear;
3 A = 235;      // Atomic weight of uranium , gm/mol
4 N_A = 6.023e+026;    // No. of atoms present in 235
                      kg of uranium-235
5 N = N_A*A;   // No. of nuclci of uranium in 5 kg of
                  U-235
6 E = N*200;    // Energy released in the fission of 5
                  kg of U-235, MeV
7 t = 24*3600;  // Time taken to consume 5 kg of U
                  -235, sec
8 P = E/t;      // Total power output of the nuclear
                  reactor , MeV per second
9 printf("\nThe total power output of the nuclear
        reactor = %4.2e MeV per second", P);
10
11 // Result
12 // The total power output of the nuclear reactor =
        2.97e+22 MeV per second
```

---

### Scilab code Exa 5.8 Average current in the GM counter circuit

```
1 // Scilab Code Ex5.8 :: Page-5.34 (2009)
2 clc;clear;
3 e = 1.6e-019;    // Electronic charge , C
4 f = 450;         // Count rate of GM counter , counts/min
5 N = f*1e+08;    // Total number of electrons
                      collected per min
6 Q = N*e;        // Charge collected per min, C
```

```

7 I = Q/60; // Average current in the GM counter , A
8 printf("\nThe average current in the GM counter= %3.1
e A", I);
9
10 // Result
11 // The average current in the GM counter= 1.2e-10 A

```

---

**Scilab code Exa 5.9** Energy needed to remove a neutron from Ca nucleus

```

1 // Scilab Code Ex5.9 :: Page-5.39 (2009)
2 clc;clear;
3 m_Ca_41 = 40.962278; // Mass of one Ca-41 nuclei ,
amu
4 m_Ca_42 = 41.958618; // Mass of one Ca-42 nuclei ,
amu
5 m_n = 1.008665; // Mass of a neutron , amu
6 delta_m = m_Ca_42 - (m_Ca_41 + m_n); //
Difference in the mass of Ca-42 and Ca-41 nuclei ,
amu
7 E = delta_m*(931.49); // Binding energy of the
missing neutron , MeV
8 printf("\nThe energy needed to remove a neutron from
Ca-42 nucleus = %5.2f MeV", abs(E));
9
10 // Result
11 // The energy needed to remove a neutron from Ca-42
nucleus = 11.48 MeV

```

---

# Chapter 6

## Semiconductors and Nano Physics

**Scilab code Exa 6.1** Resistivity of intrinsic semiconductor at 300 K

```
1 // Scilab Code Ex6.1:: Page-6.19 (2009)
2 clc; clear;
3 T = 300;      // Temperature of pure semiconductor , K
4 n_i = 2.5e+019; // Intrinsic carrier density , per
                  metre square
5 e = 1.6e-019;   // Charge on an electron , C
6 mu_e = 0.39;    // Mobility of electrons , Sq.m/V/s
7 mu_h = 0.19;    // Mobility of holes , Sq.m/V/s
8 sigma_i = e*n_i*(mu_e+mu_h); // Conductivity of
                                intrinsic semiconductor at 300 K, mho/m
9 rho_i = 1/sigma_i; // Resistivity of intrinsic
                     semiconductor at 300 K, ohm-m
10
11 printf("\nThe resistivity of intrinsic semiconductor
        at 300 K = %4.2f ohm-m", rho_i);
12
13 // Result
14 // The resistivity of intrinsic semiconductor at 300
    K = 0.43 ohm-m
```

---

### Scilab code Exa 6.2 Velocity of electron at Fermi level

```
1 // Scilab Code Ex6.2: : Page -6.19 (2009)
2 clc; clear;
3 e = 1.6e-019;      // Energy equivalent of 1 eV, J/eV
4 E_F = 2.0*e;       // Fermi level of Po, J
5 m = 9.1e-031;      // Mass of an electron , kg
6 // As E_F = 1/2*m*v^2, solving for v
7 v = sqrt(2*E_F/m); // Velocity of electron at
                      Fermi level , m/s
8 printf("\nThe Velocity of electron at Fermi level =
         %4.2e m/s", v);
9
10 // Result
11 // The Velocity of electron at Fermi level = 8.39e
     +05 m/s
```

---

# Chapter 7

## Fiber Optics

**Scilab code Exa 7.1** Critical angle and acceptance angle in an optical fibre

```
1 // Scilab Code Ex7.1:: Page-7.7 (2009)
2 clc; clear;
3 n1 = 1.6;           // Refractive index of core material
                      of fibre
4 n2 = 1.3;           // Refractive index of cladding
                      material of fibre
5 phi_C = asind(n2/n1); // Critical angle of optical
                      fibre , degrees
6 theta_Q = asind(sqrt(n1^2-n2^2)); // Acceptance
                      angle of optical fibre , degrees
7
8 printf("\nThe critical angle of optical fibre = %4.1
        f degrees", phi_C);
9 printf("\nThe angle of acceptance cone = %5.1f
        degrees", 2*theta_Q);
10
11 // Result
12 // The critical angle of optical fibre = 54.3
        degrees
13 // The angle of acceptance cone = 137.7 degrees
```

---

**Scilab code Exa 7.2** Critical angle acceptance angle and numerical aperture in an optical fibre

```
1 // Scilab Code Ex7.2:: Page-7.8 (2009)
2 clc; clear;
3 n1 = 1.50;           // Refractive index of core
                      material of fibre
4 n2 = 1.47;           // Refractive index of cladding
                      material of fibre
5 phi_C = asind(n2/n1); // Critical angle of optical
                        fibre , degrees
6 NA = sqrt(n1^2-n2^2); // Numerical aperture for
                        the fibre
7 theta_Q = asind(sqrt(n1^2-n2^2)); // Acceptance
                        angle of optical fibre , degrees
8
9 printf("\nThe critical angle of optical fibre = %4.1
       f degrees", phi_C);
10 printf("\nThe numerical aperture for the fibre = %5
        .3 f", NA);
11 printf("\nThe angle of acceptance cone = %5.1 f
        degrees", theta_Q);
12
13 // Result
14 // The critical angle of optical fibre = 78.5
       degrees
15 // The numerical aperture for the fibre = 0.298
16 // The angle of acceptance cone = 17.4 degrees
```

---

**Scilab code Exa 7.3** Parameters of an optical fibre using relative refractive index difference

```

1 // Scilab Code Ex7.3:: Page-7.8 (2009)
2 clc; clear;
3 n1 = 1.46;      // Refractive index of the core
                  material
4 delta = 0.01;    // Relative refractive index
                  difference
5 NA = n1*sqrt(2*delta);   // Numerical aperture for
                  the fibre
6 theta_Q = %pi*NA^2;    // Solid acceptance angle of
                  optical fibre for small angles , radians
7 // As relative refractive index , delta = 1-n2/n1 ,
                  solving for n2
8 n2 = n1*(1-delta);    // Refractive index of cladding
9 phi_C = asind(n2/n1);  // Critical angle of optical
                  fibre , degrees
10
11 printf("\nThe numerical aperture for the fibre = %4
.2f", NA);
12 printf("\nThe solid acceptance angle of the optical
fibre = %4.2f radians", theta_Q);
13 printf("\nThe critical angle of optical fibre = %4.1
f degrees", phi_C);
14
15 // Result
16 // The numerical aperture for the fibre = 0.21
17 // The solid acceptance angle of the optical fibre
= 0.13 radians
18 // The critical angle of optical fibre = 81.9
degrees

```

---

### Scilab code Exa 7.4 Refractive index of cladding

```

1 // Scilab Code Ex7.4:: Page-7.9 (2009)
2 clc; clear;
3 n1 = 1.54;      // Refractive index of the core

```

```

        material
4 NA = 0.45;      // Numerical aperture for the fibre
5 n2 = sqrt(n1^2-NA^2);    // Refractive index of
                           cladding
6
7 printf("\nThe refractive index of cladding = %4.2f",
       n2);
8
9 // Result
10 // The refractive index of cladding = 1.47

```

---

**Scilab code Exa 7.5** Numerical aperture for an optical fibre

```

1 // Scilab Code Ex7.5:: Page-7.9 (2009)
2 clc; clear;
3 n1 = 1.544;      // Refractive index of the core
                   material
4 n2 = 1.412;      // Refractive index of cladding
5 NA = sqrt(n1^2-n2^2);    // Numerical aperture for
                           the fibre
6
7 printf("\nThe numerical aperture for an optical
           fibre = %4.2f", NA);
8
9 // Result
10 // The numerical aperture for an optical fibre =
0.62

```

---

**Scilab code Exa 7.6** Refractive index of the cladding

```

1 // Scilab Code Ex7.6:: Page-7.9 (2009)
2 clc; clear;

```

```

3 n1 = 1.544;      // Refractive index of the core
                    material
4 theta0 = 35;     // Acceptance angel for an optical
                    fibre , degrees
5 // As theta0 = asind(sqrt(n1^2-n2^2)), solving for
n2
6 n2 = sqrt(n1^2-sind(theta0)^2);      // Refractive
                    index of cladding
7
8 printf("\nThe refractive index of the cladding = %4
.2 f", n2);
9
10 // Result
11 // The refractive index of the cladding = 1.43

```

---

**Scilab code Exa 7.7** Comparison of the acceptance angle for meridional rays with that for the skew rays

```

1 // Scilab Code Ex7.7:: Page-7.10 (2009)
2 clc; clear;
3 NA = 0.4;      // Numerical aperture of the optical
                    fibre
4 n0 = 1;        // Refractive index of fibre in air
5 theta_a = asind(NA/n0); // Acceptance angle for
                    meridional rays , degrees
6 theta = 100;    // Direction through which the skew
                    rays are bent at each reflection , degrees
7 r = theta/2;    // Angle of reflection , degrees
8 theta_as = asind(NA/(cosd(r)*n0)); // Acceptance
                    angle for skew rays , degrees
9
10 printf("\nAcceptance angle for meridional rays = %4
.1 f degrees", theta_a);
11 printf("\nAcceptance angle for skew rays = %4.1 f
degrees", theta_as);

```

```
12
13 // Result
14 // Acceptance angle for meridional rays = 23.6
15 // degrees
```

---

**Scilab code Exa 7.8** Normalized frequency for V number for the fibre

```
1 // Scilab Code Ex7.8: : Page-7.13 (2009)
2 clc; clear;
3 NA = 0.16;      // Numerical aperture of the step
                  index fibre
4 n1 = 1.50;      // Refractive index of the core
                  material
5 d = 65e-006;    // Diameter of the core , m
6 lambda = 0.9e-006; // Wavelength of transmitted
                     light , m
7 V = %pi*d/lambda*NA; // V-number for the optical
                  fibre
8
9 printf("\nThe V-number for the optical fibre = %5.2f
         ", V);
10
11 // Result
12 // The V-number for the optical fibre = 36.30
```

---

**Scilab code Exa 7.9** Number of modes in the step index fibre

```
1 // Scilab Code Ex7.9:: Page-7.13 (2009)
2 clc; clear;
3 NA = 0.28;      // Numerical aperture of the step
                  index fibre
4 d = 55e-006;    // Diameter of the core , m
```

```

5 lambda = 0.9e-006; // Wavelength of transmitted
light , m
6 M_N = (2.22*d*(NA)/lambda)^2; // Number of modes
in the step index fibre
7
8 printf("\nThe number of modes in the step index
fibre = %4d degrees", M_N);
9
10 // Result
11 // The number of modes in the step index fibre =
1442 degrees

```

---

**Scilab code Exa 7.10** Radius of core for single mode operation in step index fibre

```

1 // Scilab Code Ex7.10:: Page -7.14 (2009)
2 clc; clear;
3 n1 = 1.480; // Refractive index of core material
4 n2 = 1.47; // Refractive index of cladding
material
5 lambda = 850e-006; // Wavelength of light used , m
6 NA = sqrt(n1^2-n2^2); // Numerical aperture of
the step index fibre
7 theta0 = asind(NA); // Maximum acceptance angle
for the fibre , degrees
8 M_N = 1; // Number of modes in step index cable
9 // As number of modes, MN = 1/2*V^2, solving for V
10 V = sqrt(2*M_N); // V-number for the fibre
11 // As V = 2*pi*a/lambda*NA, solving for a
12 a = V*lambda/(2*pi*NA); // Radius of core for
single mode operation in step index fibre , m
13
14 printf("\nThe radius of core for single mode
operation in step index fibre = %3.1e" , a);
15

```

```
16 // Result  
17 // The radius of core for single mode operation in  
// step index fibre = 1.1e-03  
18 // The answer is quoted wrong in the textbook
```

---

### Scilab code Exa 7.11 Signal attenuation in optical fibre

```
1 // Scilab Code Ex7.11: : Page - 7.16 (2009)  
2 clc; clear;  
3 Pi = 1.5; // Input power to the optical fibre , mW  
4 Po = 0.5; // Output power to the optical fibre , mW  
5 L = 0.12; // Length of the optical fibre , km  
6 alpha_dB = 10/L*log10(Pi/Po); // Signal attenuation  
// in optical fibre , dB/km  
7  
8 printf("\nThe signal attenuation in optical fibre =  
%4.1f dB/km", alpha_dB);  
9  
10 // Result  
11 // The signal attenuation in optical fibre = 39.8 dB  
// /km
```

---

# Chapter 8

## Laser

**Scilab code Exa 8.1** Difference between upper and lower energy levels for the most prominent wavelength

```
1 // Scilab Code Ex8.1:: Page-8.8 (2009)
2 clc; clear;
3 lambda = 31235; // Wavelength of prominent
      emission of laser , aangstrom
4 E = 12400/lambda; // Energy difference between the
      two levels , eV
5
6 printf("\nThe difference between upper and lower
      energy levels for the most prominent wavelength
      = %5.3f eV", E);
7
8 // Result
9 // The difference between upper and lower energy
      levels for the most prominent wavelength = 0.397
      eV
```

---

**Scilab code Exa 8.2** Frequency and wavelength of carbon dioxide laser

```

1 // Scilab Code Ex8.2:: Page-8.8 (2009)
2 clc; clear;
3 E = 0.121; // Energy difference between the two
    levels , eV
4 lambda = 12400/E; // Wavelength of the radiation ,
    angstrom
5 f = 3e+08/(lambda*1e-010); // Frequency of the
    radiation , Hz
6
7 printf("\nThe wavelength of the radiation = %8.1f
    angstrom", lambda);
8 printf("\nThe frequency of the radiation = %4.2e Hz"
    , f);
9
10 // Result
11 // The wavelength of the radiation = 102479.3
    angstrom
12 // The frequency of the radiation = 2.93e+13 Hz

```

---

**Scilab code Exa 8.3** Energy of one emitted photon and total energy available per laser pulse

```

1 // Scilab Code Ex8.3:: Page-8.8 (2009)
2 clc; clear;
3 lambda = 7000; // Wavelength of the Ruby laser ,
    angstrom
4 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
5 N = 2.8e+019; // Total number of photons
6 E = 12400/lambda; // Energy of one emitted photon ,
    eV
7 E_p = E*e*N; // Total energy available per laser
    pulse , joule
8
9 printf("\nThe energy of one emitted photon = %4.2e J
    ", E*e);

```

```

10 printf("\nThe total energy available per laser pulse
      = %4.2f joule", E_p);
11
12 // Result
13 // The energy of one emitted photon = 2.83e-19 J
14 // The total energy available per laser pulse = 7.94
      joule

```

---

#### Scilab code Exa 8.4 Relative population of levels in Ruby laser

```

1 // Scilab Code Ex8.4:: Page-8.9 (2009)
2 clc; clear;
3 lambda = 7000; // Wavelength of the emitted light ,
      angstrom
4 k = 8.6e-005; // Boltzmann constant , eV/K
5 dE = 12400/lambda; // Energy difference of the
      levels , eV
6 T = [300 500]; // Temperatures of first and second
      states , K
7 for i = 1:1:2
8     N2_ratio_N1 = exp(-(dE/(k*T(i)))); // Relative
      population
9     printf("\nThe relative population at %d K = %3.1
      e", T(i), N2_ratio_N1);
10 end
11
12 // Result
13 // The relative population at 300 K = 1.5e-30
14 // The relative population at 500 K = 1.3e-18
15 // The answer is given wrong in the textbook for
      first part.

```

---

#### Scilab code Exa 8.5 Population of two states in He Ne laser

```

1 // Scilab Code Ex8.5:: Page-8.9 (2009)
2 clc; clear;
3 lambda = 7000; // Wavelength of the emitted light ,
    angstrom
4 k = 8.6e-005; // Boltzmann constant , eV/K
5 dE = 12400/lambda; // Energy difference of the
    levels , eV
6 T = 27+273; // Temperatures of the state , K
7 N2_ratio_N1 = exp(-(dE/(k*T))); // Relative
    population
8 printf("\nThe relative population of two states in
He-Ne laser at %d K = %3.1e", T, N2_ratio_N1);
9
10
11 // Result
12 // The relative population of two states in He-Ne
laser at 300 K = 1.5e-30
13 // The answer is given wrong in the textbook

```

---