

Scilab Textbook Companion for  
Applied Physics  
by P. K. Mittal<sup>1</sup>

Created by  
Mohit Garg  
B.Tech  
Electronics Engineering  
ABES Engineering College, Ghaziabad  
College Teacher  
None  
Cross-Checked by  
Lavitha Pereira

June 4, 2016

<sup>1</sup>Funded by a grant from the National Mission on Education through ICT,  
<http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab  
codes written in it can be downloaded from the "Textbook Companion Project"  
section at the website <http://scilab.in>

# **Book Description**

**Title:** Applied Physics

**Author:** P. K. Mittal

**Publisher:** I. K. International, New Delhi

**Edition:** 2

**Year:** 2006

**ISBN:** 978-81-89866-72-3

Scilab numbering policy used in this document and the relation to the above book.

**Exa** Example (Solved example)

**Eqn** Equation (Particular equation of the above book)

**AP** Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

# Contents

<b>List of Scilab Codes</b>	<b>4</b>
<b>1 Bonding in Solids</b>	<b>5</b>
<b>2 Crystal Structure</b>	<b>10</b>
<b>3 Planes in Crystals</b>	<b>15</b>
<b>4 Crystal Diffraction</b>	<b>17</b>
<b>5 Principles of Quantum Mechanics</b>	<b>26</b>
<b>6 Electron Theory and Band Theory of Metals</b>	<b>48</b>
<b>7 Dielectric Properties</b>	<b>58</b>
<b>8 Magnetic Properties</b>	<b>67</b>
<b>9 Semiconductors</b>	<b>73</b>
<b>10 Superconductivity</b>	<b>88</b>
<b>12 Fibre Optics</b>	<b>93</b>

# List of Scilab Codes

Exa 1.3	calculate potential energy . . . . .	5
Exa 1.4	calculate bond energy for NaCl . . . . .	6
Exa 1.5	calculate compressibility . . . . .	6
Exa 1.6	calculate ionic cohesive energy and atomic cohesive energy . . . . .	7
Exa 1.7	calculate contribution per ions to the cohesive energy . . . . .	8
Exa 2.1	calculate lattice constant . . . . .	10
Exa 2.2	calculate distance between two nearest Cu atoms . . . . .	10
Exa 2.3	calculate lattice constant . . . . .	11
Exa 2.4	calculate lattice constant . . . . .	12
Exa 2.5	calculate distance between adjacent atoms in NaCl . . . . .	12
Exa 2.6	calculate packing fraction and density . . . . .	13
Exa 3.11	calculate bond energy for NaCl . . . . .	15
Exa 3.12	calculate lattice spacing . . . . .	15
Exa 3.15	calculate density . . . . .	16
Exa 4.1	Find spacing constant . . . . .	17
Exa 4.2	Find glancing angle . . . . .	18
Exa 4.3	Find glancing angle . . . . .	18
Exa 4.4	Find wavelength and maximum order possible . . . . .	19
Exa 4.5	Find wavelength in XU . . . . .	20
Exa 4.6	find wavelength and energy . . . . .	21
Exa 4.7	calculate interplanar spacing . . . . .	22
Exa 4.8	calculate angle of first order diffraction maximum . . . . .	22
Exa 4.9	Show that given angles are successive order of diffraction and find spacing constant . . . . .	23
Exa 5.1	Find velocity and kinetic energy . . . . .	26
Exa 5.2	Calculate deBroglie wavelength . . . . .	27
Exa 5.3	Calculate wavelength . . . . .	27

Exa 5.4	Calculate deBroglie wavelength . . . . .	28
Exa 5.5	Calculate wavelength . . . . .	29
Exa 5.6	Calculate wavelength for photon and electron . . . . .	29
Exa 5.7	Calculate velocity and kinetic energy . . . . .	30
Exa 5.8	Calculate deBroglie wavelength . . . . .	31
Exa 5.9	Calculate deBroglie wavelength of proton . . . . .	32
Exa 5.10	Calculate wavelength . . . . .	33
Exa 5.11	Calculate glancing angle . . . . .	33
Exa 5.12	Calculate spacing of crystal . . . . .	34
Exa 5.13	Calculate velocity of electron . . . . .	35
Exa 5.14	Calculate uncertainty in the momentum and uncertainty in the velocity . . . . .	35
Exa 5.15	Compare uncertainty in the velocity of electron and proton . . . . .	36
Exa 5.16	Calculate minimum uncertainty in the momentum and minimum kinetic energy of proton . . . . .	37
Exa 5.17	Calculate percentage of uncertainty in the momentum of electron . . . . .	38
Exa 5.18	Calculate uncertainty in the position of electron . . . . .	39
Exa 5.19	Calculate uncertainty in the position of X ray photon . . . . .	40
Exa 5.20	Compare minimum uncertainty in the frequency of the photon . . . . .	41
Exa 5.21	Calculate uncertainty in the energy of the photon . . . . .	42
Exa 5.22	Calculate minimum error in the energy . . . . .	42
Exa 5.23	Calculate energy corresponding to the 2nd and 4th quantum states . . . . .	43
Exa 5.24	Calculate energy corresponding to the ground and first two excited states . . . . .	44
Exa 5.25	Calculate minimum uncertainty in the velocity of electron . . . . .	45
Exa 5.26	Calculate minimum energy of electron . . . . .	46
Exa 6.1	Calculate mean free path of electron . . . . .	48
Exa 6.2	Calculate the temperature . . . . .	49
Exa 6.3	Calculate relaxation time of conduction electrons . . . . .	50
Exa 6.4	Calculate mean free path traveeled by the electrons . . . . .	50
Exa 6.5	Calculate relaxation time of conduction electrons . . . . .	51
Exa 6.6	Calculate average kinetic energy and velocity of molecules . . . . .	52
Exa 6.7	Calculate velocity of electron and proton . . . . .	52

Exa 6.8	Calculate drift velocity of free electrons . . . . .	53
Exa 6.9	Calculate average drift velocity of electrons . . . . .	54
Exa 6.10	Calculate mobility of electrons . . . . .	55
Exa 6.11	Calculate average energy of free electron at 0K and corresponding temperature for a classical particle or an ideal gas . . . . .	55
Exa 6.12	Calculate electron density for a metal . . . . .	56
Exa 7.1	Calculate the capacitance of capacitor and charge on the plates . . . . .	58
Exa 7.2	Calculate the resultant voltage across the capacitor . .	59
Exa 7.3	Calculate the radius of electron cloud and displacement	60
Exa 7.4	Calculate the dipole moment induced in each atom and atomic polarisability . . . . .	61
Exa 7.5	Calculate the electronic polarisability of sulphur . . . .	62
Exa 7.6	Calculate the electronic polarisability of Helium atoms	62
Exa 7.7	Calculate the dielectric constant of the material . . . .	63
Exa 7.8	Calculate the atomic polarisability of sulphur . . . . .	64
Exa 7.9	calculate polarisability due to permanent dipole moment and due to deformation of the molecules . . . . .	64
Exa 7.10	determine the percentage of ionic polarisability in sodium crystal . . . . .	66
Exa 8.1	Calculate intensity of magnetism and magnetic flux density . . . . .	67
Exa 8.2	Calculate change in magnetic moment . . . . .	68
Exa 8.3	Calculate relative permeability of a ferromagnetic material . . . . .	68
Exa 8.4	Calculate magnetic force and relative permeability . .	69
Exa 8.5	Calculate current through the solenoid . . . . .	70
Exa 8.6	Calculate flux density magnetic intensity and relative permeability . . . . .	70
Exa 8.7	Calculate Hysteresis loss per cycle . . . . .	71
Exa 9.2	Find the temperature at which number of electrons becomes 10 times . . . . .	73
Exa 9.3	find the resistance of intrinsic germanium . . . . .	74
Exa 9.4	find the electrical conductivity and resistivity of germanium . . . . .	75
Exa 9.5	find the equilibrium hole concentration and conductivity	76
Exa 9.6	find intrinsic conductivity and doping conductivity . .	76

Exa 9.7	find density of hole carriers at room temperature . . . . .	78
Exa 9.8	find intrinsic carrier density and conductivity at 300K in germanium . . . . .	78
Exa 9.9	Find the energy band gap . . . . .	79
Exa 9.10	Find the electron and hole concentrations and the resistivity . . . . .	80
Exa 9.11	Find the conductivity of p type Ge crystal . . . . .	81
Exa 9.12	Find the diffusion coefficient of electron in silicon . . . . .	82
Exa 9.13	Find the probability of occupation of lowest level in conduction band . . . . .	82
Exa 9.14	Find the ratio of conductivity at 600K and at 300K . . . . .	84
Exa 9.15	Find the electron and hole densities and conductivity and the resistance . . . . .	84
Exa 9.16	Find the mobility and density . . . . .	85
Exa 9.17	Find the mobility and density of charge carrier . . . . .	86
Exa 10.1	Calculate magnitude of critical magnetic field . . . . .	88
Exa 10.2	Calculate critical current value . . . . .	88
Exa 10.3	calculate isotropic mass . . . . .	89
Exa 10.4	calculate transition temperature . . . . .	90
Exa 10.5	calculate critical current at 5K . . . . .	90
Exa 10.6	calculate frequency of EM waves . . . . .	91
Exa 10.7	calculate transition temperature of the isotopes . . . . .	92
Exa 12.1	calculate fractional index change for a given optical fibre . . . . .	93
Exa 12.2	calculate numerical aperture and the acceptance angle of an optical fibre . . . . .	93
Exa 12.3	calculate the acceptance angle of an optical fibre . . . . .	94
Exa 12.4	calculate refractive index of material of the core . . . . .	95
Exa 12.5	calculate numerical aperture acceptance angle and the critical angle of the optical fibre . . . . .	95
Exa 12.6	calculate refractive index of the core and cladding material of a fibre . . . . .	96
Exa 12.7	calculate numerical aperture and acceptance angle of the symmetrical fibre . . . . .	97
Exa 12.8	calculate numerical aperture and acceptance angle of the symmetrical fibre . . . . .	98

# Chapter 1

## Bonding in Solids

Scilab code Exa 1.3 calculate potential energy

```
1 //chapter 1
2 //example 1.3
3 //calculate potential energy
4 //page 15
5 clear;
6 clc;
7 //given
8 r=2; //in angstrom(distance)
9 e=1.6E-19; // in C (charge of electron)
10 E_o= 8.85E-12; // absolute permittivity
11 //calculate
12 r=2*1E-10; // since r is in angstrom
13 V=-e^2/(4*pi*E_o*r); // calculate potential
14 printf('\nThe potential energy is \tV=%3.3E J',V);
15 V=V/e; // changing to eV
16 printf ('\nIn electron-Volt ,\tV=%0.2f eV',V);
17 // Note: the answer in the book is wrong due to
    calculation mistake
```

---

### Scilab code Exa 1.4 calculate bond energy for NaCl

```
1 //chapter 1
2 //example 1.4
3 //calculate bond energy for NaCl
4 //page 15–16
5 clear;
6 clc;
7 //given
8 r0=0.236; //in nanometer(interionic distance)
9 e=1.6E-19; // in C (charge of electron)
10 E_o= 8.85E-12;// absolute premittivity
11 N=8; // Born constant
12 IE=5.14;// in eV (ionisation energy of sodium)
13 EA=3.65;// in eV (electron affinity of Chlorine)
14 pi=3.14; // value of pi used in the solution
15 //calculate
16 r0=r0*1E-9; // since r is in nanometer
17 PE=(e^2/(4*pi*E_o*r0))*(1-1/N); // calculate
    potential energy
18 PE=PE/e; //changing unit from J to eV
19 printf ('\nThe potential energy is \tPE=%.2f eV',PE);
20 NE=IE-EA;// calculation of Net energy
21 printf ('\nThe net energy is \tNE=%.2f eV',NE);
22 BE=PE-NE;// calculation of Bond Energy
23 printf ('\nThe bond energy is \tBE=%.2f eV',BE);
24 // Note: (1)–In order to make the answer practically
    feasible and avoid the unusual answer, I have
    used r_0=0.236 nm instead of 236 nm. because
    using this value will give very much irrelevant
    answer.
25 // (2) There is slight variation in the answer
    due to round off.
```

---

### Scilab code Exa 1.5 calculate compressibility

```

1 //chapter 1
2 //example 1.5
3 //calculate compressibility
4 //page 16
5 clear;
6 clc;
7 //given
8 r_0=.41; //in mm(lattice constant)
9 e=1.6E-19; // in C (charge of electron)
10 E_o= 8.85E-12;// absolute permittivity
11 n=0.5; // repulsive exponent value
12 alpha=1.76; // Madelung constant
13 pi=3.14; // value of pi used in the solution
14 //calculate
15 r=.41*1E-3; // since r is in mm
16 Beta=72*pi*E_o*r^4/(alpha*e^2*(n-1)); // calculation
    compressibility
17 printf ('\nThe compressibility is \tBeta=%1.2E ',Beta
        );
18 // Note: the answer in the book is wrong due to
    calculation mistake

```

---

**Scilab code Exa 1.6** calculate ionic cohesive energy and atomic cohesive energy

```

1 //chapter 1
2 //example 1.6
3 //calculate ionic cohesive energy and atomic
    cohesive energy
4 //page 16
5 clear;
6 clc;
7 //given
8 r_0=3.56; // in Angstrom
9 e=1.6E-19; // in C (charge of electron)

```

```

10 IE=3.89; //in eV (ionisation energy of Cs)
11 EA=-3.61; // in eV (electron affinity of Cl)
12 n=10.5; // Born constant
13 E_o= 8.85E-12; // absolute permittivity
14 alpha=1.763; // Madelung constant
15 pi=3.14; // value of pi used in the solution
16 //calculate
17 r_0=r_0*1E-10; // since r is in nanometer
18 U=-alpha*(e^2/(4*pi*E_o*r_0))*(1-1/n); // calculate
    potential energy
19 U=U/e; //changing unit from J to eV
20 printf ('\nThe ionic cohesive energy is \t%.2f eV',U);
21 ACE=U+EA+IE; // calculation of atomic cohesive
    energy
22 printf ('\nThe atomic cohesive energy is \t%.2f eV',
    ACE);

```

---

**Scilab code Exa 1.7** calculate contribution per ions to the cohesive energy

```

1 //chapter 1
2 //example 1.7
3 //calculate contribution per ions to the cohesive
    energy
4 //page 17
5 clear;
6 clc;
7 //given
8 r_0=2.81; // in Angstrom
9 e=1.6E-19; // in C (charge of electron)
10 n=9; // Born constant
11 E_o= 8.85E-12; // absolute permittivity
12 alpha=1.748; // Madelung constant
13 pi=3.14; // value of pi used in the solution
14 //calculate
15 r_0=r_0*1E-10; // since r is in nanometer

```

```
16 V=-alpha*(e^2/(4*pi*E_o*r_0))*(1-1/n); // calculate
      potential energy
17 V=V/e; // changing unit from J to eV
18 printf('\nThe potential energy is \tV=%f eV',V);
19 V_1=V/2; // Since only half of the energy contribute
      per ion to the cohesive energy therefore
20 printf('\nThe energy contributing per ions to the
      cohesive energy is \t%2f eV',V_1);
21 // Note: Answer in the book is wrong due to
      calculation mistake
```

---

# Chapter 2

## Crystal Structure

**Scilab code Exa 2.1** calculate lattice constant

```
1 //chapter 2
2 //example 2.1
3 //calculate lattice constant
4 //page 40–41
5 clear;
6 clc;
7 //given
8 N=6.02E26; // in /Kg-molecule (Avogadro's number)
9 n=4; // number of molecules per unit cell of NaCl
10 M=58.5; // in Kg/Kg-molecule (molecular weight of
    NaCl)
11 p=2189; // in Kg/m^3 (density)
12 //calculate
13 a=nthroot((n*M/(N*p)),3);
14 printf ('\nThe lattice constant is \ta=%1.2E m',a);
15 a=a*1E10; // changing unit to Angstrom
16 printf ('\n\t\ta=%2f Angstrom',a);
```

---

**Scilab code Exa 2.2** calculate distance between two nearest Cu atoms

```

1 //chapter 2
2 //example 2.2
3 //calculate distance between two nearest Cu atoms
4 //page 41
5 clear;
6 clc;
7 //given
8 N=6.02E23; // in /gram-atom (Avogadro's number)
9 n=4; // number of atom per unit cell for fcc
       structure
10 M=63.5; //in gram/gram-atom (atomic weight of Cu)
11 p=8.96; // in g/cm^3 (density)
12 //calculate
13 a=nthroot((n*M/(N*p)),3);
14 printf ('\nThe lattice constant is \ta=%1.2E cm',a);
15 a=a*1E8; // changing unit from cm to Angstrom
16 printf ('\n\t\ta=%f Angstrom',a);
17 d=a/sqrt(2); // distance infcc lattice
18 printf ('\nThe distance between two nearest Cu atoms
           is \td=%f Angstrom',d);

```

---

### Scilab code Exa 2.3 calculate lattice constant

```

1 //chapter 2
2 //example 2.3
3 //calculate lattice constant
4 //page 41-42
5 clear;
6 clc;
7 //given
8 N=6.02E26; // in /Kg-atom (Avogadro's number)
9 n=2; // number of molecules per unit cell for bcc
       lattice
10 M=55.85; // in Kg/Kg-atom (atomic weight of Iron)
11 p=7860; // in Kg/m^3 (density)

```

```
12 // calculate
13 a=nthroot((n*M/(N*p)),3);
14 printf ('\nThe lattice constant is \ta=%1.3E m',a);
15 a=a*1E10; // changing unit to Angstrom
16 printf ('\n\t\ta=%f Angstrom',a);
```

---

### Scilab code Exa 2.4 calculate lattice constant

```
1 //chapter 2
2 //example 2.4
3 //calculate lattice constant
4 //page 42
5 clear;
6 clc;
7 //given
8 N=6.02E26; // in /Kg-atom (Avogadro's number)
9 n=2; // number of molecules per unit cell for bcc
       lattice
10 M=6.94; // in Kg/Kg-atom (atomic weight of Iron)
11 p=530; // in Kg/m^3 (density)
12 //calculate
13 a=nthroot((n*M/(N*p)),3);
14 printf ('\nThe lattice constant is \ta=%1.3E m',a);
15 a=a*1E10; // changing unit to Angstrom
16 printf ('\n\t\ta=%f Angstrom',a);
```

---

### Scilab code Exa 2.5 calculate distance between adjacent atoms in NaCl

```
1 //chapter 2
2 //example 2.5
3 //calculate distance between adjacent atoms in NaCl
4 //page 42-43
5 clear;
```

```

6 clc;
7 //given
8 N=6.02E23; // in /gram-molecule (Avogadro's number)
9 M=58.5; //in gram/gram-molecule (atomic weight of
NaCl)
10 p=2.17; // in g/cm^3 (density)
11 // calculate
12 // since V=M/p
13 // (1/d)^-3=2N/V=2Np/M
14 // therefore d= (M/2Np)^-3
15 d=nthroot((M/(2*N*p)),3);
16 printf ('\nThe distance between two adjacent atoms of
NaCl is \td=%1.2E cm',d);
17 d=d*1E8; // changing unit from cm to Angstrom
18 printf ('\n\t\t\t\t\t\t\td=%f Angstrom',d);

```

---

**Scilab code Exa 2.6** calculate packing fraction and density

```

1 //chapter 2
2 //example 2.6
3 //calculate packing fraction and density
4 //page 43
5 clear;
6 clc;
7 //given
8 r_Na=0.98; // in Angstrom (radius of sodium ion)
9 r_Cl=1.81; // in Angstrom (radius of chloride ion)
10 M_Na=22.99; // in amu (atomic mass of sodium)
11 M_Cl=35.45; // in amu (atomic mass of chlorine)
12 //calculate
13 a=2*(r_Na+r_Cl); // lattice parameter
14 printf ('\nLattice constant is \ta=%f Angstrom',a)
;
15 //PF=volume of ions present in the unit cell/volume
of unit cell

```

```
16 PF=((4*(4/3)*%pi)*r_Na^3+(4*(4/3)*%pi)*r_Cl^3)/a^3;
17 printf ('\nPacking fraction is %.3f',PF);
18 //Density=mass of unit cell/volume of unit cell
19 p=4*(M_Na+M_Cl)*1.66E-27/(a*1E-10)^3;
20 printf ('\nDensity is \tp=%.f Kg/m^3',p);
21 p=p*1E-3; //changing unit to gm/cm^-3
22 printf ('\nDensity is \tp=%.2f g/cm^3',p);
```

---

# Chapter 3

## Planes in Crystals

**Scilab code Exa 3.11** calculate bond energy for NaCl

```
1 //chapter 3
2 //example 3.11
3 //calculate interplanar spacing
4 //page 61
5 clear;
6 clc;
7 //given
8 h=3,k=2,l=1; // miller indices
9 a=4.2E-8; // in cm (lattice constant)
10 //calculate
11 d=a/sqrt(h^2+k^2+l^2); // calculation for
    interplanar spacing
12 printf('nThe interplanar spacing is\td=%1.2E cm',d)
    ;
13 d=d*1E8; //changing unit from cm to Angstrom
14 printf('n\t\td=%f Angstrom',d);
```

---

**Scilab code Exa 3.12** calculate lattice spacing

```

1 //chapter 3
2 //example 3.12
3 //calculate lattice spacing
4 //page 61
5 clear;
6 clc;
7 //given
8 h=1,k=1,l=1; // miller indices
9 a=2.5,b=2.5,c=1.8; // in Angstrom (lattice constants
for tetragonal lattice )
10 //calculate
11 d=1/sqrt((h/a)^2+(k/b)^2+(l/c)^2); // calculation
for interplanar spacing
12 printf ('\nThe lattice spacing is \t d=%f Angstrom ',d
);

```

---

### Scilab code Exa 3.15 calculate density

```

1 //chapter 3
2 //example 3.15
3 //calculate density
4 //page 63
5 clear;
6 clc;
7 //given
8 h=1,k=0,l=0; // miller indices
9 a=2.5; // in Angstrom (lattice constant)
10 //calculate
11 a=a*1E-10; //hence a is in Angstrom
12 d=a/sqrt(h^2+k^2+l^2); // calculation for
interplanar spacing
13 p=d/a^3;
14 printf ('\nThe density is \t p=%1.1E lattice points/m^2
',p);

```

---

# Chapter 4

## Crystal Diffraction

**Scilab code Exa 4.1** Find spacing constant

```
1 //chapter 4
2 //example 4.1
3 //Find spacing constant
4 //page 75
5 clear;
6 clc;
7 //given
8 lambda=2.6; // in Angstrom (wavelength)
9 theta=20; // in Degree (angle)
10 n=2;
11 //calculate
12 lambda=lambda*1E-10; // since lambda is in Angstrom
13 // Since  $2d\sin(\theta) = n(\lambda)$ 
14 // therefore  $d = n(\lambda) / 2 \sin(\theta)$ 
15 d=n*lambda/(2*sind(theta));
16 printf('\nThe spacing constant is \td=%1.1E m',d);
17 d=d*1E10; // changing unit from m to Angstrom
18 printf('\n\t\t\td=%1.1f Angstrom',d);
```

---

### Scilab code Exa 4.2 Find glancing angle

```
1 //chapter 4
2 //example 4.2
3 //Find glancing angle
4 //page 75
5 clear;
6 clc;
7 //given
8 h=1,k=1,l=0; //miller indices
9 a=0.26; // in nanometer (lattice constant)
10 lambda=0.065; // in nanometer (wavelength)
11 n=2; // order
12 //calculate
13 d=a/sqrt(h^2+k^2+l^2); // calculation of
   interlattice spacing
14 // Since 2dsin(theta)=n(lambda)
15 // therefore we have
16 theta=asind(n*lambda/(2*d));
17 printf('\nThe glancing angle is \t%.2f degree',theta
   );
18 //Note: there is slight variation in the answer due
   to round off
```

---

### Scilab code Exa 4.3 Find glancing angle

```
1 //chapter 4
2 //example 4.3
3 //Find glancing angle
4 //page 75–76
5 clear;
6 clc;
7 //given
8 d=3.04E-10; // in mm (spacing constant)
9 lambda=0.79; // in Angstrom (wavelength)
```

```

10 n=3; // order
11 //calculate
12 // Since  $2d\sin(\theta) = n(\lambda)$ 
13 // therefore we have
14 lambda=lambda*1E-10; //since lambda is in angstrom
15 theta=asind(n*lambda/(2*d));
16 printf ('\nThe glancing angle is \t%.3f degree',theta
);
17 //Note: In question the value of d=3.04E-9 cm but in
// solution is using d=3.04E-10 m.
18 // I have used d=3.04E-10 cm as used in the solution

```

---

**Scilab code Exa 4.4** Find wavelength and maximum order possible

```

1 //chapter 4
2 //example 4.4
3 //Find wavelength and maximum order possible
4 //page 76
5 clear;
6 clc;
7 //given
8 d=0.282; // in nanometer (spacing constant)
9 n=1; // order
10 theta=8.35; // in degree (glancing angle)
11 //calculate
12 d=d*1E-9; // since d is in nanometer
13 // Since  $2d\sin(\theta) = n(\lambda)$ 
14 // therefore we have
15 lambda=2*d*sind(theta)/n;
16 printf ('\nThe wavelength is \t%1.2E m',lambda);
17 lambda_1=lambda*1E10; //changing unit from m to
// Angstrom
18 printf ('\n\t\t\t= %.3f Angstrom',lambda_1);
19 theta_1=90; // in degree (for maximum order theta
=90)

```

```

20 n_max=2*d*sind(theta_1)/lambda; // calculation of
   maximum order .
21 printf ('\nThe maximum order possible is \tn=%f ,'
   n_max );
22 //Note: In question value of theta=8 degree and 35
   minutes but solution uses theta=8.35 degree
23 // I am using theta=8.35 degree

```

---

### Scilab code Exa 4.5 Find wavelength in XU

```

1 //chapter 4
2 //example 4.5
3 //Find wavelength in X.U.
4 //page 76-77
5 clear;
6 clc;
7 //given
8 theta=6; // in degree (glancing angle)
9 p=2170; // in Kg/m^3 (density)
10 M=58.46; // Molecular weight of NaCl
11 N=6.02E26; // in Kg-molecule (Avogadro's number)
12 n=1; // order
13 XU=1E-12; //since 1X.U.= 1E-12m
14 //calculate
15 d=(M/(2*N*p))^(1/3); //calclation of lattice constant
16 printf ('\nThe spacing constant is \td=%1.3E m',d);
17 // Since 2dsin(theta)=n(lambda)
18 // therefore we have
19 lambda=2*d*sind(theta)/n; //calculation of
   wavelength
20 printf ('\n\nThe wavelength is \t\t=%1.2E m',lambda);
21 lambda=lambda/XU;
22 printf ('\n\t\t\t\t\t=%1.1f X.U.',lambda);
23 // Note: The answer in the book is wrong due to
   calculation mistake

```

---

### Scilab code Exa 4.6 find wavelength and energy

```
1 //chapter 4
2 //example 4.6
3 //find wavelength and energy
4 //page 77
5 clear;
6 clc;
7 //given
8 h=1,k=1,l=1; // miller indices
9 a=5.63; // in Angstrom (lattice constant)
10 theta=27.5; // in degree (Glancing angle)
11 n=1; //order
12 H=6.625E-34; // in J-s (Plank's constant)
13 c=3E8; // in m/s (velocity of light)
14 e=1.6E-19; // charge of electron
15 //calculate
16 d=a/sqrt(h^2+k^2+l^2); // calculation for
    interplanar spacing
17 printf('\nThe lattice spacing is \td=%f Angstrom ',d
    );
18 // Since 2dsin(theta)=n(lambda)
19 // therefore we have
20 lambda=2*d*sind(theta)/n; // calculation for
    wavelength
21 printf('\nThe wavelength is \t=%f Angstrom ',lambda);
22 E=H*c/(lambda*1E-10); //calculation of Energy
23 printf('\nThe energy of X-rays is E=%1.3E J ',E);
24 E=E/e; // changing unit from J to eV
25 printf('\n\t\tE=%1.3E eV ',E);
26 // Note: c=3E8 m/s but in solution c=3E10 m/s has
    been used that's why answer is different
```

---

### Scilab code Exa 4.7 calculate interplanar spacing

```
1 //chapter 4
2 //example 4.7
3 //calculate interplanar spacing
4 //page 77–78
5 clear;
6 clc;
7 //given
8 V=344; // in V (accelerating voltage)
9 theta=60; // in degree (glancing angle)
10 m=9.1E-31; // in Kg (mass of electron)
11 h=6.625e-34; // in J-s (Plank's constant)
12 n=1; //order
13 e=1.6E-19; // charge on electron
14 //calculate
15 //Since K=m*v^2/2=e*V
16 // therefore v=sqrt(2*e*V/m)
17 // since lambda=h/(m*v)
18 //therefore we have lambda=h/sqrt(2*m*e*V)
19 lambda=h/sqrt(2*m*e*V); // calculation of lambda
20 printf ('\nThe wavelength is \t\t =%1.2E m',lambda);
21 lambda=lambda*1E10; //changing unit from m to
    Angstrom
22 printf ('\n\t\t\t\t =%.2 f Angstrom',lambda);
23 // Since 2dsin(theta)=n(lambda)
24 // therefore we have
25 d=n*lambda/(2*sind(theta));
26 printf ('\nThe interplanar spacing is \t d=% .2 f
    Angstrom',d);
```

---

### Scilab code Exa 4.8 calculate angle of first order diffraction maximum

```

1 //chapter 4
2 //example 4.8
3 //calculate angle of first order diffraction maximum
4 //page 78–79
5 clear;
6 clc;
7 //given
8 K=0.02; // in eV (kinetic energy)
9 d=2.0; // in Angstrom (Bragg's spacing)
10 m=1.00898; // in amu (mass of neutron)
11 amu=1.66E-27; // in Kg (1amu=1.66E-27 Kg)
12 h=6.625e-34; // in J-s (Plank's constant)
13 n=1; //order
14 e=1.6E-19; // charge on electron
15 //calculate
16 //Since  $K=m*v^2/2$ 
17 // therefore  $v=sqrt(2*K/m)$ 
18 // since  $\lambda=h/(m*v)$ 
19 // therefore we have  $\lambda=h/sqrt(2*m*K)$ 
20 m=m*amu; //changing unit from amu to Kg
21 K=K*e; //changing unit to J from eV
22 lambda=h/sqrt(2*m*K); // calculation of lambda
23 printf('\nThe wavelength is \t\t =%.1E m',lambda);
24 lambda=lambda*1E10; //changing unit from m to
    Angstrom
25 printf('\n\t\t\t\t =%.1f Angstrom',lambda);
26 // Since  $2d\sin(\theta)=n(\lambda)$ 
27 // therefore we have
28 theta=asin(n*lambda/(2*d)); // calculation of angle
    of first order diffraction maximum
29 printf('\nThe angle of first order diffraction
    maximum is %.f Degree',theta);

```

---

**Scilab code Exa 4.9** Show that given angles are successive order of diffraction and find spacing constant

```

1 //chapter 4
2 //example 4.9
3 //Show that given angles are successive order of
   diffraction and find spacing constant
4 //page 79
5 clear;
6 clc;
7 //given
8 lambda=0.586; // in Angstrom (wavelength of X-rays)
9 n1=1, n2=2, n3=3; // orders of diffraction
10 theta1=5+(58/60); // in degree (Glancing angle for
    first order of diffraction)
11 theta2=12+(01/60); //in degree (Glancing angle for
    second order of diffraction)
12 theta3=18+(12/60); //in degree (Glancing angle for
    third order of diffraction)
13 //calculate
14 K1=sind(theta1);
15 K2=sind(theta2);
16 K3=sind(theta3);
17 printf('The value of sine of different angle of
   diffraction is \nK1=%f\nK2=%f\nK3=%f',K1,K2,
   K3);
18 // Taking the ratios of K1:K2:K3
19 // We get K1:K2:K3=1:2:3
20 //Therefore we have
21 printf('\n\nOr we have \tK1:K2:K3=1:2:3');
22 printf('\nHence these angles of incidence are for
   Ist , 2nd and 3rd order reflections respectively')
   ;
23 // Since 2dsin(theta)=n(lambda)
24 // therefore we have
25 d1=n1*lambda/(2*K1);
26 d2=n2*lambda/(2*K2);
27 d3=n3*lambda/(2*K3);
28 d1=d1*1E-10; //changing unit from Angstrom to m
29 d2=d2*1E-10; //changing unit from Angstrom to m
30 d3=d3*1E-10; //changing unit from Angstrom to m

```

```
31 printf( '\n\nThe spacing constants are \nd1=%1.3E m\  
      nd2=%1.3E m\nd3=%1.3E m' ,d1 ,d2 ,d3 );  
32 d=(d1+d2+d3)/3;  
33 printf( '\n\nThe mean value of crystal spacing is d=  
      %1.3E m' ,d );
```

---

# Chapter 5

## Principles of Quantum Mechanics

**Scilab code Exa 5.1** Find velocity and kinetic energy

```
1 //chapter 5
2 //example 5.1
3 //Find velocity and kinetic energy
4 //page 102–103
5 clear;
6 clc;
7 //given
8 lambda=1; //in Angstrom (wavelength)
9 m=1.67E-27; // in Kg (mass of neutron)
10 h=6.625E-34; // in J-s (Planck's constant)
11 e=1.6E-19; // in C (charge of electron)
12 //calculate
13 lambda=lambda*1E-10; //since lambda is in Angstrom
14 // Since lambda=h/(m*v)
15 // Therefore we have
16 v=h/(m*lambda); //calculation of velocity
17 printf('\nThe velocity is \t v=%1.2E m/s ',v);
18 K=m*v^2/2; //calculation of kinetic energy
19 printf('\nThe kinetic energy is \tK=%1.2E J ',K);
```

```
20 K=K/e; // changing unit from J to eV
21 printf ('\n\t\tt=%f eV',K);
22 //Note: Due to round off, there is slight variation
      in the answer
```

---

### Scilab code Exa 5.2 Calculate deBroglie wavelength

```
1 //chapter 5
2 //example 5.2
3 //Calculate de-Broglie wavelength
4 //page 103–104
5 clear;
6 clc;
7 //given
8 K=50; // in eV (Kinetic energy)
9 m0=9.1E-31; // in Kg (mass of electron)
10 h=6.625E-34; // in J-s (Planck's constant)
11 e=1.6E-19; // in C (charge of electron)
12 //calculate
13 K=K*e; //changing unit from eV to J
14 //Since K=m*v^2/2
15 // Therefore v=sqrt(2*K/m)
16 // Since lambda=h/(m*v)
17 // Therefore we have
18 lambda=h/sqrt(2*m0*K); //calculation of wavelength
19 printf ('\nThe wavelength is \t=%f m',lambda);
20 lambda=lambda*1E10; //changing unit from m to
      Angstrom
21 printf ('\n\t\tt=%f Angstrom',lambda);
```

---

### Scilab code Exa 5.3 Calculate wavelength

```
1 //chapter 5
```

```

2 //example 5.3
3 //Calculate wavelength
4 //page 104
5 clear;
6 clc;
7 //given
8 E=2000; // in eV (Kinetic energy)
9 m=9.1E-31; // in Kg (mass of electron)
10 h=6.625E-34; // in J-s (Planck's constant)
11 e=1.6E-19; // in C (charge of electron)
12 //calculate
13 E=E*e; //changing unit from eV to J
14 //Since E=m*v^2/2
15 // Therefore v=sqrt(2*E/m)
16 // Since lambda=h/(m*v)
17 // Therefore we have
18 lambda=h/sqrt(2*m*E); //calculation of wavelength
19 printf ('\nThe wavelength is \t=%1.3E m',lambda);
20 lambda=lambda*1E9; //changing unit from m to
    nanometer
21 printf ('\n\t\t\t\t=%.4 f nm',lambda);

```

---

### Scilab code Exa 5.4 Calculate deBroglie wavelength

```

1 //chapter 5
2 //example 5.3
3 //Calculate de-Broglie wavelength
4 //page 104
5 clear;
6 clc;
7 //given
8 m_e=9.1E-31; // in Kg (mass of electron)
9 m_n=1.676E-27; // in Kg (mass of neutron)
10 h=6.625E-34; // in J-s (Planck's constant)
11 c=3E8; // in m/s (velocity of light)

```

```

12 // calculate
13 E_e=m_e*c^2; // rest mass energy of electron
14 E_n=2*E_e; // given (kinetic energy of neutron)
15 // Since K=m*v^2/2
16 // Therefore v=sqrt(2*K/m)
17 // Since lambda=h/(m*v)
18 // Therefore we have
19 lambda=h/sqrt(2*m_n*E_n); // calculation of
    wavelength
20 printf ('\nThe wavelength is \t=%1.1E m',lambda);
21 lambda=lambda*1E10; // changing unit from m to
    Angstrom
22 printf ('\n\t\t\t\t=%1.1E Angstrom',lambda);

```

---

### Scilab code Exa 5.5 Calculate wavelength

```

1 // chapter 5
2 //example 5.4
3 //Calculate wavelength
4 //page 104
5 clear;
6 clc;
7 //given
8 V=1600; // in V ( Potential )
9 // calculate
10 lambda=12.27/sqrt(V); // calculation of wavelength
    in Angstrom
11 printf ('\nThe wavelength is \t=%.3f Angstrom',lambda)
    ;
12 // Note: The answer in the book is wrong due to
    calculation mistake

```

---

### Scilab code Exa 5.6 Calculate wavelength for photon and electron









```

6  clc;
7  //given
8 V=100; // in V ( Potential)
9 n=1; // order of diffraction
10 d=2.15; // in Angstrom (lattice spacing)
11 //calculate
12 lambda=12.27/sqrt(V); // calculation of wavelength
    in Angstrom
13 printf ('\nThe wavelength is \t=%.3f Angstrom',lambda)
    ;
14 // Since 2*d*sind(theta)=n*lambda
15 //therefore we have
16 theta=asind(n*lambda/(2*d)); // calculation of
    glancing angle
17 printf ('\nThe glancing angle is \t=%.1f degree',theta)
    );
18 // Note: In question V=100 eV but the solution is
    using V=100V in the book and I have also used V
    =100V

```

---

### Scilab code Exa 5.12 Calculate spacing of crystal

```

1 //chapter 5
2 //example 5.12
3 //Calculate spacing of crystal
4 //page 107
5 clear;
6 clc;
7 //given
8 V=344; // in V ( Potential)
9 n=1; // order of diffraction
10 theta=60; // in degree (glancing angle)
11 //calculate
12 lambda=12.27/sqrt(V); // calculation of wavelength
    in Angstrom

```

```

13 printf ('\nThe wavelength is \t\t=%f Angstrom ,  

14 lambda);  

14 // Since 2*d*sind(theta)=n*lambda  

15 // therefore we have  

16 d=n*lambda/(2*sind(theta)); // calculation of  

    spacing constant  

17 printf ('\nThe spacing of the crystal is \td=%f  

    Angstrom ',d);

```

---

**Scilab code Exa 5.13** Calculate velocity of electron

```

1 //chapter 5  

2 //example 5.13  

3 //Calculate velocity of electron  

4 //page 107-108  

5 clear;  

6 clc;  

7 //given  

8 r=0.53E-10; // in m (radius of first Bohr orbit)  

9 h=6.6E-34; // in J-s (Planck's constant)  

10 m=9.1E-31; // in Kg (mass of electron)  

11 n=1; // First Bohr orbit  

12 pi=3.14; // value of pi used in the solution  

13 //calculate  

14 // Since 2*pi*r=n*lambda and lambda=h/(m*v)  

15 //Therefore we have v=h*n/(2*pi*r*m)  

16 v=h*n/(2*pi*r*m); //calculation of velocity  

17 printf ('\nThe velocity of electron is \tv=%f m/s ',  

    v);

```

---

**Scilab code Exa 5.14** Calculate uncertainty in the momentum and uncertainty in the velocity

```

1 //chapter 5
2 //example 5.14
3 //Calculate uncertainty in the momentum and
   uncertainty in the velocity
4 //page 108
5 clear;
6 clc;
7 //given
8 dx=0.2; // in Angstrom (uncertainty in the position)
9 h=6.6E-34; // in J-s (Planck's constant)
10 m0=9.1E-31; // in Kg (mass of electron)
11 pi=3.14; // value of pi used in the solution
12 //calculate
13 dx=dx*1E-10; //since dx is in Angstrom
14 // Since  $dx \cdot dp = h / 4 \pi$  (uncertainty relation)
15 dp=h/(4*pi*dx); // calculation of uncertainty in the
   momentum
16 printf('\nThe uncertainty in the momentum is \t dp=%1
   .2E Kg-m/s',dp);
17 //since  $dp = m \cdot dv$ 
18 dv=dp/m0; // calculation of uncertainty in the
   velocity
19 printf('\nThe uncertainty in the velocity is \t dv=%1
   .2E m/s',dv);

```

---

**Scilab code Exa 5.15** Compare uncertainty in the velocity of electron and proton

```

1 //chapter 5
2 //example 5.15
3 //Compare uncertainty in the velocity of electron
   and proton
4 //page 108
5 clear;
6 clc;

```

```

7 // given
8 m_e=9.1E-31; // in Kg (mass of electron)
9 m_p=1.67E-27; // in Kg (mass of proton)
10 dx_p=1; // in nanometer (uncertainty in position of
    electron)
11 dx_n=1; // in nanometer (uncertainty in position of
    proton)
12 // calculate
13 // since  $dp = h/(4\pi dx)$ 
14 // since  $h/(4\pi)$  is constant and  $dx$  is same for
    electron and proton
15 // therefore both electron and proton have same
    uncertainty in the momentum
16 // since  $dv = dp/m$  and  $dp$  is same for both
17 // therefore  $dv_e/dv_p = m_p/m_e$ 
18 // therefore
19 K=m_p/m_e; // ratio of uncertainty in the velocity
    of electron and proton
20 printf('\nThe ratio of uncertainty in the velocity
    of electron to that of proton is \t=%f',K);

```

---

**Scilab code Exa 5.16** Calculate minimum uncertainty in the momentum and minimum kinetic energy of proton

```

1 // chapter 5
2 //example 5.16
3 //Calculate minimum uncertainty in the momentum and
    minimum kinetic energy of proton
4 //page 108–109
5 clear;
6 clc;
7 //given
8 dx=5E-15; // in m (radius of nucleus or uncertainty
    in the position)
9 h=6.6E-34; // in J-s (Planck's constant)

```



```

12 pi=3.14; // value of pi used in the solution
13 e=1.6E-19; // in C (charge of electron)
14 //calculate
15 dx=dx*1E-10; // since dx is in Angstrom
16 // Since dx*dp=h/4*pi (uncertainty relation)
17 dp=h/(4*pi*dx); // calculation of uncertainty in the
    momentum
18 printf('\nThe uncertainty in the momentum of
        electron is \tdp=%1.2E Kg-m/s ',dp);
19 K=K*1E3*1.6E-19; // changing unit from KeV to J
20 p=sqrt(2*m*K); // calculation of momentum
21 printf('\nThe momentum of electron is \t\t\t p=%1.2E
        Kg-m/s ',p);
22 poc=(dp/p)*100; // calculation of percentage of
    uncertainty
23 printf('\nThe percentage of uncertainty in the
        momentum is =%1f ',poc);

```

---

**Scilab code Exa 5.18** Calculate uncertainty in the position of electron

```

1 //chapter 5
2 //example 5.18
3 //Calculate uncertainty in the position of electron
4 //page 109–110
5 clear;
6 clc;
7 //given
8 v=6.6E4; // m/s (speed of electron)
9 poc=0.01; // percentage of uncertainty
10 h=6.63E-34; // in J-s (Planck's constant)
11 m=9E-31; // in Kg (mass of electron)
12 pi=3.14; // value of pi used in the solution
13 //calculate
14 p=m*v; // calculation of momentum
15 printf('\nThe momentum of electron is \t\t\t p=%1.2E

```

```

    Kg-m/s',p);
16 dp=(poc/100)*p; // calculation of uncertainty in the
                     momentum
17 printf('\nThe uncertainty in the momentum of
          electron is \tdp=%1.2E Kg-m/s',dp);
18 // Since dx*dp=h/4*pi (uncertainty relation)
19 dx=h/(4*pi*dp); // calculation of uncertainty in the
                     position
20 printf('\nThe uncertainty in the position of
          electron is \tdx=%1.2E Kg-m/s',dx);
21 // Note; solution is incomplete in the book

```

---

**Scilab code Exa 5.19** Calculate uncertainty in the position of X ray photon

```

1 //chapter 5
2 //example 5.19
3 //Calculate uncertainty in the position of X-ray
   photon
4 //page 111-112
5 clear;
6 clc;
7 //given
8 lambda=1; // in Angstrom (wavelength)
9 pi=3.14; // value of pi used in the solution
10 dlambd=1E-6; // uncertainty in wavelength
11 //calculate
12 lambda=lambda*1E-10; // sinc lambda is in Angstrom
13 // By uncertainty principle , dx*dp>=h/(4*pi) --(1)
14 // since p=h/lambda ----- (2)
15 // Or p*lambda=h
16 // differentiting this equation
17 // p*dlambda+lambda*dp=0
18 // dp=-p*dlambda/lambda ----- (3)
19 //from (2) and (3) dp=-h*dlambda/lambda^2 ----- (4)

```

```

20 // from (1) and (4) dx*dlambda>=lambda^2/4*pi
21 // Or dx=lambda^2/(4*pi*dlambda)
22 dx=lambda^2/(4*pi*dlambda); // calculation of
    uncertainty in the position
23 printf('\nThe uncertainty in the position of X-ray
    photon is \tdx=%1.0E m',dx);
24 // Note: 1. In the question , wavelength accuracy is
    given as 1 in 1E8 but in book solution has used 1
    in 1E6 and same has been used by me.
25 //          2. ANSWER IS WRONG DUE TO CALCULATION
    MISTAKE

```

---

**Scilab code Exa 5.20** Compare minimum uncertainty in the frequency of the photon

```

1 //chapter 5
2 //example 5.20
3 //Compare minimum uncertainty in the frequency of
    the photon
4 //page 111
5 clear;
6 clc;
7 //given
8 dt=1E-8; // in sec (average life time)
9 pi=3.14; // value of pi used in the solution
10 //calculate
11 // Since dE*dt>=h/(4*pi) (uncertainty relation for
    energy)
12 // and E=h*v      v is the frequency
13 // therefore we have dv>=1/(4*pi*dt)
14 dv=1/(4*pi*dt); // calculation of minimum
    uncertainty in the frequency
15 printf('\nThe minimum uncertainty in the frequency
    of the photon is \tdv=%1.1E sec^-1',dv);

```

---





```

    corresponding to the 2nd quantum state
17 printf ('\nThe energy corresponding to the 2nd
        quantum state is \tE2=%1.3E J',E2);
18 E2=E2/e; //changing unit from J to eV
19 printf ('\n\t\t\t\t\t\t\t\t\t\t =%1.4E eV',E2);
20 E4=n4^2*h^2/(8*m*a^2); // calculation of energy
    corresponding to the 4nd quantum state
21 printf ('\nThe energy corresponding to the 4nd
        quantum state is \tE4=%1.3E J',E4);
22 E4=E4/e; //changing unit from J to eV
23 printf ('\n\t\t\t\t\t\t\t\t\t\t =%1.4E eV',E4);
24 // Note: The answer in the book is wrong due to
    calculation mistake

```

---

**Scilab code Exa 5.24** Calculate energy corresponding to the ground and first two excited states

```

1 //chapter 5
2 //example 5.24
3 //Calculate energy corresponding to the ground and
    first two excited states
4 //page 113
5 clear;
6 clc;
7 //given
8 a=1E-10; // in m (width of the well)
9 m=9.1E-31; // in Kg (mass of electron)
10 h=6.626E-34; // in J-s (Planck's constant)
11 n1=1, n2=2, n3=3; // ground and first two excited
    states
12 e=1.6*1E-19; // in C (charge of electron)
13 //calculate
14 // Since E_n=n^2*h^2/(8*m*a^2) (Energy
    corresponding to nth quantum state)
15 E1=n1^2*h^2/(8*m*a^2); // calculation of energy

```

```

        corresponding to the Ground state
16 printf ('\nThe energy corresponding to the ground
           state is \tE1=%1.3E J',E1);
17 E1=E1/e; //changing unit from J to eV
18 printf ('\n\t\t\t\t\t\t\t\t\t\t =%.2f eV',E1);
19 E2=n2^2*h^2/(8*m*a^2); // calculation of energy
   corresponding to the 1st excited state
20 printf ('\nThe energy corresponding to the 1st
           excited state is \tE2=%1.3E J',E2);
21 E2=E2/e; //changing unit from J to eV
22 printf ('\n\t\t\t\t\t\t\t\t\t\t =%.2f eV',E2);
23 E3=n3^2*h^2/(8*m*a^2); // calculation of energy
   corresponding to the 2nd excited state
24 printf ('\nThe energy corresponding to the 2nd
           excited state is \tE3=%1.3E J',E3);
25 E3=E3/e; //changing unit from J to eV
26 printf ('\n\t\t\t\t\t\t\t\t\t\t =%.2f eV',E3);
27 // Note: There is slight variation in the answer due
       to round off

```

---

**Scilab code Exa 5.25** Calculate minimum uncertainty in the velocity of electron

```

1 //chapter 5
2 //example 5.25
3 //Calculate minimum uncertainty in the velocity of
   electron
4 //page 113
5 clear;
6clc;
7 //given
8 dx=1E-8; // in m (length of box or uncertainty in
   the position)
9 h=6.626E-34; // in J-s (Planck's constant)
10 m=9.1E-31; // in Kg (mass of electron)

```



19 // Note: The answer in the book corresponding to J  
is wrong due to printing error.

---

# Chapter 6

## Electron Theory and Band Theory of Metals

**Scilab code Exa 6.1** Calculate mean free path of electron

```
1 //chapter 6
2 //example 6.1
3 //Calculate mean free path of electron
4 //page 146
5 clear;
6 clc;
7 //given
8 n=8.5E28; // in 1/m^3 (density of electron)
9 m_e=9.11E-31; // in Kg (mass of electron)
10 k=1.38E-23; // in J/K (Boltzmann's constant)
11 e=1.6E-19; // in C (charge of electron)
12 T=300; // in K (temperature)
13 p=1.69E-8; // in ohm-m (resistivity)
14 //calculate
15 lambda=sqrt(3*k*m_e*T)/(n*e^2*p); // calculation of
   mean free path
16 printf('\nThe mean free path of electron is \t=%1.2E
   m',lambda);
17 lambda=lambda*1E9; // changing unit from m to
```

```
    nanometer
18 printf ('\n\t\t\t\t\t\t\t\t\t\tt=% .2 f nm' , lambda);
19 // Note: answer in the book is wrong due to printing
        mistake
```

---

### Scilab code Exa 6.2 Calculate the temperature

```
1 //chapter 6
2 //example 6.2
3 //Calculate the temperature
4 //page 146
5 clear;
6 clc;
7 //given
8
9 k=1.38E-23; // in J/K (Boltzmann's constant)
10 e=1.6E-19; // in C (charge of electron)
11 P_E=1; // in percentage (probability that a state
        with an energy 0.5 eV above Fermi energy will be
        occupied)
12 E=0.5; // in eV (energy above Fermi level)
13 //calculate
14 P_E=1/100; // changing percentage into ratio
15 E=E*e; // changing unit from eV to J
16 // P_E=1/(1+exp((E-E_F)/k*T))
17 // Rearranging this equation, we get
18 // T=(E-E_F)/k*log((1/P_E)-1)
19 // Since E-E_F has been denoted by E therefore
20 T=E/(k*log((1/P_E)-1));
21 printf ('\nThe temperature is \tT=% .2 f K' , T);
22 // Note: There is slight variation in the answer due
        to logarithm function
```

---

### Scilab code Exa 6.3 Calculate relaxation time of conduction electrons

```
1 //chapter 6
2 //example 6.3
3 //Calculate relaxation time of conduction electrons
4 //page 147
5 clear;
6 clc;
7 //given
8 n=5.8E28; // in 1/m^3 (density of electron)
9 m=9.1E-31; // in Kg (mass of electron)
10 e=1.6E-19; // in C (charge of electron)
11 p=1.54E-8; // in ohm-m (resistivity)
12 //calculate
13 t=m/(n*e^2*p); // calculation of relaxation time
14 printf('\nThe relaxation time of conduction
electrons is %1.2E sec ',t);
```

---

### Scilab code Exa 6.4 Calculate mean free path traveeled by the electrons

```
1 //chapter 6
2 //example 6.4
3 //Calculate mean free path traveeled by the
electrons
4 //page 147
5 clear;
6 clc;
7 //given
8 n=8.5E28; // in 1/m^3 (density of electron)
9 m=9.1E-31; // in Kg (mass of electron)
10 e=1.6E-19; // in C (charge of electron)
11 sigma=6E7; // in 1/ohm-m (conductivity)
12 E_F=7; // in E=eV (Fermi energy of Copper)
13 //calculate
14 E_F=E_F*e; // changing unit from eV to J
```

```

15 v_F=sqrt(2*E_F/m); // calculation of velocity of
   electrons
16 printf('\nThe velocity of the electrons is \t\t\tv_F
   =%1.1E m/s ',v_F);
17 // Since sigma=n*e^2*lambda/(2*m*v_F)
18 // Therefore we have
19 lambda=2*m*v_F*sigma/(n*e^2); // calculation of mean
   free path
20 lambda=lambda*1E10; // changing unit from m to
   Angstrom
21 printf('\n\nThe mean free path traveled by the
   electrons is \t%.f Angstrom ',lambda);
22 // Note: Answer in the book is wrong due to the use
   of round-off value of v_F as calculated in the
   first part.

```

---

**Scilab code Exa 6.5** Calculate relaxation time of conduction electrons

```

1 //chapter 6
2 //example 6.5
3 //Calculate relaxation time of conduction electrons
4 //page 147–148
5 clear;
6 clc;
7 //given
8 n=6.5E28; // in 1/m^3 (density of electron)
9 m=9.1E-31; // in Kg (mass of electron)
10 e=1.6E-19; // in C (charge of electron)
11 p=1.43E-8; // in ohm-m (resistivity)
12 //calculate
13 t=m/(n*e^2*p); // calculation of relaxation time
14 printf('\nThe relaxation time of conduction
   electrons is %1.2E sec ',t);

```

---

**Scilab code Exa 6.6** Calculate average kinetic energy and velocity of molecules

```

1 //chapter 6
2 //example 6.6
3 //Calculate average kinetic energy and velocity of
molecules
4 //page 148
5 clear;
6clc;
7 //given
8 T=30; // in Celcius (temperature)
9 k=1.38E-23; // in J/K (Boltzmann's constant)
10 m_p=1.67E-27; // in Kg (mass of proton)
11 e=1.6E-19; // in C (charge of electron)
12 //calculate
13 T=T+273; // changing temperature from Celcius to
Kelvin
14 KE=(3/2)*k*T; // calculation of average kinetic
energy
15 printf ('\nThe average kinetic energy of gas ,\n
molecules is \tKE=%3.2E J ',KE);
16 KE=KE/e; // changing unit from J to eV
17 printf ('\n\t\t\t\t\t\t\t\t\t\t\t =%f eV ',KE);
18 m=1.008*2*m_p; // calculating mass of hydrogen gas
molecule
19 c=sqrt(3*k*T/m); // calculation of velocity
20 printf ('\n\nThe velocity of molecules is \tc=%.2f m/\n
s ',c);
21 // Note: There is calculation mistake in the answer
of energy given in eV and that of velocity

```

---

**Scilab code Exa 6.7** Calculate velocity of electron and proton

```

1 //chapter 6
2 //example 6.7
3 //Calculate velocity of electron and proton
4 //page 148–149
5 clear;
6 clc;
7 //given
8 E=10; // in eV (kinetic energy for each electron and
      proton)
9 m_e=9.1E-31; // in Kg (mass of electron)
10 m_p=1.67E-27; // in Kg (mass of proton)
11 e=1.6E-19; // in C (charge of electron)
12 //calculate
13 E=E*e; // changing unit from eV to J
14 // since  $E=m*v^2/2$ 
15 // therefore  $v=\sqrt{2E/m}$ 
16 v_e=sqrt(2*E/m_e); // calculation of kinetic energy
      of electron
17 printf('\nThe kinetic energy of electron is \tv_e=%1
      .3E m/s',v_e);
18 v_p=sqrt(2*E/m_p); // calculation of kinetic energy
      of proton
19 printf('\nThe kinetic energy of proton is \tv_p=%1.3
      E m/s',v_p);
20 // Note: The answer in the book for both kinetic
      energy of electron and that of proton is wrong
      due to calculation mistake

```

---

### Scilab code Exa 6.8 Calculate drift velocity of free electrons

```

1 //chapter 6
2 //example 6.8
3 //Calculate drift velocity of free electrons
4 //page 149
5 clear;

```

```

6 clc;
7 //given
8 I=100; // in A (current in the wire)
9 e=1.6E-19; // in C (charge of electron)
10 A=10; // in mm^2 (cross-sectional area)
11 n=8.5E28; // in 1/m^3 (density of electron)
12 //calculate
13 A=A*1E-6; // changing unit from mm^2 to m^2
14 v_d=I/(n*A*e);
15 printf('\nThe drift velocity of free electrons is \
tv_d=%1.3E m/s',v_d);

```

---

### Scilab code Exa 6.9 Calculate average drift velocity of electrons

```

1 //chapter 6
2 //example 6.9
3 //Calculate average drift velocity of electrons
4 //page 149
5 clear;
6 clc;
7 //given
8 I=4; // in A (current in the conductor)
9 e=1.6E-19; // in C (charge of electron)
10 A=1E-6; // in m^2 (cross-sectional area)
11 N_A=6.02E23; // in atoms/gram-atom (Avogadro's
    number)
12 p=8.9; // in g/cm^3 (density)
13 M=63.6; // atomic mass of copper
14 //calculate
15 n=N_A*p/M; // Calculation of density of electrons in
    g/cm^3
16 printf('\nThe density of copper atoms is \tn=%1.2E
    atoms/m^3',n);
17 n=n*1E6; // changing unit from g/cm^3 to g/m^3
18 printf('\n\t\t\t\t=%1.2E atoms/m^3',n);

```

```
19 v_d=I/(n*A*e);
20 printf ('\n\nThe average drift velocity of free
          electrons is \tv_d=%1.1E m/s',v_d);
```

---

**Scilab code Exa 6.10** Calculate mobility of electrons

```
1 //chapter 6
2 //example 6.10
3 //Calculate mobility of electrons
4 //page 149–150
5 clear;
6 clc;
7 //given
8 n=9E28; // in 1/m^3 (density of valence electrons)
9 sigma=6E7; // in mho/m (conductivity of copper)
10 e=1.6E-19; // in C (charge of electron)
11 //calculate
12 // Since sigma=n*e*mu therefore
13 mu=sigma/(n*e); // calculation of mobility of
                     electron
14 printf ('\n\nThe mobility of electrons is \t%1.2E m
          ^2/V-s',mu);
```

---

**Scilab code Exa 6.11** Calculate average energy of free electron at 0K and corresponding temperature for a classical particle or an ideal gas

```
1 //chapter 6
2 //example 6.11
3 //Calculate average energy of free electron at 0K
   and corresponding temperature for a classical
   particle (an ideal gas)
4 //page 150
5 clear;
```

```

6 clc;
7 //given
8 E_F=5.51; // in eV (Fermi energy in Silver)
9 k=1.38E-23; // in J/K (Boltzmann's constant)
10 e=1.6E-19; // in C (charge of electron)
11 //calculate
12 // part-(a)
13 Eo=(3/5)*E_F; // calculation of average energy of
    free electron at 0K
14 printf ('\n\nThe average energy of free electron at 0
    K is \tEo=%f eV',Eo);
15 // part-(b)
16 Eo=Eo*e; // changing unit from eV to J
17 // Since for a classical particle  $E=(3/2)*k*T$ 
18 // therefroe we have
19 T=(2/3)*Eo/k; // calculation of temperature for a
    classical particle (an ideal gas)
20 printf ('\n\nThe temperature at which a classical
    particle have this much energy is \t T=%1.3E K',T
);

```

---

### Scilab code Exa 6.12 Calculate electron density for a metal

```

1 //chapter 6
2 //example 6.12
3 //Calculate electron density for a metal
4 //page 150
5 clear;
6 clc;
7 //given
8 E_F_L=4.7; // in eV (Fermi energy in Lithium)
9 E_F_M=2.35; // in eV (Fermi energy in a metal)
10 n_L=4.6E28; // in  $1/m^3$  (density of electron in
    Lithium)
11 //calculate

```

```
12 // Since n=((2*m/h) ^ 3/2)*E_F^(3/2)*(8*pi/3) and all
   things except E_F are constant
13 // Therefore we have n=C*E_F^(3/2) where C is
   proportionality constant
14 // n1/n2=(E_F_1/E_F_2)^(3/2)
15 // Therefore we have
16 n_M=n_L*(E_F_M/E_F_L); // calculation of electron
   density for a metal
17 printf ('\nThe lectron density for a metal is \t=%1.1
   E 1/m^3 ', n_M);
18 //Note: Answer in the book is wrong due to priting
   error
```

---

# Chapter 7

## Dielectric Properties

**Scilab code Exa 7.1** Calculate the capacitance of capacitor and charge on the plates

```
1 //chapter 7
2 //example 7.1
3 //Calculate the capacitance of capacitor and charge
   on the plates
4 //page 187
5 clear;
6 clc;
7 //given
8 A=100; // in cm^2 (cross-sectional area)
9 d=1; // in cm (seperation between plates)
10 Eo=8.85E-12; // in F/m (absolute permittivity)
11 V=100; // in V (potential difference)
12 //calculate
13 A=A*1E-4; // changing unit from cm^2 to m^2
14 d=d*1E-2; // changing unit from cm to m
15 C=Eo*A/d; // calculation of capacitance
16 Q=C*V; // calculation of charge
17 printf ('\nThe capacitance of capacitor is \t C=%1.2E
          C',C);
18 C=C*1E12; // changing unit of capacitance from F to
```

```
pF  
19 printf ('\n\t\t\t\t\t\t\t\t =%.2f pF',C);  
20 printf ('\n\nThe charge on the plates is \t\t Q=%1.2E  
                              C',Q);  
-----
```

**Scilab code Exa 7.2** Calculate the resultant voltage across the capacitor

```
1 //chapter 7  
2 //example 7.2  
3 //Calculate the resultant voltage across the  
                capacitor  
4 //page 187  
5 clear;  
6 clc;  
7 //given  
8 A=650; // in mm^2 (cross-sectional area)  
9 d=4; // in mm (seperation between plates)  
10 Eo=8.85E-12; // in F/m (absolute permittivity)  
11 Er=3.5; // di-electric constant of the material  
12 Q=2E-10; // in C (charge on plates)  
13 //calculate  
14 A=A*1E-6; // changing unit from mm^2 to m^2  
15 d=d*1E-3; // changing unit from mm to m  
16 C=Er*Eo*A/d; // calculation of capacitance  
17 V=Q/C; // calculation of charge  
18 printf ('\nThe capacitance of capacitor is \t C=%1.2E  
                              C',C);  
19 C=C*1E12; // changing unit of capacitance from F to  
                    pF  
20 printf ('\n\t\t\t\t\t\t\t\t\t\t =%.2f pF',C);  
21 printf ('\n\nThe resultant voltage across the  
                capacitor is \t V=%1.2f V',V);  
22 // NOTE: The answer is wrong due to calculation  
                mistake. The mistake is that in the book Value of  
                cross-sectional area and seperation
```

23 // between plates is considered in cm and dielectric constant has not been considered.

---

**Scilab code Exa 7.3** Calculate the radius of electron cloud and displacement

```
1 //chapter 7
2 //example 7.3
3 //Calculate the radius of electron cloud and
   displacement
4 //page 188
5 clear;
6 clc;
7 //given
8 N=2.7E25; // in 1/m^3 (density of atoms)
9 E=1E6; // in V/m (electric field)
10 Z=2; // atomic number of Helium
11 Eo=8.85E-12; // in F/m (absolute permittivity)
12 Er=1.0000684; // (dielectric constant of the
   material)
13 e=1.6E-19; // in C (charge of electron)
14 pi=3.14; // value of pi used in the solution
15 //calculate
16 // since alpha=Eo*(Er-1)/N=4*pi*Eo*r_0^3
17 // Therefore we have r_0^3=(Er-1)/(4*pi*N)
18 r_0=((Er-1)/(4*pi*N))^(1/3); // calculation of radius
   of electron cloud
19 printf('\nThe radius of electron cloud is \t r_0=%1
   .2E m',r_0);
20 x=4*pi*Eo*E*r_0/(Z*e); // calculation of
   displacement
21 printf('\n\nThe displacement is x=%1.2E m',x);
22 // NOTE: The answer is wrong due to calculation
   mistake.
```

---

**Scilab code Exa 7.4** Calculate the dipole moment induced in each atom and atomic polarisability

```
1 //chapter 7
2 //example 7.4
3 //Calculate the dipole moment induced in each atom
   and atomic polarisability
4 //page 188–189
5 clear;
6 clc;
7 //given
8 K=1.000134; // di-elecrtic constant of the neon gas
   at NTP
9 E=90000; // in V/m (electric field)
10 Eo=8.85E-12; // in C/N·m^2 (absolute premittivity)
11 N_A=6.023E26; // in atoms/Kg–mole (Avogadro's number
   )
12 V=22.4; // in m^3 (volume of gas at NTP
13 //calculate
14 n=N_A/V; // calculaton of density of atoms
15 // Since P=n*p=(k-1)*Eo*E
16 // therefore we have
17 p=(K-1)*Eo*E/n; // calculation of dipole moment
   induced
18 printf('\nThe dipole moment induced in each atom is
   \tp=%1.2E C·m',p);
19 alpha=p/E; // calculation of atomic polarisability
20 printf('\n\nThe atomic polarisability of neon is \t=
   %1.2E c·m^2/V',alpha);
21 // NOTE: The answer of atomic polarisability is
   wrong due to printing error
```

---

**Scilab code Exa 7.5** Calculate the electronic polarisability of sulphur

```
1 //chapter 7
2 //example 7.5
3 //Calculate the electronic polarisability of sulphur
4 //page 189
5 clear;
6 clc;
7 //given
8 Er=3.75; // di-elecrtic constant of sulphur at 27
degree Celcius
9 gama=1/3; // internal field constant
10 p=2050; // in Kg/m^3 (density)
11 M_A=32; // in amu (atomic weight of sulphur)
12 Eo=8.85E-12; // in F/m (absolute permittivity)
13 N=6.022E23; // Avogadro's number
14 //calculate
15 // Since ((Er-1)/(Er+2))*(M_A/p)=(N/(3*Eo))*alpha_e
16 // therefore we have
17 alpha_e=((Er-1)/(Er+2))*(M_A/p)*(3*Eo/N); //
calculation of electronic polarisability of
sulphur
18 printf('\nThe electronic polarisability of sulphur
is \t=%1.2E Fm^2',alpha_e);
19 // NOTE: There is slight variation in the answer due
to round off
```

---

**Scilab code Exa 7.6** Calculate the electronic polarisability of Helium atoms

```
1 //chapter 7
2 //example 7.6
3 //Calculate the electronic polarisability of Helium
atoms
4 //page 189-190
5 clear;
```

```

6 clc;
7 //given
8 Er=1.0000684; // di-electric constant of Helium gas
      at NTP
9 Eo=8.85E-12; // in F/m (absolute permittivity)
10 N=2.7E25; // number of atoms per unit volume
11 // calculate
12 // Since  $Er-1 = (N/Eo) * \alpha_e$ 
13 // therefore we have
14 alpha_e=Eo*(Er-1)/N; // calculation of electronic
      polarisability of Helium
15 printf ('\nThe electronic polarisability of Helium
      gas is \t=%1.2E Fm^2 ',alpha_e);
16 // NOTE: There is slight variation in the answer due
      to round off

```

---

**Scilab code Exa 7.7** Calculate the dielectric constant of the material

```

1 //chapter 7
2 //example 7.7
3 //Calculate the dielectric constant of the material
4 //page 190
5 clear;
6 clc;
7 //given
8 N=3E28; // in atoms/m^3 (density of atoms)
9 alpha_e=1E-40; // in F-m^2 (electronic
      polarisability)
10 Eo=8.85E-12; // in F/m (absolute permittivity)
11 // calculate
12 // Since  $(Er-1)/(Er+2) = N * \alpha_e / (3 * Eo)$ 
13 // therefore we have
14 Er=(2*(N*alpha_e/(3*Eo))+1)/(1-(N*alpha_e/(3*Eo)));
15 // calculation of dielectric constant of the
      material

```

```
16 printf ('\nThe dielectric constant of the material is  
    \tEr=%f F/m',Er);  
17 // NOTE: The answer in the book is wrong due to  
    calculation mistake
```

---

**Scilab code Exa 7.8** Calculate the atomic polarisability of sulphur

```
1 //chapter 7  
2 //example 7.8  
3 //Calculate the atomic polarisability of sulphur  
4 //page 190  
5 clear;  
6 clc;  
7 //given  
8 Er=4; // relative permittivity of sulphur  
9 Eo=8.85E-12; // in F/m (absolute permittivity)  
10 NA=2.08E3; // in Kg/m^3 (density of atoms in sulphur  
    )  
11 //calculate  
12 // Since ((Er-1)/(Er+2))*(M_A/p)=(N/(3*Eo))*alpha_e  
13 // therefore we have  
14 alpha_e=((Er-1)/(Er+2))*(3*Eo/NA); // calculation of  
    electronic polarisability of sulphur  
15 printf ('\nThe electronic polarisability of sulphur  
    is \t=%f Fm^2',alpha_e);  
16 // NOTE: The answer in the book is wrong due to  
    calculation mistake. Also one point to be  
    mentioned is that wrong formula has been used in  
    the solution but i have used the formula as used  
    in the solution.
```

---

**Scilab code Exa 7.9** calculate polarisability due to permanent dipole moment and due to deformation of the molecules

```

1 // chapter 7
2 // example 7.9
3 // calculate polarisability due to permanent dipole
   moment and due to deformation of the molecules
4 // page 190–191
5 clear;
6 clc;
7 // given
8 alpha1=2.5E-39; // in C^2–m/N (dielectric constant
   at 300K)
9 alpha2=2.0E-39; // in C^2–m/N (dielectric constant
   at 400K)
10 T1=300; // in K(first temperature)
11 T2=400; // in K(second temperature)
12 //calculate
13 // since alpha=alpha_d+alpha0 and alpha0=Beta/T
14 // therefore alpha=alpha_d+(Beta/T)
15 // since alpha1=alpha_d+(Beta/T1) and alpha2=
   alpha_d+(Beta/T2)
16 // therefore alpha1–alpha2=Beta*((1/T1)–(1/T2))
17 // or Beta= (alpha1–alpha2)/ ((1/T1)–(1/T2))
18 Beta= (alpha1–alpha2)/ ((1/T1)–(1/T2)); //
   calculation of Beta
19 alpha_d=alpha1–(Beta/T1); // calculation of
   polarisability due to defromation
20 alpha0_1=Beta/T1; // calculation of polarisability
   due to permanent dipole moment at 300K
21 alpha0_2=Beta/T2; // calculation of polarisability
   due to permanent dipole moment at 400K
22 printf('\nThe polarisability due to permanent dipole
   moment at 300K is \t %1.2E C^2–m/N',alpha0_1);
23 printf('\nThe polarisability due to permanent dipole
   moment at 400K is \t %1.2E C^2–m/N',alpha0_2);
24 printf('\n\nThe polarisability due to deformation of
   the molecules is \t %1.2E C^2–m/N',alpha_d);

```

---

**Scilab code Exa 7.10** determine the percentage of ionic polarisability in sodium crystal

```
1 // chapter 7
2 // example 7.10
3 // determine the percentage of ionic polarisability
   in sodium crystal
4 // page 191–192
5 clear;
6 clc;
7 // given
8 n=1.5; // refractive index
9 Er=5.6; // dielectric constant
10 //calculate
11 // since  $(Er-1)/(Er+2)=N*(\alpha_e+\alpha_i)/(3*E_0)$ 
   Clausius–Mossotti equation
12 // and  $(n^2-1)/(n^2+2)=N*\alpha_e/(3*E_0)$ 
13 // from above two equations , we get  $((n^2-1)/(n^2+2)) * ((Er+2)/(Er-1)) = \alpha_e / (\alpha_e + \alpha_i)$ 
14 // or  $\alpha_i / (\alpha_e + \alpha_i) = 1 - ((n^2-1)/(n^2+2)) * ((Er+2)/(Er-1)) = (\text{say } P)$ 
15 // where P is fractional ionisational polarisability
16 P=1-((n^2-1)/(n^2+2))*((Er+2)/(Er-1)); //
   calculation of fractional ionisational
   polarisability
17 P=P*100; // calculation of percentage of
   ionisational polarisability
18 printf('\nThe percentage of ionisational
   polarisability is \t%.1f percent ',P);
```

---

# Chapter 8

## Magnetic Properties

**Scilab code Exa 8.1** Calculate intensity of magnetism and magnetic flux density

```
1 //chapter 8
2 //example 8.1
3 //Calculate intensity of magnetism and magnetic flux
   density
4 //page 236
5 clear;
6 clc;
7 //given
8 X=-0.5E-5; // magnetic susceptibility of silicon
9 H=0.9E4; // in A/m (magnetic field intensity)
10 mu0=4*pi*1E-7; // in H/m (absolute permeability)
11 //calculate
12 I=X*H; // calculation of intensity of magnetism
13 printf('\nThe intensity of magnetism is \tI=%.3f A/m
   ',I);
14 B=mu0*H*(1+X); // calculation of magnetic flux
   density
15 printf('\nThe magnetic flux density is \tB=%^.3f Wb/m
   ^2 ',B);
16 // Note: The answer in the book is wrong. This is
```

because the value of H given in the question is  $H = 0.9E4$  A/m but in the solution the value of H that has been used is  $H=9.9E4$  A/m.

---

### Scilab code Exa 8.2 Calculate change in magnetic moment

```
1 //chapter 8
2 //example 8.2
3 //Calculate change in magnetic moment
4 //page 236
5 clear;
6 clc;
7 //given
8 r=0.052; // in nm (radius of orbit)
9 B=1; // in Wb/m^2 (magnetic field of induction)
10 e=1.6E-19; // in C (charge of electron)
11 m=9.1E-31; // in Kg (mass of electron)
12 //calculate
13 r=0.052*1E-9; // changing unit from nm to m
14 d_mu=(e^2*r^2*B)/(4*m); // calculation of change in
    magnetic moment
15 printf ('\nThe change in magnetic moment is \t%1.4E
    Am^2',d_mu);
16 // Note: The answer in the book is wrong due to
    caluclation mistake
```

---

### Scilab code Exa 8.3 Calculate relative permeability of a ferromagentic material

```
1 //chapter 8
2 //example 8.3
3 //Calculate relative permeability of a ferromagentic
    material
```

```

4 //page 236
5 clear;
6 clc;
7 //given
8 H=220; // in A/m (magnetic field intensity)
9 I=3300; // in A/m (intensity of magnetisation)
10 //calculate
11 mu_r=1+(I/H); // calculation of relative
    permeability
12 printf ('\nThe relative permeability of a
    ferromagnetic material is %.f',mu_r);

```

---

**Scilab code Exa 8.4** Calculate magnetic force and relative permeability

```

1 //chapter 8
2 //example 8.4
3 //Calculate magnetic force and relative permeability
4 //page 236–237
5 clear;
6 clc;
7 //given
8 I=3000; // in A/m (intensity of magnetisation)
9 B=0.005; // in Wb/m^2 (magnetic flux intensity)
10 pi=3.14; // value of pi used in the solution
11 mu0=4*pi*1E-7; // in H/m (absolute permeability)
12 //calculate
13 H=(B/mu0)-I; // calculation of magnetic force
14 printf ('\nThe magnetic force is \tH=%f',H);
15 mu_r=(I/H)+1; // calculation of relative
    permeability
16 printf ('\nThe relative permeability is \t%.3f',mu_r)
;
```

---

**Scilab code Exa 8.5** Calculate current through the solenoid

```
1 //chapter 8
2 //example 8.5
3 //Calculate current through the solenoid
4 //page 237
5 clear;
6 clc;
7 //given
8 H=4E3; // in A/m (magnetic field intensity)
9 N=60; // number of turns
10 l=12; // in cm (length of solenoid)
11 //calculate
12 n=N/(l*1E-2); // calculation of number of turns per
    unit metre
13 // Snice H=n*i;
14 i=H/n; // calculation of current through the
    solenoid
15 printf('\nThe current through the solenoid is \ti=%.
    f A',i);
```

---

**Scilab code Exa 8.6** Calculate flux density magnetic intensity and relative permeability

```
1 //chapter 8
2 //example 8.6
3 //Calculate flux density , magnetic intensity and
    relative permeability
4 //page 237
5 clear;
6 clc;
7 //given
8 l=30; // in cm (length of solenoid)
9 A=1; // in cm^2 (cross-sectional area)
10 N=300; // number of turns
```

```

11 i=0.032; // in A (current through the winding)
12 phi_B=2E-6; // in Wb (magnetic flux)
13 pi=3.14; // value of pi used in the solution
14 mu0=4*pi*1E-7; // in H/m (absolute permeability)
15 //calculate
16 l=1*1E-2; // changing unit from cm to m
17 A=A*1E-4; // changing unit from cm^2 to m^2
18 B=phi_B/A; // calculation of flux density
19 printf ('\nThe flux density is \tB=%1.0E Wb/m^2',B);
20 H=N*i/l; // calculation of magnetic intensity
21 printf ('\nThe magnetic intensity is \tH=%f A-turns/
    m',H);
22 mu=B/H; // calcluation of absolute permeability of
    iron
23 mu_r=mu/mu0; // calcluation of relative permeability
    of iron
24 printf ('\nThe relative permeability of iron is \t%.
    f ',mu_r);
25 // Note: The value of relative permeability varies
    slightly due to the use of round off value mu as
    calculated

```

---

### Scilab code Exa 8.7 Calculate Hysteresis loss per cycle

```

1 //chapter 8
2 //example 8.7
3 //Calculate Hysteresis loss per cycle
4 //page 238
5 clear;
6 clc;
7 //given
8 A=100; // in m^2 (area of Hysteresis loop)
9 B=0.01; // in Wb/m^2 (unit space along vertical axis
    or magnetic flux density)
10 H=40; // in A/m (unit space along horizontal axis or

```

```
    magnetic field intensity)
11 // calculate
12 H_L=A*B*H; // calculation of magnetic intensity
13 printf('\nThe Hysteresis loss per cycle is %.f J/m^2
      ',H_L);
```

---

# Chapter 9

## Semiconductors

**Scilab code Exa 9.2** Find the temperature at which number of electrons becomes 10 times

```
1 // chapter 9
2 // example 9.2
3 // Find the temperature at which number of electrons
   becomes 10 times
4 // page 272
5 clear;
6 clc;
7 //given
8 Eg=0.67; // in eV (Energy band gap)
9 k=1.38E-23; // in J/K (Boltzmann's constant)
10 T1=298; // in K (room temperature)
11 e=1.6E-19; // in C (charge of electron)
12 K=10; // ratio of number of electrons at different
   temperature
13 // calculate
14 Eg=Eg*e; // changing unit from eV to Joule
15 // since ne=Ke*exp(-Eg/(2*k*T))
16 // and ne/ne1=exp(-Eg/(2*k*T))/exp(-Eg/(2*k*T1)) and
   ne/ne1=K=10
17 // therefore we have 10=exp(-Eg/(2*k*T))/exp(-Eg/(2*
```

```

        k*T1))
18 // re-arranging the equation for T, we get T2=1/((1/
    T1)-((2*k*log(10))/Eg))
19 T=1/((1/T1)-((2*k*log(10))/Eg)); // calculation of
    the temperature
20 printf ('\nThe temperature at which number of
    electrons in the conduction band of a
    semiconductor increases by a factor of 10 is \tT=
    %.f K',T);
21 // Note: there s slight variation in the answer due
    to round off calculation

```

---

### Scilab code Exa 9.3 find the resistance of intrinsic germanium

```

1 // chapter 9
2 // example 9.3
3 // find the resistance of intrinsic germanium
4 // page 272-273
5 // given
6 clear;
7 clc;
8 ni=2.5E13; // in /cm^3 (intrinsic carrier density)
9 ue=3900; // in cm^2/(V-s) (electron mobilities)
10 uh=1900; // in cm^2/(V-s) (hole mobilities)
11 e=1.6E-19; // in C (charge of electron)
12 l=1; // in cm (length of the box)
13 b=1,h=1; // in mm (dimensions of germanium rod )
14 // calculate
15 ni=ni*1E6; // changing unit from 1/cm^3 to 1/m^3
16 ue=ue*1E-4; // changing unit from cm^2 to m^2
17 uh=uh*1E-4; // changing unit from cm^2 to m^2
18 sigma=ni*e*(ue+uh); // calculation of conductivity
19 rho=1/sigma; // calculation of resistivity
20 l=l*1E-2; // changing unit from mm to m for length
21 A=(b*1E-3)*(h*1E-3); // changing unit from mm to m

```

```

        for width and height and calculation of cross-
        sectional area
22 R=rho*l/A; // calculation of resistance
23 printf ('\nThe resistance of intrinsic germanium is \
           tR=%1.1E ohm',R);

```

---

**Scilab code Exa 9.4** find the electrical conductivity and resistivity of germanium

```

1 // chapter 9
2 // example 9.4
3 // find the electrical conductivity and resistivity
   of germanium
4 // page 273
5 clear;
6 clc;
7 // given
8 ne=2.5E19; // in /m^3 (electron density)
9 nh=2.5E19; // in /m^3 (hole density)
10 ue=0.36; // in m^2/(V-s) (electron mobilities)
11 uh=0.17; // in m^2/(V-s) (hole mobilities)
12 e=1.6E-19; // in C (charge of electron)
13 // calculate
14 // since ne=nh=ni , therefore we have
15 ni=nh;
16 sigma=ni*e*(ue+uh); // calculation of conductivity
17 printf ('\nThe conductivity of germanium is %.2f /
           ohm-m',sigma);
18 rho=1/sigma; // calculation of resistivity
19 printf ('\nThe resistivity of germanium is %.2f ohm-
           m',rho);
20 // Note: In the question , the value of ni has been
   misprinted as 2.5E-19 /m^3 rather it should be
   2.5E19 /m^3. I have used 2.5E19 /m^3

```

---

**Scilab code Exa 9.5** find the equilibrium hole concentration and conductivity

```
1 // chapter 9
2 // example 9.5
3 // find the equilibrium hole concentration and
   conductivity
4 // page 273–274
5 clear;
6 clc;
7 // given
8 ni=1.5E16; // in /m^3 (intrinsic carrier density)
9 ue=0.135; // in m^2/(V-s) (electron mobilities)
10 uh=0.048; // in m^2/(V-s) (hole mobilities)
11 e=1.6E-19; // in C (charge of electron)
12 ND=1E23; // in atom/m^3 (doping concentration)
13 // calculate
14 sigma_i=ni*e*(ue+uh); // calculation of intrinsic
   conductivity
15 printf('\n\nThe intrinsic conductivity for silicon is
   %1.2E S',sigma_i);
16 sigma=ND*ue*e; // calculation of conductivity after
   doping
17 printf('\n\nThe conductivity after doping with
   phosphorus atoms is %1.2E S',sigma);
18 rho=ni^2/ND; // calculation of equilibrium hole
   concentration
19 printf('\n\nThe equilibrium hole concentration is
   %1.2E /m^3',rho);
```

---

**Scilab code Exa 9.6** find intrinsic conductivity and doping conductivity

```

1 // chapter 9
2 // example 9.6
3 // find intrinsic conductivity and doping
   conductivity
4 // page 274
5 clear;
6 clc;
7 // given
8 ni=1.5E16; // in /m^3 (intrinsic carrier density)
9 ue=0.13; // in m^2/(V-s) (electron mobilities)
10 uh=0.05; // in m^2/(V-s) (hole mobilities)
11 e=1.6E-19; // in C (charge of electron)
12 ne=5E20; // in /m^3 (concentration of donor type
   impurity)
13 nh=5E20; // in /m^3 (concentration of acceptor type
   impurity)
14 // calculate
15 // part-i
16 sigma=ni*e*(ue+uh); // calculation of intrinsic
   conductivity
17 printf ('\nThe intrinsic conductivity for silicon is
   %1.2E (ohm-m)^-1',sigma);
18 // part-ii
19 // since 1 donor atom is in 1E8 Si atoms, hence
   holes concentration can be neglected
20 sigma=ne*e*ue; // calculation of conductivity after
   doping with donor type impurity
21 printf ('\n\nThe conductivity after doping with donor
   type impurity is %.1f (ohm-m)^-1',sigma);
22 // part-iii
23 // since 1 acceptor atom is in 1E8 Si atoms, hence
   electron concentration can be neglected
24 sigma=nh*e*uh; // calculation of conductivity after
   doping with acceptor type impurity
25 printf ('\n\nThe conductivity after doping with
   acceptor type impurity is %.f (ohm-m)^-1',sigma)
   ;
26 // Note: In question the value of ne and nh has

```

been misprinted as 5E28 atoms/m<sup>3</sup> which is too big but the solution has used the correct value 5 E20 atoms/m<sup>3</sup>. I have also used this value.

---

**Scilab code Exa 9.7** find density of hole carriers at room temperature

```
1 // chapter 9
2 // example 9.7
3 // find density of hole carriers at room temperature
4 // page 274–275
5 clear;
6 clc;
7 // given
8 ni=1E20; // in /m^3 (intrinsic carrier density)
9 ND=1E21; // in /m^3 (donor impurity concentration)
10 // calculate
11 nh=ni^2/ND; // calculation of density of hole
    carriers at room temperature
12 printf('\nThe density of hole carriers at room
    temperature is \tnh=%1.0E /m^3 ',nh);
13 // Note: answer in the book is wrong due to printing
    mistake
```

---

**Scilab code Exa 9.8** find intrinsic carrier density and conductivity at 300K in germanium

```
1 // chapter 9
2 // example 9.8
3 // find intrinsic carrier density and conductivity
    at 300K in germanium
4 // page 275
5 clear;
6 clc;
```

```

7 M=72.6; // atomic mass of germanium
8 P=5400; // in Kg/m^3 (density)
9 ue=0.4; // in m^2/V-s (mobility of electrons)
10 uh=0.2; // in m^2/V-s (mobility of holes)
11 Eg=0.7; // in eV (Band gap)
12 m=9.1E-31; // in Kg (mass of electron)
13 k=1.38E-23; // in J/K (Boltzmann s constant)
14 T=300; // in K (temperature)
15 h=6.63E-34; // in J/s (Planck s constant)
16 pi=3.14; // value of pi used in the solution
17 e=1.6E-19; // in C(charge of electron)
18 // calculate
19 Eg=Eg*e; // changing unit from eV to J
20 ni=2*(2*pi*m*k*T/h^2)^(3/2)*exp(-Eg/(2*k*T));
21 printf ('\nThe intrinsic carrier density for
germanium at 300K is \tni=%1.1E /m^3',ni);
22 sigma=ni*e*(ue+uh);
23 printf ('\nThe conductivity of germanium is \t%1.2f (
ohm-m)^-1',sigma);
24 // Note: Answer in the book is wrong due to
calculation mistake

```

---

**Scilab code Exa 9.9** Find the energy band gap

```

1 // chapter 9
2 // example 9.9
3 // Find the energy band gap
4 // page 275
5 clear;
6clc;
7 //given
8 rho1=4.5;// in ohm-m (resistivity at 20 degree
Celcius)
9 rho2=2.0;// in ohm-m (resistivity at 32 degree
Celcius)

```

```

10 k=1.38E-23; // in J/K ( Boltzmann s constant)
11 T1=20, T2=32; // in degree Celcius (two temperatures
12 e=1.6E-19; // in C (charge of electron)
13 // calculate
14 T1=T1+273; // changing unit from degree Celcius to K
15 T2=T2+273; // changing unit from degree Celcius to K
16 // since sigma=e*u*C*T^(3/2)*exp(-Eg/(2*k*T))
17 // therefore sigma1/sigma2=(T1/T2)^(3/2)*exp((-Eg/(2*k
18 //)*(1/T1)-(1/T2)))
19 // and sigma=1/rho
20 // therefore we have rho2/rho1=(T1/T2)^(3/2)*exp((-Eg
21 // /(2*k)*(1/T1)-(1/T2)))
22 // re-arranging above equation for Eg, we get Eg=(2*
23 k/((1/T1)-(1/T2)))*((3/2)*log(T1/T2)-log(rho2/
24 rho1));
25 Eg=(2*k/((1/T1)-(1/T2)))*((3/2)*log(T1/T2)-log(rho2/
26 rho1));
27 printf ('\nThe energy band gap is \tEg=%1.2E J ',Eg);
28 Eg=Eg/e;// changing unit from J to eV
29 printf ('\n\t\t=%0.2f eV ',Eg);

```

---

**Scilab code Exa 9.10** Find the electron and hole concentrations and the resistivity

```

1 // chapter 9
2 // example 9.10
3 // Find the electron and hole concentrations and the
3 // resistivity
4 // page 276
5 clear;
6 clc;
7 //given
8 rho=2300; // in ohm-m (resistivity of pure silicon)
9 ue=0.135; // in m^2/V-s (mobility of electron)

```

```

10 uh=0.048; // in m^2/V-s (mobility of electron)
11 Nd=1E19; // in /m^3 (doping concentration)
12 e=1.6E-19; // in C (charge of electron)
13 //calculate
14 // since sigma=ni*e*(ue+uh) and sigma=1/rho
15 // therefore ni=1/(rho*e*(ue+uh))
16 ni=1/(rho*e*(ue+uh)); // calculation of intrinsic
    concentration
17 ne=Nd; // calculation of electron concentration
18 printf ('\nThe electron concentration is \tne=%1.1E /
    m^3',ne);
19 nh=ni^2/Nd; // calculation of hole concentration
20 printf ('\nThe hole concentration is \tnh=%1.1E /m^3',
    nh);
21 sigma=ne*ue*e+nh*uh*e; // calculation of
    conductivity
22 rho=1/sigma; // calculation of resistivity
23 printf ('\nThe resistivity of the specimen is \t%.2f
    ohm-m',rho);

```

---

**Scilab code Exa 9.11** Find the conductivity of p type Ge crystal

```

1 // chapter 9
2 // example 9.11
3 // Find the conductivity of p-type Ge crystal
4 // page 276-277
5 clear;
6 clc;
7 //given
8 uh=1900; // in cm^2/V-s (mobility of electron)
9 Na=2E17; // in /m^3 (acceptor doping concentration)
10 e=1.6E-19; // in C(charge of electron)
11 //calculate
12 uh=uh*1E-4; // changing unit from cm^2/V-s to m^2/V-
    s

```

```

13 Na=Na*1E6; // changing unit from 1/cm^3 to 1/m^3
14 nh=Na; // hole concentration
15 // since sigma=ne*ue*e+nh*uh*e and nh>>ne
16 // therefore sigma=nh*uh*e
17 sigma=nh*uh*e; // calculation of conductivity
18 printf ('\nThe conductivity of p-type Ge crystal is \
t%.f /ohm-m',sigma);
19 // Note: there is slight variation in the answer due
to round off calculation

```

---

**Scilab code Exa 9.12** Find the diffusion coefficient of electron in silicon

```

1 // chapter 9
2 // example 9.12
3 // Find the diffusion co-efficient of electron in
silicon
4 // page 277
5 clear;
6 clc;
7 //given
8 ue=0.19; // in m^2/V-s (mobility of electron)
9 T=300; // in K (temperature)
10 k=1.38E-23; // in J/K (Boltzmann s constant)
11 e=1.6E-19; // in C(charge of electron)
12 //calculate
13 Dn=ue*k*T/e; // calculation of diffusion co-
efficient
14 printf ('\nThe diffusion co-efficient of electron in
silicon is \tDn=%1.1E m^2/s',Dn);

```

---

**Scilab code Exa 9.13** Find the probability of occupation of lowest level in conduction band

```

1 // chapter 9
2 // example 9.13
3 // Find the probability of occupation of lowest
   level in conduction band
4 // page 277–278
5 clear;
6 clc;
7 //given
8 Eg=0.4; // in eV (Band gap of semiconductor)
9 k=1.38E-23; // in J/K (Boltzmann s constant)
10 T1=0; // in degree Celcius (first temperature)
11 T2=50; // in degree Celcius (second temperature)
12 T3=100; // in degree Celcius (third temperature)
13 e=1.602E-19; //in C (charge of electron)
14 // calculate
15 T1=T1+273; // changing temperature form Celcius to
   Kelvin
16 T2=T2+273; // changing temperature form Celcius to
   Kelvin
17 T3=T3+273; // changing temperature form Celcius to
   Kelvin
18 Eg=Eg*e; // changing unit from eV to Joule
19 //Using F_E=1/(1+exp(Eg/2*k*T))
20 F_E1=1/(1+exp(Eg/(2*k*T1))); // calculation of
   probability of occupation of lowest level at 0
   degree Celcius
21 F_E2=1/(1+exp(Eg/(2*k*T2))); // calculation of
   probability of occupation of lowest level at 50
   degree Celcius
22 F_E3=1/(1+exp(Eg/(2*k*T3))); // calculation of
   probability of occupation of lowest level at 100
   degree Celcius
23 printf('\nThe probability of occupation of lowest
   level in conduction band is\n\n');
24 printf('\t\t at 0 degree Celcius , F_E=%1.3E eV\n',
   F_E1);
25 printf('\t\t at 50 degree Celcius , F_E=%1.2E eV\n',
   F_E2);

```

```
26 printf ('\t\t at 100 degree Celcius , F_E=%1.3E eV' ,  
F_E3);
```

---

**Scilab code Exa 9.14** Find the ratio of conductivity at 600K and at 300K

```
1 // chapter 9  
2 // example 9.14  
3 // Find the ratio of conductivity at 600K and at 300  
K  
4 // page 278  
5 clear;  
6 clc;  
7 //given  
8 Eg=1.2; // in eV (Energy band gap)  
9 k=1.38E-23; // in J/K (Boltzmann s constant)  
10 T1=600, T2=300; // in K (two temperatures)  
11 e=1.6E-19; // in C (charge of electron)  
12 // calculate  
13 Eg=Eg*e; // changing unit from eV to Joule  
14 // since sigma is proportional to exp(-Eg/(2*k*T))  
15 // therefore ratio=sigma1/sigma2=exp(-Eg/(2*k*((1/T1  
)-(1/T2))));  
16 ratio= exp((-Eg/(2*k))*((1/T1)-(1/T2))); //  
calculation of ratio of conductivity at 600K and  
at 300K  
17 printf ('\nThe ratio of conductivity at 600K and at  
300K is \t%1.2E',ratio);
```

---

**Scilab code Exa 9.15** Find the electron and hole densities and conductivity and the resistance

```
1 // chapter 9  
2 // example 9.15
```

```

3 // Find the electron and hole densities and
   conductivity and the resistance
4 // page 278–279
5 clear;
6 clc;
7 //given
8 ue=0.39; // in m^2/V-s (mobility of electron)
9 n=5E13; // number of donor atoms
10 ni=2.4E19; // in atoms/m^3 (intrinsic carrier
    density)
11 l=10; // in mm (length of rod)
12 a=1; // in mm (side of square cross-section)
13 e=1.6E-19; // in C (charge of electron)
14 //calculate
15 l=l*1E-3; // changing unit from mm to m
16 a=a*1E-3; // changing unit from mm to m
17 A=a^2; // calculation of cross-section area
18 Nd=n/(l*A); // calculation of donor concentration
19 ne=Nd; // calculation of electron density
20 nh=ni^2/Nd; // calculation of hole density
21 printf ('\nThe electron density is \tne=%1.0E /m^3 , ne);
22 printf ('\nThe hole density is \tnh=%1.2E /m^3 , nh);
23 // since sigma=ne*e*ue+nh*e*ue and since ne>>nh
24 // therefore sigma=ne*e*ue
25 sigma=ne*e*ue; // calculation of conductivity
26 printf ('\nThe conductivity is \t%.f /ohm-m' ,sigma);
27 rho=1/sigma; // calculation of resistivity
28 R=rho*l/A; // calculation of resistance
29 printf ('\nThe resistance is \tR=% .f ohm' ,R);

```

---

**Scilab code Exa 9.16** Find the mobility and density

```

1 // chapter 9
2 // example 9.16

```

```

3 // Find the mobility and density
4 // page 279
5 clear;
6 clc;
7 //given
8 RH=3.66E-4; // in m^3/C (Hall coefficient)
9 rho=8.93E-3; // in ohm-m (resistivity)
10 e=1.6E-19; // in C (charge of electron)
11 // calculate
12 u=RH/rho; // calculation of mobility
13 n=1/(RH*e); // calculation of density
14 printf ('\nThe mobility is \tu=%f m^2/(V-s) ',u);
15 printf ('\nThe density is \tn=%1.1E /m^3 ',n);

```

---

**Scilab code Exa 9.17** Find the mobility and density of charge carrier

```

1 // chapter 9
2 // example 9.17
3 // Find the mobility and density of charge carrier
4 // page 279–280
5 clear;
6 clc;
7 //given
8 RH=3.66E-4; // in m^3/C (Hall coefficient)
9 rho=8.93E-3; // in ohm-m (resistivity)
10 e=1.6E-19; // in C (charge of electron)
11 // calculate
12 nh=1/(RH*e); // calculation of density of charge
    carrier
13 uh=1/(rho*nh*e); // calculation of mobility of
    charge carrier
14 printf ('\nThe density of charge carrier is \tnh=%1.4
    E /m^3 ',nh);
15 printf ('\nThe mobility of charge carrier is \tuh=%f
    f m^2/(V-s) ',uh);

```



# Chapter 10

## Superconductivity

**Scilab code Exa 10.1** Calculate magnitude of critical magnetic field

```
1 //chapter 10
2 //example 10.1
3 //Calculate magnitude of critical magnetic field
4 //page 313
5 clear;
6 clc;
7 //given
8 Tc=7.2; // in K (critical temperature)
9 T=5; // in K (given temperature)
10 H0=6.5E3; // in A/m (critical magnetic field at 0K)
11 //calculate
12 Hc=H0*(1-(T/Tc)^2); // calculation of magnitude of
   critical magnetic field
13 printf('\nThe magnitude of critical magnetic field
   is \tHc=%1.3E A/m',Hc);
```

---

**Scilab code Exa 10.2** Calculate critical current value

```

1 // chapter 10
2 //example 10.2
3 //Calculate critical current value
4 //page 313
5 clear;
6 clc;
7 //given
8 r=0.02; // in m (radius of ring)
9 Hc=2E3; // in A/m (critical magnetic field at 5K)
10 pi=3.14; // value of pi used in the solution
11 //calculate
12 Ic=2*pi*r*Hc; // calculation of critical current
   value
13 printf ('\nThe critical current value is \tIc=%f A',
   ,Ic);

```

---

### Scilab code Exa 10.3 calculate isotropic mass

```

1 // chapter 10
2 // example 10.3
3 // calculate isotropic mass at 5.1K
4 // page 313
5 clear;
6 clc;
7 // given
8 M1=199.5; // in amu (isotropic mass at 5K)
9 T1=5; // in K (first critical temperature)
10 T2=5.1; // in K (second critical temperature)
11 //calculate
12 // since  $T_c = C * (1 / \sqrt{M})$ 
13 // therefore  $T_1 * \sqrt{M_1} = T_2 * \sqrt{M_2}$ 
14 // therefore we have  $M_2 = (T_1 / T_2)^2 * M_1$ 
15 M2=(T1/T2)^2*M1; // calculation of isotropic mass at
   5.1K
16 printf ('\nThe isotropic mass at 5.1K is \t M2=%f a

```

.m. u. ',M2);

---

### Scilab code Exa 10.4 calculate transition temperature

```
1 // chapter 10
2 // example 10.4
3 // calculate transition temperature
4 // page 314
5 // given
6 clear;
7 clc;
8 T=6; // in K (given temperature)
9 Hc=5E3; // in A/m (critical magnetic field at 5K)
10 H0=2E4; // in A/m (critical magnetic field at 0K)
11 //calculate
12 // since  $H_c = H_0 \cdot (1 - (T/T_c)^2)$ 
13 // therefor we have  $T_c = T / \sqrt{1 - (H_c/H)^2}$ 
14 Tc=T/sqrt(1-(Hc/H0)); // calculation of transition
   temperature
15 printf ('\nThe transition temperature is \tTc=%.3f K',
          Tc);
16 // Note: answer in the book is wrong due to
   calculation mistake
```

---

### Scilab code Exa 10.5 calculate critical current at 5K

```
1 // chapter 10
2 // example 10.5
3 // calculate critical current at 5K
4 // page 314
5 // given
6 clear;
7 clc;
```

```

8 T=5; // in K (given temperature)
9 d=3; // in mm (diameter of the wire)
10 Tc=8; // in K (critical temperature for Pb)
11 H0=5E4; // in A/m (critical magnetic field at 0K)
12 pi=3.14; // value of pi used in the solution
13 //calculate
14 Hc=H0*(1-(T/Tc)^2); // calculation of critical
    magnetic field at 5K
15 printf ('\nThe critical magnetic field at 5K is \tHc=%1.3E A/m',Hc);
16 r=(d*1E-3)/2; // calculation of radius in m
17 Ic=2*pi*r*Hc; // calculation of critical current at
    5K
18 printf ('\nThe critical current at 5K is \tIc=%.4f A',
    ,Ic);
19 //Note: there is slight variation in the answer due
    to round off

```

---

### Scilab code Exa 10.6 calculate frequency of EM waves

```

1 // chapter 10
2 // example 10.6
3 // calculate frequency of EM waves
4 // page 314
5 clear;
6 clc;
7 // given
8 V=8.50; // in micro V (voltage across Josephson
    junction )
9 e=1.6E-19; // in C (charge of electron)
10 h=6.626E-34; // in J/s (Planck s constant)
11 //calculate
12 V=V*1E-6; // changing unit from V to microVolt
13 v=2*e*V/h; // calculation of frequency of EM waves
14 printf ('\nThe frequency of EM waves is \tv=%1.3E Hz'

```

```
,v);  
15 // Note: the answer in the book is wrong due to  
// calculation misatke
```

---

### Scilab code Exa 10.7 calculate transition temperature of the isotopes

```
1 // chapter 10  
2 // example 10.7  
3 // calculate transition temperature of the isotopes  
4 // page 315  
5 clear;  
6 clc; // given  
7 M1=200.59; // in amu (average atomic mass at 4.153K  
//)  
8 Tc1=4.153; // in K (first critical temperature)  
9 M2=204; // in amu (average atomic mass of isotopes)  
10 //calculate  
11 // since  $T_c = C * (1 / \sqrt{M})$   
12 // therefore  $T_1 * \sqrt{M_1} = T_2 * \sqrt{M_2}$   
13 // therefore we have  $T_c = T_1 * \sqrt{M_1 / M_2}$   
14 Tc2=Tc1*sqrt(M1/M2); //calculation of transition  
// temperature of the isotopes  
15 printf('\nThe transition temperature of the isotopes  
is \t Tc2=%f K',Tc2);
```

---

# Chapter 12

## Fibre Optics

**Scilab code Exa 12.1** calculate fractional index change for a given optical fibre

```
1 // chapter 12
2 // example 12.1
3 // calculate fractional index change for a given
   optical fibre
4 // page 360
5 clear;
6 clc;
7 // given
8 u1=1.563; // refractive index of core
9 u2=1.498; // refractive index of cladding
10 //calculate
11 d=(u1-u2)/u1; // calculation of fractional index
   change
12 printf ('\nThe fractional index change for a given
   optical fibre is %.4f ',d);
```

---

**Scilab code Exa 12.2** calculate numerical aperture and the acceptance angle of an optical fibre

```

1 // chapter 12
2 // example 12.2
3 // calculate numerical aperture and the acceptance
   angle of an optical fibre
4 // page 360
5 clear;
6 clc;
7 // given
8 u1=1.55; // refractive index of core
9 u2=1.50; // refractive index of cladding
10 //calculate
11 d=(u1-u2)/u1; // calculation of fractional index
   change
12 NA=u1*sqrt(2*d); // calculation of numerical
   aperture
13 printf('\nThe numerical aperture of the fibre is \
   tNA=%f',NA);
14 theta=asind(NA); // calculation of acceptance angle
15 printf('\nThe acceptance angle of the optical fibre
   is \t%.1f degree',theta);

```

---

**Scilab code Exa 12.3** calculate the acceptance angle of an optical fibre

```

1 // chapter 12
2 // example 12.3
3 // calculate the acceptance angle of an optical
   fibre
4 // page 360
5 // given
6 clear;
7 clc;
8 u1=1.563; // refractive index of core
9 u2=1.498; // refractive index of cladding
10 //calculate
11 NA=sqrt(u1^2-u2^2); // calculation of numerical

```

```

        aperture
12 printf('\nThe numerical aperture of the fibre is \
          tNA=%f ,NA);
13 theta=asind(NA); // calculation of acceptance angle
14 printf('\nThe acceptance angle of the optical fibre
          is \t%.2f degree ',theta);

```

---

**Scilab code Exa 12.4** calculate refractive index of material of the core

```

1 // chapter 12
2 // example 12.4
3 // calculate refractive index of material of the
   core
4 // page 360–361
5 clear;
6 clc;
7 // given
8 NA=0.39; //numerical aperture of the optical fibre
9 d=0.05; // difference in the refractive index of the
           material of the core and cladding
10 //calculate
11 // since NA=u1*sqrt(2*d)
12 //we have u1=NA/sqrt(2*d)
13 u1= NA/sqrt(2*d); // calculation of refractive index
           of material of the core
14 printf('\nThe refractive index of material of the
           core is \tu1=%f ',u1);

```

---

**Scilab code Exa 12.5** calculate numerical aperture acceptance angle and the critical angle of the optical fibre

```

1 // chapter 12
2 // example 12.5

```

```

3 // calculate numerical aperture , acceptance angle and
   the critical angle of the optical fibre
4 // page 361
5 clear;
6 clc;
7 // given
8 u1=1.50; // refractive index of core
9 u2=1.45; // refractive index of cladding
10 //calculate
11 d=(u1-u2)/u1; // calculation of fractional index
   change
12 NA=u1*sqrt(2*d); // calculation of numerical
   aperture
13 printf ('\nThe numerical aperture of the fibre is \
   tNA=%.3f',NA);
14 theta_0=asind(NA); // calculation of acceptance
   angle
15 printf ('\nThe acceptance angle of the optical fibre
   is \t%.2f degree',theta_0);
16 theta_c=asind(u2/u1); // calculation of critical
   angle
17 printf ('\nThe critical angle of the optical fibre is
   \t%.1f degree',theta_c);

```

---

**Scilab code Exa 12.6** calculate refractive index of the core and cladding material of a fibre

```

1 // chapter 12
2 // example 12.6
3 // calculate refractive index of the core and
   cladding material of a fibre
4 // page 361
5 clear;
6 clc;
7 // given

```

```

8 NA=0.33; // numerical aperture
9 d=0.02; // difference in the refractive index of the
    core and cladding of the material
10 // calculate
11 // since  $NA = u_1 \sqrt{2d}$ 
12 // therefore we have
13 u1=NA/sqrt(2*d); // calculation of refractive index
    of the core
14 // since  $d = (u_1 - u_2)/u_2$ 
15 // therefore we have
16 u2=(1-d)*u1; // calculation of refractive index of
    the cladding
17 printf('\nThe refractive index of the core is \tu1=%
    .1f',u1);
18 printf('\nThe refractive index of the cladding is \%
    tu2=%.3f',u2);
19 // Note: In the question, it is given that NA=0.33
    but in the book NA=0.22 has been used in the
    solution. That's why answer in the book is
    different from that of generated from the code

```

---

**Scilab code Exa 12.7** calculate numerical aperture and acceptance angle of the symmetrical fibre

```

1 // chapter 12
2 // example 12.7
3 // calculate numerical aperture and acceptance angle
    of the symmetrical fibre
4 // page 361
5 clear;
6 clc;
7 // given
8 u1=3.5; // refractive index of core
9 u2=3.45; // refractive index of cladding
10 u0=1; // refractive index of the air

```

```
11 // calculate
12 NA=sqrt(u1^2-u2^2); // calculation of numerical
    aperture
13 NA=NA/u0;
14 printf('\nThe numerical aperture of the fibre is \
    tNA=%f',NA);
15 alpha=asind(NA); // calculation of acceptance angle
16 printf('\nThe acceptance angle of the optical fibre
    is \t%.2f degree',alpha);
```

---

**Scilab code Exa 12.8** calculate numerical aperture and acceptance angle of the symmetrical fibre

```
1 // chapter 12
2 // example 12.8
3 // calculate numerical aperture and acceptance angle
    of an optical fibre
4 // page 361–362
5 clear;
6 clc;
7 // given
8 u1=1.48; // refractive index of core
9 u2=1.45; // refractive index of cladding
10 //calculate
11 NA=sqrt(u1^2-u2^2); // calculation of numerical
    aperture
12 printf('\nThe numerical aperture of the fibre is \
    tNA=%f',NA);
13 theta=asind(NA); // calculation of acceptance angle
14 printf('\nThe acceptance angle of the optical fibre
    is \t%.2f degree',theta);
15 // Note: there is slight variation in the answer due
    to round off
```

---