

Scilab Textbook Companion for
Modern Physics
by K. S. Krane¹

Created by
Balakavi Venkata Siva Sai Krishna
B.TECH
Electrical Engineering
VNIT,NAGPUR.
College Teacher
V S Kale
Cross-Checked by

August 10, 2013

¹Funded by a grant from the National Mission on Education through ICT, <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab codes written in it can be downloaded from the "Textbook Companion Project" section at the website <http://scilab.in>

Book Description

Title: Modern Physics

Author: K. S. Krane

Publisher: John Wiley & Sons

Edition: 2

Year: 1996

ISBN: 978-81-265-0826-6

Scilab numbering policy used in this document and the relation to the above book.

Exa Example (Solved example)

Eqn Equation (Particular equation of the above book)

AP Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

Contents

| | |
|--|----|
| List of Scilab Codes | 5 |
| 1 Introduction | 9 |
| 2 The Special Theory of Relativity | 11 |
| 3 Review of Electromagnetic waves | 21 |
| 4 The Wavelike properties of particles | 25 |
| 5 The Schrodinger Equation | 33 |
| 6 The Rutherford Bohr model of an atom | 36 |
| 7 The Hydrogen atom in wave mechanics | 41 |
| 8 Many Electron Atoms | 45 |
| 9 Molecular Structure | 49 |
| 10 Statistical Physics | 53 |
| 11 Properties of Ionic Crystals | 57 |
| 12 Nuclear Structure and Radioactivity | 60 |
| 13 Nuclear Reaction and Applications | 69 |
| 14 The Four Basic Forces | 73 |

| | |
|--|----|
| 15 Astrophysics and General Relativity | 78 |
| 16 The Cosmic Microwave Background Radiation | 80 |

List of Scilab Codes

| | | |
|----------|---|----|
| Exa 1.1 | Mass difference between a proton and a neutron | 9 |
| Exa 1.2 | Total mass of proton and electron | 9 |
| Exa 1.3 | Value of hc | 10 |
| Exa 2.1 | Speed of A wrt B | 11 |
| Exa 2.2 | Velocity of plane wrt ground | 11 |
| Exa 2.3 | Time taken in each case | 12 |
| Exa 2.4 | Minimum speed for survival of muons | 12 |
| Exa 2.5 | Apparent thickness of Earth atmosphere | 13 |
| Exa 2.6 | Solution for a b c d e | 13 |
| Exa 2.7 | Speed of missile wrt earth | 14 |
| Exa 2.8 | Speed of galaxy wrt earth | 15 |
| Exa 2.9 | Velocity of rocket2 wrt rocket1 | 15 |
| Exa 2.10 | Time interval between the events | 16 |
| Exa 2.11 | momentum of proton | 16 |
| Exa 2.12 | Various energies of proton | 17 |
| Exa 2.13 | Velocity and momentum of electron | 17 |
| Exa 2.14 | Solution for a b | 18 |
| Exa 2.15 | Rate of decrease of the mass of Sun | 18 |
| Exa 2.16 | Kinetic energy of pion in each case | 19 |
| Exa 2.17 | Threshold kinetic energy to produce antiprotons | 20 |
| Exa 3.1 | Atomic Spacing of Nacl | 21 |
| Exa 3.2 | Time taken to release an electron | 21 |
| Exa 3.3 | Solution for a and b | 22 |
| Exa 3.4 | Solution for a b and c | 22 |
| Exa 3.5 | Solution for a b and c | 23 |
| Exa 3.6 | Solution for a b c and d | 24 |
| Exa 4.1 | Solution for a b c d and e | 25 |
| Exa 4.2 | Minimum uncertainty in wavelength | 26 |

| | | |
|----------|---|----|
| Exa 4.3 | Validity of the claim | 27 |
| Exa 4.4 | Solution for a and b | 27 |
| Exa 4.5 | Solution for a and b | 28 |
| Exa 4.6 | Uncertainty in x component | 29 |
| Exa 4.7 | Range of kinetic energy of an electron | 29 |
| Exa 4.8 | Solution for a b and c | 30 |
| Exa 4.9 | minimum velocity of the billiard ball | 31 |
| Exa 4.10 | Group velocity of a wave packet in terms of phase velocity | 31 |
| Exa 5.1 | Displacement and velocity of the object | 33 |
| Exa 5.2 | Solution for a and b | 33 |
| Exa 5.3 | Proof for average value of x | 34 |
| Exa 6.1 | Average deflection angle per collision | 36 |
| Exa 6.2 | Solution for a and b | 36 |
| Exa 6.3 | Distance of closest approach | 37 |
| Exa 6.4 | Three longest wavelengths of the Paschen series | 38 |
| Exa 6.5 | Various wavelegths in Balmer and Lymann series | 38 |
| Exa 6.6 | wavelengths of transition | 39 |
| Exa 6.7 | Two longest wavelengths of triply ionized beryllium | 39 |
| Exa 7.1 | Proof | 41 |
| Exa 7.2 | Probability of finding an electron closer to nucles than Bohrs orbit | 41 |
| Exa 7.3 | Probability of finding an electron inisde Bohr Radius | 42 |
| Exa 7.4 | Length of angular momentum vectors | 42 |
| Exa 7.5 | possible Z components of the vector L | 43 |
| Exa 7.6 | Seperation of beams as they leave the magnet | 43 |
| Exa 7.7 | Change in wavelength | 44 |
| Exa 8.1 | Energy of Ka X ray of sodium | 45 |
| Exa 8.2 | Solution for a and b | 45 |
| Exa 8.3 | Total orbital and spin quantum numbers of carbon | 46 |
| Exa 8.4 | Total orbital and spin quantum numbers of nitrogen | 46 |
| Exa 8.5 | Hunds rule to find ground state quantum numbers of nitrogen | 47 |
| Exa 8.6 | Ground state L and S of oxygen | 47 |
| Exa 9.1 | Charge on the sphere | 49 |
| Exa 9.2 | Solution for a and b | 49 |
| Exa 9.3 | vibrational frequency and photon energy of H2 | 50 |
| Exa 9.4 | Energies and wavelengths of 3 lowest radiations emitted by molecular H2 | 51 |

| | | |
|-----------|---|----|
| Exa 9.5 | Rotational Inertia of molecule | 51 |
| Exa 9.6 | Solution for a and b | 52 |
| Exa 10.1 | Various Speeds obtained from maxwell speed distribution | 53 |
| Exa 10.2 | Frequency distribution of emitted light | 53 |
| Exa 10.3 | Solution for a and b | 54 |
| Exa 10.4 | Solution for a and b | 55 |
| Exa 10.5 | Fermi Energy E_f for sodium | 55 |
| Exa 11.1 | Solution for a and b | 57 |
| Exa 11.2 | Energy per neytral atom to take apart a crystal of Nacl | 58 |
| Exa 11.3 | Solution for a and b | 58 |
| Exa 12.1 | Solution for a and b | 60 |
| Exa 12.2 | Approximate nuclear radii | 61 |
| Exa 12.3 | Density of typical nucleus and resultant mass | 61 |
| Exa 12.4 | Total Binding Energy | 62 |
| Exa 12.5 | Solution for a b c and d | 62 |
| Exa 12.6 | Atoms at the time of solidification | 63 |
| Exa 12.7 | Kinetic energy of alpha particle emitted in alpha decay | 64 |
| Exa 12.8 | Q value of ^{14}C emission | 64 |
| Exa 12.9 | Maximum Kinetic energy of emitted electron | 65 |
| Exa 12.10 | Q value of various decays | 65 |
| Exa 12.11 | Maximum kinetic energy of emitted beta particle | 66 |
| Exa 12.12 | Rate of energy production per gram of uranium | 67 |
| Exa 12.13 | Ages of the given rocks | 67 |
| Exa 12.14 | Solution for a and b | 68 |
| Exa 13.1 | Rate of production of neutron | 69 |
| Exa 13.2 | Resultant activity of ^{198}Au | 70 |
| Exa 13.3 | Solution for a and b | 70 |
| Exa 13.4 | Solution for a and b | 71 |
| Exa 13.5 | Solution for a and b | 71 |
| Exa 14.1 | Solution for a and b | 73 |
| Exa 14.2 | Energy of the proton and pi meson | 73 |
| Exa 14.3 | Maximum kinetic energy of the electron emitted in the decay | 74 |
| Exa 14.4 | maximum energy of the positron nad pi mesons | 74 |
| Exa 14.5 | Q values for reaction | 75 |
| Exa 14.6 | Threshold Kinetic energy to produce pi mesons | 75 |
| Exa 14.7 | Threshold Energy of the given reaction | 76 |
| Exa 14.8 | Solution for a and b | 76 |

| | | |
|----------|---|----|
| Exa 15.1 | Change in wavelength in solar spectrum due to gravitational shift | 78 |
| Exa 15.2 | Maximum energy of neutrino in the first reaction of proton proton cycle | 79 |
| Exa 16.1 | Resultin temperature of interstellar space | 80 |
| Exa 16.2 | Solution for a and b | 81 |
| Exa 16.3 | Relative number of neutrons and protons among the nucleus | 81 |

Chapter 1

Introduction

Scilab code Exa 1.1 Mass difference between a proton and a neutron

```
1 clear
2 clc
3 disp('Exa-1.1');
4 Mn=1.008665;Mp=1.007276
                                     //Given mass
   of an electron and a proton in terms of u
5 Md= Mn-Mp;

   //mass difference
6 printf('Mass difference in terms of U is %f ',Md);
7 Md=Md*931.50;
                                     //
   converting u into Mev/c^2 by multiplying by 931.5
   MeV/c^2
8 printf('which equals %.3f Mev/c^2. ',Md);
```

Scilab code Exa 1.2 Total mass of proton and electron

```

1 clear
2 clc
3 disp('Exa-1.2');
4 Mp=1.007276 ; Me=5.4858*10^-4; //mass of proton and
   electron in terms of U
5 Mt=Mp+Me; //Total mass= sum of
   above masses
6 printf('The combined mass of an electron and a
   proton was found out to be %f U.',Mt);

```

Scilab code Exa 1.3 Value of hc

```

1 clear
2 clc
3 disp('Exa-1.3');
4 h=6.621*10^-34 ; c=2.9979*10^8; //
   h is in J/s and c is in m/s
5 hc=h*c*((10^9)/(1.6022*10^-19)); //1e
   =1.602*10^-19 J and 1 m=10^9 nm
6 printf('The value of hc is %f eV.nm\n',hc);
7 printf('Rounding off to 4 digits, we obtain %4.f eV.
   nm.',hc);
8 disp('Hence zero at the end is significant.');
```

Chapter 2

The Special Theory of Relativity

Scilab code Exa 2.1 Speed of A wrt B

```
1 clear
2 clc
3 disp('Exa-2.1');
4 v1=60; v2=40 //Velocities of cars wrt to
  observer in km/hr
5 vr=v1-v2; //relative velocity
6 printf('The value of relative velocity is %4.f km/h.
  ',vr);
```

Scilab code Exa 2.2 Velocity of plane wrt ground

```
1 clear
2 clc
3 disp('Exa-2.2');
```

```

4 Va_w=[320 0]; Vw_g=[0 65]; //Vp/q=[X Y]=>
    velocity of object p wrt q along X(east) and Y(
    north) directions.
5 Va_g=Va_w + Vw_g; //net velocity
6 k=norm(Va_g); //magnitude
7 s=atan(Va_g(1,2)/Va_g(1,1))*180/%pi; //angle in
    rad*180/pi for conversion to degrees
8 printf('The magnitude of velocity Va/g(airplane wrt
    ground) is %.3f Km/h at %.3f degrees north of
    east.',k,s);

```

Scilab code Exa 2.3 Time taken in each case

```

1 clear
2 clc
3 disp('Exa-2.3'); // The problem is entirely
    theoretical hence following the standard
    procedure we obtain
4 printf('The time required for round trip is 2*L/(c
    *(1-(u/c)^2)). \n');
5 printf('The time required to swim across and return
    is 2*L/(c*sqrt((1-(u/c)^2)))');

```

Scilab code Exa 2.4 Minimum speed for survival of muons

```

1 clear
2 clc
3 disp('Exa-2.3');

```

```

4 Lo=100*(10^3);c=3*(10^8); //Given values//all the
  quantities are converted to SI units
5 d=2.2*(10^-6); //time between its birth and
  decay
6 t=Lo/c //where Lo is the distance from
  top of atmosphere to the Earth. c is the velocity
  of light. t is the time taken
7 u=sqrt(1-((d/t)^2)); // using time dilation formula
  for finding u where u is the minimum velocity in
  terms of c;
8 printf('Hence the minimum speed required is %f c.',u
  );

```

Scilab code Exa 2.5 Apparent thickness of Earth atmosphere

```

1 clear
2 clc
3 disp('Exa-2.5');
4 Lo=100*(10^3); //Lo is converted to Km
5 u=0.999978; //u/c is taken as u since u is
  represented in terms of c.
6 L=Lo*(sqrt(1-u^2)); // from the length contraction
  formula
7 printf('Hence the apparent thickness of the Earth''s
  surface is %.2f metres.',L);

```

Scilab code Exa 2.6 Solution for a b c d e

```

1 clear all

```

```

2  clc
3  disp('Exa-2.6(a)');
4  L=65; c=3*10^8; u=0.8*c;
5  t=L/u ;           //The value of time taken as
                    measured by the observer
6  printf('The time for rocket to pass a point as
          measured by O is %.2e.\n',t);           //The value
          of time taken as measured by the observer
7  disp('Exa-2.6(b)');
8  Do=65;           //given length
9  Lo= L/sqrt(1-(u/c)^2);           //contracted
          length of rocket
10 printf('Actual length according to O is %.2f.\n',Lo)
    ;
11 disp('Exa-2.6(c)');
12 D=Do*(sqrt(1-(u/c)^2));           //contracted length of
          platform.
13 printf('Contracted length according to O'' is %.2e.\n
          n',D);
14 disp('Exa-2.6(d)');
15 t1=Lo/u;           //time needed to pass
          according to O'.
16 printf('Time taken according to O is %.2e.\n',t1);
17 disp('Exa-2.6(e)');
18 t2=(Lo-D)/u;           //time intervals
          between the two instanscs
19 printf('Time taken according to O'' is %.2e.\n',t2);
20 disp('The value of t1 and t2 did not match');

```

Scilab code Exa 2.7 Speed of missile wrt earth

```

1  clear all
2  clc

```

```

3 disp('Exa-2.7');
4 v1=0.6; u=0.8; c=1; // all the values are measured
    in terms of c hence c=1
5 v= (v1+u)/(1+(v1*u/c^2));
6 printf('The speed of missile as measured by an
    observer on earth is %.2f c.',v);

```

Scilab code Exa 2.8 Speed of galaxy wrt earth

```

1 clear all
2 clc
3 disp('Exa-2.8');
4 w1=600;w2=434; // w1=recorded wavelength;w2=actual
    wavelength
5 // c/w1 = c/w2 *(sqrt(1-u/c)/(1+u/c))
6 k=w2/w1;
7 x=(1-k^2)/(1+k^2); //solving for u/c
8 printf('The speed of galaxy wrt earth is %.2f c',x);

```

Scilab code Exa 2.9 Velocity of rocket2 wrt rocket1

```

1 clear all
2 clc
3 disp('Exa-2.9');
4 v1x=0.6; v1y=0; v2x=0; v2y=.8; c=1; // all the
    velocities are taken wrt c
5 v21x=(v2x-v1x)/(1-(v1x*v2x/c^2)); //using lorentz
    velocity transformation
6 v21y=(v2y*(sqrt(1-(v1x*c)^2)/c^2))/(1-v1y*v2y/c^2)

```



```

7 printf('The velocity of rocket 2 wrt rocket 1 along
  x and y directions is %.2f c & %.2f c
  respectively ',v21x,v21y);

```

Scilab code Exa 2.10 Time interval between the events

```

1 clear all
2 clc
3 disp('Exa-2.10');
4 u=0.8*c;L=65;c=3*10^8; //all values are
  in terms of c
5 t=u*L/(c^2*(sqrt(1-((u/c)^2)))); //from the
  equation 2.31
6 printf('The time interval between the events is %e
  sec which equals %.2f usec.',t,t*10^6);

```

Scilab code Exa 2.11 momentum of proton

```

1 clear all
2 clc
3 disp('Exa-2.11');
4 m=1.67*10^-27;c= 3*10^8;v=0.86*c; //all the
  given values and constants
5 p=m*v/(sqrt(1-((v/c)^2))); // in terms of
  Kgm/sec
6 printf('The value of momentum was found out to be %
  .3e Kg-m/sec.\n',p);
7 c=938;v=0.86*c;mc2=938 // all the
  energies in MeV where mc2= value of m*c^2

```

```

8 pc=(mc2*(v/c))/(sqrt(1-((v/c)^2))); //expressing
   in terms of Mev
9 printf('The value of momentum was found out to be %
   .2f Mev.',pc);

```

Scilab code Exa 2.12 Various energies of proton

```

1 clear all
2 clc
3 disp('Exa-2.12');
4 pc=1580; mc2=938;E0=938; // all the energies in
   MeV mc2=m*c^2 and pc=p*c
5 E=sqrt(pc^2+mc2^2);
6 printf('The relativistic total energy is %.2f MeV.\n
   ',E); //value of Energy E
7 K=E-E0; //value of possible
   kinetic energy
8 printf('The kinetic energy of the proton is %.1f MeV
   .',K);

```

Scilab code Exa 2.13 Velocity and momentum of electron

```

1 clear all
2 clc
3 disp('Exa-2.13');
4 E=10.51; mc2=0.511; //all the values are in MeV
5 p=sqrt(E^2-mc2^2); //momentum of the electron
6 printf('The momentum of electron is %.1f MeV/c\n',p)
   ;

```

```
7 v=sqrt(1-(mc2/E)^2); //velocity in terms of c
8 printf('The velocity of electron is %.4f c',v);
```

Scilab code Exa 2.14 Solution for a b

```
1 clear all
2 clc
3 disp('Exa-2.14');
4 k=50;mc2=0.511*10^-3;c=3*10^8; // all the values of
   energy are in GeV and c is in SI units
5 v=sqrt(1-(1/(1+(k/mc2))^2)); //speed of the
   electron in terms of c
6 k=c-(v*c); //difference in
   velocities
7 printf('Speed of the electron as a fraction of c is
   %.12f*10^-12.\n',v*10^12); // v=(v*10^12)
   *10^-12; so as to obtain desired accuracy in the
   result
8 printf('The difference in velocities is %.1f cm/s.',
   k*10^2);
```

Scilab code Exa 2.15 Rate of decrease of the mass of Sun

```
1 clear all
2 clc
3 disp('Exa-2.15');
4 r=1.5*10^11; I=1.4*10^3; //radius and intensity
   of sun
```

```

5 s=4*%pi*r^2 //surface area of the
   sun
6 Pr=s*I // Power radiated in J/
   sec
7 c=3*10^8; //velocity of light
8 m=Pr/c^2 //rate od decrease of mass
9 printf('The rate of decrease in mass of the sun is %
   .1e kg/sec.',m);

```

Scilab code Exa 2.16 Kinetic energy of pion in each case

```

1 clear all
2 clc
3 disp('Exa-2.16');
4 K=325; mkc2=498; //kinetic energy and rest mass
   energy of kaons
5 mpic=140; //given value
6 Ek=K+mkc2;
7 pkc=sqrt(Ek^2-mkc2^2);
8 //consider the law of conservation of energy which
   yields Ek=sqrt(plc^2+mpic^2)+sqrt(p2c^2+mpic^2)
9 // The above equations (4th degree,hence no direct
   methods)can be solved by assuming the value of
   p2c=0.
10 plc=sqrt(Ek^2-(2*mpic*Ek));
11 //consider the law of conservation of momentum.
   which gives plc+p2c=pkc implies
12 p2c=pkc-plc;
13 k1=(sqrt(plc^2+(mpic^2))-mpic); //corresponding
   kinetic energies
14 k2=(sqrt((p2c^2)+(mpic^2))-mpic);
15 printf('The corresponding kinetic energies of the
   pions are %.0f MeV and %.1f MeV.',k1,k2);

```

Scilab code Exa 2.17 Threshold kinetic energy to produce antiprotons

```
1 clear all
2 clc
3 disp('Ex-2.17');
4 mpc2=938;c=3*10^8; //mpc2=mp*c^2,mp=mass of proton
5 Et=4*mpc2; //final total energy
6 E1=Et/2;E2=E1; //applying conservation of
   momentum and energy
7 v2=c*sqrt(1-(mpc2/E1)^2); //lorentz
   transformation
8 u=v2;v=(v2+u)/(1+(u*v2/c^2));
9 E=mpc2/(sqrt(1-(v/c)^2));
10 K=E-mpc2;
11 printf('The threshold kinetic energy is %.3f Gev',K
   /10^3);
```

Chapter 3

Review of Electromagnetic waves

Scilab code Exa 3.1 Atomic Spacing of Nacl

```
1 clear
2 clc
3 disp('Exa-3.1');
4 w=0.250; theta=26.3;n=1 // n=1 for hydrogen atom and
    rest all are given values
5 d=n*w/(2*sind(theta)); // bragg's law
6 printf('Hence the atomic spacing is %.3f nm.',d);
```

Scilab code Exa 3.2 Time taken to release an electron

```
1 clear
2 clc
3 disp('Exa-3.2');
4 I=120;r=0.1*10^-9;Eev=2.3 //I-intensity in W/m^2 r
    in m & E in electron volt
```

```

5 A=%pi*r^2;K=1.6*10^-19; // A=area and K is
  conversion factor from ev to joules
6 t= Eev*K/(I*A); //time interval
7 printf('The value of time interval was found out to
  be %.1f sec',t);

```

Scilab code Exa 3.3 Solution for a and b

```

1 clear
2 clc
3 disp('Exa-3.3(a)');
4 w=650*10^-9;h=6.63*10^-34;c=3*10^8; //given values
  and constant taken in comfortable units
5 E=h*c/w; printf('The Energy of the electron is %.3e
  J ',E);
6 E=E/(1.6*10^-19);printf('which is equivalent to %f
  eV\n',E);
7 printf('The momentum of electron is p=E/c i.e %.2f/c
  \n',E);
8 disp('Exa-3.3(b)');
9 E2=2.40; //given energy of
  photon.
10 w2=h*c*10^9/(E2*1.6*10^-19); //converting the
  energy in to eV and nm
11 printf('The wavelength of the photon is %.2f nm',w2)
  ;

```

Scilab code Exa 3.4 Solution for a b and c

```

1 clear
2 clc
3 disp('Exa-3.4(a)');
4 hc=1240; phi=4.52 //both the values
   are in eV
5 w1=hc/phi;
6 printf('The cutoff wavelength of the tungsten metal
   is %.3f nm\n ',w1);
7 disp('Exa-3.4(b)');
8 w2=198; //given value of wavelength
9 Kmax=(hc/w2)-phi; printf('The max value of kinetic
   energy is %.3f eV\n ',Kmax);
10 disp('Exa-3.4(c)');
11 Vs=Kmax; printf('The numerical value of the max
   kinetic energy is same as stopping potential in
   volts. Hence %.2f V',Vs);

```

Scilab code Exa 3.5 Solution for a b and c

```

1 clear
2 clc
3 disp('Exa-3.5(a)');
4 T1=293; Kw=2.898*10^-3;
5 w1=Kw/T1;
6 printf('The wavelength at which emits maximum
   radiation is %.2f um.\n ',w1*10^6);
7 disp('Exa-3.5(b)');
8 w2=650*10^-9;
9 T2=Kw/w2;
10 printf('The temperature of the object must be raised
   to %.0f K.\n ',T2);
11 disp('Exa-3.5(c)');
12 x=(T2/T1)^4; printf('Thus the thermal radiation at

```


higher temperature is $1.2e$ times the room (lower) temperature.\n',x);

Scilab code Exa 3.6 Solution for a b c and d

```
1 clear
2 clc
3 disp('Exa-3.6(a)');
4 w1=0.24;wc=0.00243;theta=60; //given values w=
   wavelength(lambda)
5 w2=w1+(wc*(1-cosd(theta)));
6 printf('The wavelength of x-rays after scattering is
   %.4f nm\n',w2);
7 disp('Exa-3.6(b)');
8 hc=1240;
9 E2=hc/w2;E1=hc/w1; printf('The energy of scattered x
   -rays is %.0f eV\n',E2);
10 disp('Exa-3.6(c)');
11 K= E1-E2; //The kinetic energy is the difference in
   the energy before and after the collision;
12 printf('The kinetic energy of the x-rays is %.3f eV\
   n',K);
13 disp('Exa-3.6(d)');
14 phi2=atand(E2*sind(theta)/(E1-E2*cosd(theta)))
15 printf('The direction of the scattered electron is %
   .1f degrees',phi2);
```

Chapter 4

The Wavelike properties of particles

Scilab code Exa 4.1 Solution for a b c d and e

```
1 clear
2 clc
3 disp("Ex: 4.1 ");
4 h=6.6*10^-34; // h(
    planck's constant)= 6.6*10^-34
5 m1= 10^3;v1=100;; // for
    automobile
6 w1= h/(m1*v1); // ['w'-
    wavelength in metre 'm'-mass in Kg 'v'-velocity in
    metres/sec.] of the particles
7 printf("Wavelength of the automobile is %1.2e m\n",
    w1 );
8 m2=10*(10^-3);v2= 500; // for
    bullet
9 w2=h/(m2*v2);
10 printf("Wavelength of the bullet is %1.2e m\n ",w2 )
    ;
11 m3=(10^-9)*(10^-3); v3=1*10^-2;
12 w3=h/(m3*v3);
```

```

13 printf("Wavelength of the smoke particle is %1.2e m\n
    n",w3 );
14 m4=9.1*10^-31;k=1*1.6*10^-19; // k-
    kinetic energy of the electron & using 1ev =
    1.6*10^-19 joule
15 p=sqrt(2*m4*k); // p=
    momentum of electron ;from K=1/2*m*v^2
16 w4=h/p;
17 printf("Wavelength of the electron(1ev) is %1.2fm\n
    ",w4*10^9 );
18 hc=1240;pc=100 // In the
    extreme relativistic realm , K=E=pc; Given pc=100
    MeV, hc=1240MeV
19 w5= hc/pc;
20 printf("Wavelength of the electron (100Mev) is %1.2 f
    fm\n",w5);

```

Scilab code Exa 4.2 Minimum uncertainty in wavelength

```

1 clear
2 clc
3 disp('Ex-4.2');
4 // w=wavelength; consider k=2*(pi/w);
5 // differentiate k w.r.t w and replace del(k)/del(w)
    = 1 for equation.4.3
6 // which gives del(w)= w^2 /(2*pi*del(x)), hence
7 w=20; delx=200; // delx=200cm and w=20cm
8 delw=(w^2)/(delx*2*pi);
9 printf('Hence uncertainty in length is %1.2 f cm',
    delw);

```

Scilab code Exa 4.3 Validity of the claim

```
1 clear
2 clc
3 disp('Ex-4.3')
4 delw=1; //consider time interval of 1
   sec
5 delw=1/delt; // since delw*delt =1 from
   equation 4.4
6 delf=0.01 //calculated accuracy is 0.01Hz
7 delwc =2*%pi*delf // delwc-claimed accuracy from w
   =2*pi*f
8 printf('The minimum uncertainty calculated is 1rad/
   sec. The claimed accuracy is %.3f rad/sec\n',
   delwc);
9 if delw==delwc then disp('Valid claim');
10 end
11 if delw~=delwc then disp('Invalid claim');
12 end
```

Scilab code Exa 4.4 Solution for a and b

```
1 clear
2 clc
3 disp('Ex-4.4(a)');
4 m=9.11*10^-31; v=3.6*10^6; // 'm', 'v' - mass an
   velocity of the electron in SI units
5 h=1.05*10^-34; //planck's constant in SI
```

```

6 p=m*v; //momentum
7 delp=p*0.01;//due to 1% precision in p
8 delx = h/delp//uncertainty in position
9 printf('Uncertainty in position is %1.2f nm',delx
    *10^9);
10 disp('Ex-4.4((b)')
11 printf('Since the motion is strictly along X-
    direction , its velocity in Y direction is
    absolutely zero.\n So uncertainty in velocity
    along y is zero=> uncertainty in position along
    y is infinite. \nSo nothing can be said about its
    position/motion along Y')

```

Scilab code Exa 4.5 Solution for a and b

```

1 clear
2 clc
3 disp('Ex-4.5(a)');
4 m=0.145;v=42.5; //'m','v' - mass an velocity of the
    electron in SI units
5 h=1.05*10^-34; //planck's constant in SI
6 p=m*v; //momentum
7 delp=p*0.01;//due to 1% precision in p
8 delx = h/delp//uncertainty in position
9 printf('Uncertainty in position is %1.2e',delx);
10 disp('Ex-4.5(b)');
11 printf('Motion along y is unpredictable as long as
    the velocity along y is exactly known(as zero).');

```

Scilab code Exa 4.6 Uncertainty in x component

```
1 clc
2 clear
3 disp('Ex-4.6')
4 printf('The uncertainty in the poosition of
        electron after it passes through the slit is
        reduced to width of the slit\n delx=a\n');
5 printf('The uncertainty in momentum = h/a\n');
6 printf('Position of landing(angle t) = sin t = tan t
        = delz/dely =(h/a)/2*pi*a= w/2*pi*a \nwhere w=
        wavelength\n');
7 printf('Rewriting the above expression a*sint = w
        /(2*pi)\n which is similar to a*sint = w (
        neglect 2*pi)as found out by first minimum in
        diffraction by a slit of width a');
8 disp('It proves a close connection between wave
        behaviour and uncertainty principle');
```

Scilab code Exa 4.7 Range of kinetic energy of an electron

```
1 clear
2 clc
3 disp('Ex-4.7');
4 mc2=2.15*10^-4; //mc2 is the mass of the
                  electron, considered in Mev for the simplicity in
                  calculations
5 hc=197 // The value of h*c in Mev.
         fm for simplicity
6 delx= 10 // Given uncertainty in
           position=diameter of nucleus= 10 fm
7 delp= hc/delx ; //Uncertainiy in momentum per
                 unit 'c' i.e (Mev/c) delp= h/delx =(h*c)/(c*delx)
```

```

        ;hc=197 Mev.fm  1Mev=1.6*10^-13 Joules ')
8  p=delp;           // Equating delp to p as a
    consequence of equation 4.10
9  K1=[[p]^2]+[mc2]^2 // The following 3 steps are
    the steps involved in calculating K.E= sqrt((p*c
    )^2 + (mc^2)^2)- m*c^2
10 K1=sqrt(K1)
11 K1= K1-(mc2);
12 printf('Kinetic energy was found out to be %d Mev',
    K1)

```

Scilab code Exa 4.8 Solution for a b and c

```

1  clear
2  clc
3  disp('Ex-4.8')
4  h=6.58*10^-16; // plack's constant
5  delT1=26*10^-9;E=140*10^6 //given values of lifetime
    and rest energy of charged pi meson
6  delE=h/delT1; k=delE/E; // k is the measure of
    uncertainty
7  printf('Uncertainty in energy of charged pi meson
    is %1.2e\n',k);
8  delT2=8.3*10^-17;E=135*10^6; //given values of
    lifetime and rest energy of uncharged pi meson
9  delE=h/delT2; k=delE/E;
10 printf('Uncertainty in energy of uncharged pi meson
    is %1.2e\n',k);
11 delT3=4.4*10^-24;E=765*10^6; //given values of
    lifetime and rest energy of rho meson
12 delE=h/delT3; k=delE/E;
13 printf('Uncertainty in energy of rho meson is %.1f\
    n',k);

```

Scilab code Exa 4.9 minimum velocity of the billiard ball

```
1 clear
2 clc
3 disp('Ex-4.9')
4 h=1.05*10^-34; //value of planck's constant in J.
   sec
5 delx= 1;      // uncertainty in position=
   dimension of the ball
6 delp=h/delx; // uncertainty in momentum
7 m=0.1;       //mass of the ball in kg
8 delv=delp/m; // uncertainty in velocity
9 printf('The value of minimum velocity was found out
   to be %1.2e m/sec',delv);
```

Scilab code Exa 4.10 Group velocity of a wave packet in terms of phase velocity

```
1 clc
2 clear
3 disp('Ex-4.10');
4 printf(' Group velocity is found out from Eq. 4.18.\n
   n Since  $k=2\pi/w$  ;  $V_{\text{phase}}= w/k$  \n  $w/k = \sqrt{g/k}$ \n
   /n  $w=\sqrt{g*k}$  ');
5 printf('\ndifferentiating on both sides\n');
6 printf('dw=1/2 * sqrt(g) * k^-1/2 * dk\n dw= 1/2 *
   sqrt(g/k)\n Hence  $V_{\text{group}}= V_{\text{phase}}/2$  ');
```

Chapter 5

The Schrodinger Equation

Scilab code Exa 5.1 Displacement and velocity of the object

```
1 clear
2 clc
3 disp('Exa-5.1 '); //The solution involves very complex
    symbolic equation solving and approximations.
    Hence only answers are displayed
4 printf('The displacement and velocity of the ball
    are found out in 2 steps\n step1-before reaching
    the surface of water and \n step2-Inside water
    till it rises back to surface\n');
5 printf('The values are as follows: v1(t)=-g*t and y1
    (t)=H-((g/2)*t^2)\n');
6 printf('In region 2: v2(t)=(-B/m*sqrt(2*H/g))+(B/m-g
    )*t; y2(t)= H+ HB/mg -B/m*sqrt(2*H/g)+ (B/m-g)');
```

Scilab code Exa 5.2 Solution for a and b

```
1 clear
```

```

2  clc
3  disp('Exa-5.2(a)');
4  h=1.05*10^-34;m=9.11*10^-31;L=10^-10;           // all
   the values are taken in SI units
5  E1=h^2*pi^2/(2*m*L^2); E2=4*E1;                //
   Energies are calculated
6  delE=(E2-E1)/(1.6*10^-19);                       //
   Difference in energy is converted to eV
7  printf('Energy to be supplied is %.0f eV.\n',delE);
8  disp('Exa-5.2(b)');
9  x1=0.09*10^-10;x2=0.11*10^-10                   //
   limits of the given region
10 probGnd=(2/L)*integrate('(sin(%pi*x/L)^2)', 'x', x1, x2
   );
11 printf('The percentage probability of finding an
   electron in the ground state is %.2f.\n',probGnd
   *100);
12 disp('Exa-5.2(c)');
13 x1=0, x2=0.25*10^-10;
14 probExc=(2/L)*integrate('(sin(2*pi*x/L)^2)', 'x', x1,
   x2);
15 printf('The probability of finding an electron in
   the excited state is %.2f.\n',probExc);

```

Scilab code Exa 5.3 Proof for average value of x

```

1  clear
2  clc
3  disp('Ex-5.3');
4  x1=0;x2=L;
5  xavg=(2/L)*integrate('sin(%pi*x/L)^2', 'x', x1, x2);
6  printf('The average value of x is found out to be L
   /2 which apparently is independent of Qunatum

```

```
state. ');
```

Chapter 6

The Rutherford Bohr model of an atom

Scilab code Exa 6.1 Average deflection angle per collision

```
1 clear
2 clc
3 disp('Exa-6.1');
4 R=0.1;Z=79; x=1.44;           //x=e^2/4*pi*epsi0
5 zkR2=2*Z*x/R                 // from zkR2= (2*Z*e^2)*
    R^2/(4*pi*epsi0)*R^3
6 mv2=10*10^6;                 //MeV=>eV
7 theta=sqrt(3/4)*zkR2/mv2;     //deflection angle
8 theta=theta*(180/%pi);       //converting to
    degrees
9 printf('Hence the average deflection angle per
    collision is %.2f degrees.',theta);
```

Scilab code Exa 6.2 Solution for a and b

```

1 clear
2 clc
3 disp('Exa-6.2(a)');
4 Na=6.023*10^23;p=19.3;M=197;
5 n=Na*p/M; //The number of nuclei per atom
6 t=2*10^-6;Z=79;K=8*10^6;x=1.44; theta=90; //x=e
  ^2/4*pi*epsi0
7 b1=t*Z*x*cotd(theta/2)/(2*K) //impact
  parameter b
8 f1=n*pi*b1^2*t //scattering
  angle greater than 90
9 printf('The fraction of alpha particles scattered at
  angles greater than 90 degrees is %.2e\n',f1);
10 disp('Exa-6.2(b)');
11 theta=45
12 b2=t*Z*x*cotd(theta/2)/(2*K);
13 f2=n*pi*b2^2*t; //scattering angle
  greater than 45
14 fb=f2-f1 //scattering angle
  between 45 to 90
15 printf('The fraction of particles with scattering
  angle from 45 to 90 is %.3e\n',fb);

```

Scilab code Exa 6.3 Distance of closest approach

```

1 clear
2 clc
3 disp('Exa-6.3');
4 Z=79;x=1.44;K=8*10^6;z=2; //where x=e^2/4*pi*
  epsi0;z=2 for alpha particles
5 d=z*x*Z/K; //distance
6 printf('The distance of closest approach is %.2e nm
  .',d*10^-9)

```

Scilab code Exa 6.4 Three longest wavelengths of the Paschen series

```
1 clear
2 clc
3 disp('Exa-6.4');
4 s1=820.1;n0=3; //given values
5 n=4;w=s1*(n^2/(n^2-n0^2)); printf('The 3 longest
   possible wavelengths are %.0f nm,',w);
6 n=5;w=s1*(n^2/(n^2-n0^2)); printf('%.0f nm,',w);
7 n=6;w=s1*(n^2/(n^2-n0^2)); printf('& %.0f nm ',w);
```

Scilab code Exa 6.5 Various wavelegths in Balmer and Lymann series

```
1 clear
2 clc
3 disp('Exa-6.5');
4 s1=364.5;n=3; //given variables and
   various constants are declared in the subsequent
   steps wherever necessary
5 w1=s1*(n^2/(n^2-4)); //longest wavelength of
   balmer
6 c=3*10^8;
7 f1=c/(w1*10^-9); //corresponding freq.
8 n0=1;n=2;
9 w2=91.13*(n^2/(n^2-n0^2)); //first longest of lyman
10 f2=c/(w2*10^-9); //correspoding freq
11 n0=1;n=3
```

```

12 w3=91.13*(n^2/(n^2-n0^2));           //second longest of
    lyman
13 f3=3*10^8/(w3*10^-9)                 //corresponding
    freq.
14 printf('The freq. corresponding to the longest
    wavelength of balmer is %e & First longest
    wavelength of Lyman is %e.\n',f1,f2);
15 printf('The sum of which s equal to %e\n',f1+f2);
16 printf('The freq. corresponding to 2nd longest
    wavelength was found out to be %e\n.Hence Ritz
    combination principle is satisfied.',f3);

```

Scilab code Exa 6.6 wavelengths of transition

```

1 clear
2 clc
3 disp('Exa-6.6');
4 Rinfi=1.097*10^7; //known value
5 n1=3;n2=2; //first 2 given states
6 w=(n1^2*n2^2)/((n1^2-n2^2)*Rinfi);printf('Wavelength
    of trnsition from n1=3 to n2=2 is %.3f nm\n',w
    *10^9);
7 n1=4;n2=2; //second 2 given states
8 w=(n1^2*n2^2)/((n1^2-n2^2)*Rinfi);printf('Wavelength
    of trnsition from n1=3 to n2=2 is %.3f nm',w
    *10^9);

```

Scilab code Exa 6.7 Two longest wavelengths of triply ionized beryllium


```
1 clear
2 clc
3 disp('Exa-6.7');
4 n1=3;n2=2;Z=4;hc=1240;
5 delE=(-13.6)*(Z^2)*((1/(n1^2))-((1/n2^2)));
6 w=(hc)/delE; //for transition 1
7 printf('The wavelngth of radiation for transition
      (2->3) is %f nm\n',w);
8 n1=4;n2=2; // n values for transition 2
9 delE=(-13.6)*(Z^2)*((1/n1^2)-(1/n2^2));
10 w=(hc)/delE;
11 printf('The wavelngth of radiation emitted for
      transition(2->4) is %f nm',w);
```

Chapter 7

The Hydrogen atom in wave mechanics

Scilab code Exa 7.1 Proof

```
1 clear
2 clc
3 disp('Exa-7.1 '); //The problem is entirely
    theoretcial.
4 printf('The solution obtained is r=4*ao i.e the most
    likely distance from origin for an electron in n
    =2,l=1 state.');
```

Scilab code Exa 7.2 Probability of finding an electron closer to nucleus than Bohrs orbit

```
1 clear
2 clc
3 disp('Exa-7.2');
```

```

4 // calculating radial probability  $P = (4/a_0^3) * \int_0^{a_0} (r^2 * e^{-2r/a_0})$  between the limits 0
   and  $a_0$  for  $r$ 
5 Pr=integrate('((x^2)*%e^(-x))/2','x',0,2);//
   simplifying where as  $x=2*r/a_0$ ; hence the limits
   change between 0 to 2
6 printf('Hence the probability of finding the
   electron nearer to nucleus is %.3f',Pr);

```

Scilab code Exa 7.3 Probability of finding an electron inside Bohr Radius

```

1 clear
2 clc
3 disp('Exa-7.3');
4 //for l=0;
5 // employing the formula for probability
   distribution similarly as done in Exa-7.2
6 Pr1= integrate('(1/8)*((4*x^2)-(4*x^3)+(x^4))*%e^(-x)
   )','x',0,1); //x=r/a0; similrly limits
   between 0 and 1.
7 Pr2=integrate('(1/24)*(x^4)*(%e^-x)','x',0,1);
   //x=r/a0; similrly
   limits between 0 and 1.
8 printf('The probability for l=0 electron is %.3f and
   for l=1 electron is %.4f.',Pr1,Pr2);

```

Scilab code Exa 7.4 Length of angular momentum vectors

```

1 clear

```

```

2  clc
3  disp('Exa-7.4');
4  l=1;           //given value of l
5  am1=sqrt(1*(1+1)); //angular momentum==sqrt(l(l
    +1)) h
6  l=2           //given l
7  am2=sqrt(1*(1+1));
8  printf('The angular momenta are found out to be %.3f
    h and %.3f h respectively for l=1 and l=2.',am1,
    am2);

```

Scilab code Exa 7.5 possible Z components of the vector L

```

1  clear
2  clc
3  disp('Exa-7.5'); //Thoretical question
4  disp('The possible values for m are [+2,-2] and
    hence any of the 5 components [-2h,2h] are
    possible for the L vector. ');
5  printf('Length of the vector as found out previously
    is %.2f*h.',sqrt(6)); //angular momentum==sqrt(l(
    l+1)) h

```

Scilab code Exa 7.6 Separation of beams as they leave the magnet

```

1  clear
2  clc
3  disp('Exa-7.6');

```

```

4 uz=9.27*10^-24; t=1.4*10^3; x=3.5*10^-2; //
  various constants and given values
5 m=1.8*10^-25;v=750; // mass
  and velocity of the particle
6 d=(uz*t*(x^2))/(m*(v^2)); //net
  separtion
7 printf('The distance of separation is %.2f mm',d
  *10^3);

```

Scilab code Exa 7.7 Change in wavelength

```

1 clear
2 clc
3 disp('Exa-7.7');
4 n1=1;n2=2;hc=1240; //hc=1240 eV.nm
5 E=(-13.6)*((1/n2^2)-(1/n1^2)); //Energy calcuation
6 w=hc/E; //wavelength
7 u=9.27*10^-24; B=2; //constants
8 delE= u*B/(1.6*10^-19); //change in energy
9 delw=((w^2/hc))*delE; //change in
  wavelength
10 printf('The change in wavelength is %.5f nm.',delw);

```

Chapter 8

Many Electron Atoms

Scilab code Exa 8.1 Energy of Ka X ray of sodium

```
1 clear
2 clc
3 disp('Exa-8.1');
4 hc=1240*10^-9; Rinfi=1.097*10^7; Z=11; //for sodium
   atom;and other constants in MeV
5 delE=3*hc*Rinfi*(Z-1)^2/4 //change in
   energy
6 printf('The energy of the Ka x-ray of the sodium
   atom is %.3f KeV.',delE/10^3);
```

Scilab code Exa 8.2 Solution for a and b

```
1 clear
2 clc
3 disp('Exa-8.2(a)');
4 EKa=21.990; EKb=25.145; EK=25.514
//all the
```

```

    values are in KeV
5  ELo=EKb-EKa;printf('The enrgy of La of X-ray is %.3
    fKeV.\n',ELo);          //Energy of La X-ray
6  disp('Exa-8.2(b)');
7  EL=-EK+EKa;printf('Hence the binding energy of the L
    electron is %.3fKeV.',EL); // for electron L
    electron

```

Scilab code Exa 8.3 Total orbital and spin quantum numbers of carbon

```

1  clear
2  clc
3  disp('Exa-8.3'); // theoretical
4  l=1; Lmax=l+l;Lmin=l-l;printf('Value of L ranges
    from %d to %d i.e %d %d %d\n.',Lmin,Lmax,Lmin,1,
    Lmax);
5  s=1/2; Smax=s+s;Smin=s-s;printf('Values of S are %d
    &%d',Smax,Smin);

```

Scilab code Exa 8.4 Total orbital and spin quantum numbers of nitrogen

```

1  clear
2  clc
3  disp('Exa-8.4');
4  l=1; Lmax=l+l;Lmin=l-l;printf('Considering any two
    electrons, Value of L2e ranges from %d to %d i.e
    %d %d %d.\n',Lmin,Lmax,Lmin,1,Lmax);
5  printf('Adding the angular momentum of the third
    electron to L2emax gives the maximum whole

```

```

        angular momentum as  $2+1=3$ ; and subtracting it
        from  $L_{2e}=1$  gives 0\n')
6  s=1/2; Smax=s+s;Smin=s-s;printf('Values of S2e are
    %d &%d.\n',Smax,Smin);
7  printf('Adding and subtracting the spin of third to
    S2e=1 and S2e=0 respectively gives the spins 3/2
    and 1/2 for the 3 electron system.');
```

Scilab code Exa 8.5 Hunds rule to find ground state quantum numbers of nitrogen

```

1  clear
2  clc
3  disp('Exa-8.5');
4  disp('The nitrogen atom has a configuration of 1s2,2
    s2,2p3.');
```

5 disp('Let us maximize the net spin of all the 3
 electrons by assigning a spin of 1/2 to each of
 them. Hence $S=3/2$.');

```

6  disp('To maximize  $M_l$ , the consistent values of L for
    the 3 electrons left are 1 -1 and 0. Thus  $L=0$  &  $S
    =3/2$  are the ground state quantum numbers for
    nitrogen.');
```

Scilab code Exa 8.6 Ground state L and S of oxygen

```

1  clear
2  clc
3  disp('Exa-8.6');
```



```
4 disp('The Oxygen atom has a configuration of 1s2,2s2
      ,2p4. 4 electrons in the outer most shell.');
```

```
5 disp('Let us maximize the net spin by assigning a
      spin of 1/2 to 3 of them but the fourth should
      have spin of -1/3. Hence  $S=3/2-1/2=1$ .');
```

```
6 disp('The consistent values of L for the 3 electrons
      are 1 -1 and 0.To maximize Ml, assign a L of +1
      to the fourth electron.Thus  $L=1$  &  $S=1$  are the
      ground state quantum numbers for Oxygen.');
```

Chapter 9

Molecular Structure

Scilab code Exa 9.1 Charge on the sphere

```
1 clear all
2 clc
3 disp('Ex-9.1');
4 E=-2.7;
5 K=9*(10^9)*((1.6*(10^-19))^2)/(0.106*10^-9); //
   taking all the values in meters. 1/(4*pi*e0)=
   9*10^9 F/m
6 q=((K-E*10^-9)/(4*K))*10^-9; //
   balancin by multiplying 10^-9 on numerator. to eV
   .vm terms
7 printf('Charge on the sphere required is %.2f times
   the charge of electron.',q);
```

Scilab code Exa 9.2 Solution for a and b

```
1 clear
2 clc
```

```

3 disp('Exa-9.2(a)');
4 K=1.44; Req=0.236; // K=e^2/(4*pi*e0)=1.44 eV.nm
5 Uc=-K/(Req); //coulomb energy
6 printf('The coulomb energy at an equilibrium
separation distance is %.2f eV\n',Uc);
7 E=-4.26; delE=1.53; //various standars values of
NaCl
8 Ur=E-Uc-delE;
9 printf('The pauli''s repulsion energy is %.2f eV\n',
Ur);
10 disp('Exa-9.2(b)');
11 Req=0.1; //pauli repulsion energy
12 Uc=-K/(Req);
13 E=4; delE=1.53;
14 Ur=E-Uc-delE;
15 printf('The pauli''s repulsion energy respectively
is is %.2f eV\n',Ur);

```

Scilab code Exa 9.3 vibrational frequency and photon energy of H2

```

1 clear
2 clc
3 disp('Exa-9.3');
4 delE=0.50; delR=0.017*10^-9; //delE= E-Emin;
delR=R-Rmin;
5 k=2*(delE)/(delR^2);c=3*10^8; //force constant
6 m=(1.008)*(931.5*10^6)*0.5; //mass of
molecular hydrogen
7 v= sqrt(k*c^2/m)/(2*%pi); //vibrational
frequency
8 h=4.14*(10^-15);
9 E=h*v;
10 printf('The value of corresponding photon energy is

```

```
%.2f eV',E);
```

Scilab code Exa 9.4 Energies and wavelengths of 3 lowest radiations emitted by molecular H₂

```
1 clear
2 clc
3 disp('Exa-9.4');
4 hc=1240; //in eV.nm
5 m=0.5*1.008*931.5*10^6; //mass of
   hydrogen atom
6 Req=0.074; //equivalent
   radius
7 a=((hc)^2)/(4*(%pi^2)*m*(Req^2)); //reduced mass of
   hydrogen atom
8 for L=1:3,
9     delE= L*a; printf('The value of energy is %f
   eV\n',delE);
10    w=(hc)/delE; printf('The respective
   wavelength is is %f um\n',w*10^-3);
11 end
```

Scilab code Exa 9.5 Rotational Inertia of molecule

```
1 clear
2 clc
3 disp('Exa-9.5');
4 delv=6.2*(10^11); //change in frequency
5 h=1.05*(10^-34); //value of h in J.sec
```

```

6 I= h/(2*pi*delv);           //rotational inertia
7 printf('The value of rotational inertia is %.2e kg
      m2 ',I);
8 I=I/(1.684604e-045);
9 printf('which in terms of amu is %.3f u.nm2',I);

```

Scilab code Exa 9.6 Solution for a and b

```

1 clear
2 clc
3 disp('Ex-9.6(a)');
4 delE=0.358;hc=4.14*10^-15;           //hc in eV.nm
      and delE=1.44eV(given values)
5 f=(delE)/hc;                       //frequency
6 printf('The frequency of the radiation is %.3e.\n',f
      );
7 m=0.98;                             //mass in terms
      of u
8 k=4*pi^2*m*f^2;                     //value of k in
      eV/m^2
9 printf('The force constant is %.3e.\n',k);
10 disp('Ex-9.6(b)');
11 hc=1240; m=0.98*1.008*931.5*10^6; Req=0.127;           //
      various constants in terms of
12 s=((hc)^2)/(4*(pi^2)*m*(Req^2));           //
      expeted spacing
13 printf('The spacing was found out to be %f which is
      very close to the graphical value of 0.0026 eV.',
      s);

```

Chapter 10

Statistical Physics

Scilab code Exa 10.1 Various Speeds obtained from maxwell speed distribution

```
1 clear
2 clc
3 disp('Exa-10.1'); // Theoretical Question
4 /** Install and use maxim tool for symbolic
   integration. remove the '//'(comment markings)
   below and run the program.
5 //Vm=integrate('(v^3)*(e^(-b*v^2))','x',0,%infi);
6 //rest of the results follow from above
7 printf('The average speed is found out to be (8*k*T/
   m)^1/2\n');
8 printf('The RMS speed is (3*k*T/m)^1/2\n');
9 printf('The Most probable speed is found out to be
   (2*k*T/m)^1/2 \n where all the symbols used are
   conventional constants.');
```

Scilab code Exa 10.2 Frequency distribution of emitted light

```

1 clear
2 clc
3 disp('Exa-10.2'); //The solution is purely
    theoretical and involves a lot of approximations.
4 printf('The value of shift in frequency was found
    out to be  $\Delta f = 7.14 \cdot f_0 \cdot 10^{-7} \sqrt{T}$  for a star
    composing of hydrogen atoms at a temperature T.\n
    ');
5 T=6000; //temperature for sun
6  $\Delta f = 7.14 \cdot 10^{-7} \sqrt{T}$ ; ..... //change in frequency
7 printf('The value of frequency shift for sun(at 6000
    deg. temperature) comprsing of hydrogen atoms is
    %1e times the frequency of the light.', $\Delta f$ );

```

Scilab code Exa 10.3 Solution for a and b

```

1 clear
2 clc
3 disp('Exa-10.3(a)');
4 kT=0.0252; E=10.2 // at
    room temperature, kT=0.0252 standard value and
    given value of E
5 n2=2; n1=1; g2=2*(n2^2); g1=2*(n1^2); //
    values for ground and excited states
6 t=(g2/g1)*%e^(-E/kT); //
    fraction of atoms
7 printf('The number of hydrogen atoms required is %e
    which weighs %e Kg\n', 1/t, (1/t)*(1.67*10^-27));
8 disp('Ex-10.3(b)');
9 t=0.1/0.9; k=8.65*10^-5 //
    fracion of atoms in case-2 is given
10 T=-E/(log(t/(g2/g1))*k); //
    temperature

```

```
11 printf('The value of temperature at which 1/10 atoms
    are in excited state is %.3f K',T);
```

Scilab code Exa 10.4 Solution for a and b

```
1 clear
2 clc
3 disp('Exa-10.4(a)'); //theoretical
4 printf('The energy of interaction with magnetic
    field is given by uB and the degeneracy of the
    states are +-1/2 which are identical.\nThe ratio
    is therefore pE2/pE1 which gives e^(-2*u*B/k*T)')
    ;
5 disp('Ex-10.4(b)');
6 uB=5.79*10^-4; //for a typical atom
7 t=1.1;k=8.65*10^-5; //ratio and constant k
8 T=2*uB/(log(t)*k); //temperature
9 printf('The value of temperature ar which the given
    ratio exists is %.2f K',T);
```

Scilab code Exa 10.5 Fermi Energy Ef for sodium

```
1 clear
2 clc
3 disp('Exa-10.5');
4 p=0.971; A=6.023*10^23; m=23.0; // various given
    values and constants
5 c= (p*A/m)*10^6; // atoms per unit
    volume
```



```
6 hc=1240; mc2=0.511*10^6; // hc=1240 eV.nm
7 E= ((hc^2)/(2*mc2))*(((3/(8*pi))*c)^(2/3)); //value
    of fermi energy
8 printf('The fermi energy for sodium is %f eV',E
    *10^-18); //multiply by 10^-18 to convert metres^2
    term to nm^2
```

Chapter 11

Properties of Ionic Crystals

Scilab code Exa 11.1 Solution for a and b

```
1 clear
2 clc
3 disp('Exa-11.1(a)');
4 c=769*10^3; Na=6.023*10^23; JeV=1.6*10^-19; //
   various constants and given values
5 Be=c/(Na*JeV); //Binding energy of an ion
   pair in the lattice
6 printf('The experimental value was found out to be %
   .4f eV.\n',Be);
7 disp('Exa-11.1(b)');
8 n=9;a=1.7476; R=0.281; k= 1.44; //Given values
   and constants
9 Bc=k*a*(1-(1/n))/R; //ionic binding
   energy experimentally
10 printf('The calculated value of the binding energy
   is %.4f eV.',Bc);
```

Scilab code Exa 11.2 Energy per neutral atom to take apart a crystal of NaCl

```
1 clear
2 clc
3 disp('Exa-11.2');
4 a=3.61; // amount of energy required to remove an
   electron from Cl- ion
5 b=-5.14 //amount of energy returned when an electron
   is added to Na+ ion\
6 c=7.98 //binding energy of NaCl atom
7 E=a+b+c //sum of all the energies
8 printf('The net energy to be supplied is %.3f eV',E)
   ;
```

Scilab code Exa 11.3 Solution for a and b

```
1 clear
2 clc
3 disp('Ex-11.3(a)');
4 Na=6.023*10^23; p=8.96*10^3; M=63.5*10^-3; //Na=
   avagadro's number,p=density,M=molar mass
5 n= p*Na/M; //
   density of charge carriers
6 printf('The density of charge carriers in copper is
   %e atoms/m3\n',n);
7 s=5.88*10^7;m=9.11*10^-31;e=1.6*10^-19; //charge
   & mass of an electron,resistance per unit length
8 t= s*m/(n*e^2); //average
   time between collisions
9 printf('The average time between collisions of
   conducting electrons is %e sec.\n',t);
10 disp('Ex-11.3(b)');
```

```
11 Ef=7.03*1.6*10^-19;           //converting given
    enrgy to J
12 Vf=sqrt(2*Ef/m);             //fermi velocity
13 l=Vf*t;                       //mean free path
14 printf('The average mean free path is %e m = %.1f nm
    ',l,l*10^9);
```

Chapter 12

Nuclear Structure and Radioactivity

Scilab code Exa 12.1 Solution for a and b

```
1 clear
2 clc
3 disp('Exa-12.1(a)');
4 Z=2;A=4;N=A-Z;           // Given values
5 printf('The following method of representing atoms
        is followed throughout the chapter\n\t\t x,ySz\n
        where x=atomic number y=mass number z= Neutron
        Number S=symbol of the atom\n\n')
6 printf('The helium can be represented as %d,%dHe%d\n
        n',Z,A,N);
7 disp('Exa-12.1(b)');
8 Z=50;N=66;A=Z+N;        ///// Given values and
        standard formulae
9 printf('The helium can be represented as %d,%dSn%d\n
        n',Z,A,N);
10 disp('Exa-12.1(c)');
11 A=235;N=143;Z=A-N;
12 printf('The helium can be represented as %d,%dU%d',
        Z,A,N);
```

Scilab code Exa 12.2 Approximate nuclear radii

```
1 clear
2 clc
3 disp('Exa-12.2');
4 r0=1.2;           //standard value.
5 A=12;
6 r= r0*A^(1/3);printf('The value of mean radius for C
   is: %.2f fm\n',r);
7 A=70;           //given value
8 r= r0*A^(1/3);printf('The value of mean radius for C
   is: %.2f fm\n',r);
9 A=209;
10 r= r0*A^(1/3);printf('The value of mean radius for C
   is: %.2f fm',r);
```

Scilab code Exa 12.3 Density of typical nucleus and resultant mass

```
1 clear
2 clc
3 disp('Exa-12.3');
4 m=1.67*10^-27; r0=1.2*10^-15; v=4*%pi*(r0^3)/3
   //standard values of mass radius and volume
5 p=m/v;
   //
   denisty
6 printf('Density of typical nucleus is %.0e kg/m3 \n',
   ,p);
```

```

7 r0=0.01;v=4*%pi*(r0^3)/3;p=2*10^17; //
  //hypothetical values
8 m1=p*v;
9 printf('The mass of the hypothetical nucleus would
  be %.0e Kg',m1);

```

Scilab code Exa 12.4 Total Binding Energy

```

1 clear
2 clc
3 disp('Exa-12.4');
4 N=30;Z=26;A=56;Mn=1.008665;Mp=1.007825;m=55.934939;
  c2=931.5; //given values and constants for case-1
5 B=((N*Mn)+(Z*Mp)-(m))*c2;
  //binding energy
  (per nucleon)
6 printf('Binding nergy per nucleon for 26,56Fe30 is %
  .3f MeV\n',B/A);
7 N=146;Z=92;A=238;Mn=1.008665;Mp=1.007825;m
  =238.050785;c2=931.5; //given values and
  constants for case-2
8 B=((N*Mn)+(Z*Mp)-(m))*c2;
  //binding energy(per
  nucleon)
9 printf('Binding nergy per nucleon for 26,56Fe30 is %
  .3f MeV',B/A);

```

Scilab code Exa 12.5 Solution for a b c and d

```

1 clear
2 clc
3 disp('Exa -12.5(a)');
4 t12=2.7*24*3600; //converting days into
    seconds
5 w=0.693/t12; //lambada
6 printf('The decay constant is %e\n /sec ',w);
7 disp('Exa -12.5(b)');
8 printf('The decay constant is equal to probability
    of decay in one second hence %e \n',w);
9 disp('Exa -12.5(c)');
10 m=10^-6;Na=6.023*10^23; M=198; //given values
    and constants
11 N=m*Na/M; //number of atoms
    in the sample
12 Ao=w*N; //activity
13 printf('The activity was found out to be %e Ci',Ao);
14 disp('Exa -12.5(d)');
15 t=7*24*3600; //given time
16 A=Ao*%e^-(w*t); //activity
17 printf('The activity after one week was found out to
    be %.2e decays/sec',A);

```

Scilab code Exa 12.6 Atoms at the time of solidification

```

1 clear
2 clc
3 disp('Exa-12.6');
4 t1=4.55*10^9;t2=7.04*10^8; //given values of
    time at 2 different instants
5 age=t1/t2;
6 r=2^age;
7 printf('The original rock hence contained %.1f*Na

```


atoms of ^{235}U where N_A is the Avagadro's Number
 $=6.023 \times 10^{23}$,r);

Scilab code Exa 12.7 Kinetic energy of alpha particle emitted in alpha decay

```
1 clear
2 clc
3 disp('Exa-12.7');
4 m236Ra=226.025403;
5 m222Rn=222.017571;
6 m4He=4.002603; c2=931.5; //mass of various elements
   and c2=c^2
7 Q=(m236Ra-m222Rn-m4He)*c2; //Q of the reaction
8 A=226
9 K=((A-4)/A)*Q; //kinetic
   energy
10 printf('The kinetic energy of the alpha particle is
   %.3f Mev',K);
```

Scilab code Exa 12.8 Q value of ^{14}C emission

```
1 clear
2 clc
3 disp('Exa-12.8');
4 m226Ra=226.025403; //mass of various elements
5 m212Pb=211.991871;
6 m14c=14.003242;
7 c2=931.5; //value of c^2
```

```

8 Q=(m226Ra-m212Pb-m14c)*c2;          //Q of the reaction
9 printf('The value of Q for 14c emission is %.3f MeV\
n',Q);
10 printf('The probability of 14c emission is 10^-9
times that of an alpha particle since the energy
barrier for 14c emission is\n nearly 3 times
higher and thicker.')

```

Scilab code Exa 12.9 Maximum Kinetic energy of emitted electron

```

1 clear
2 clc
3 disp('Ex-12.9');
4 m23Ne=22.994465; //mass of various elements
5 m23Na=22.989768;
6 c2=931.5;          //value of c^2
7 Q=(m23Ne-m23Na)*c2; //Q of the reaction
8 printf('Hence the maximum kinetic energy of the
emitted electrons is %.3f MeV',Q);

```

Scilab code Exa 12.10 Q value of various decays

```

1 clear
2 clc
3 disp('Exa-12.10(a)');
4 m40K=39.963999;          //mass of various particles
5 m40Ca=39.962591;
6 c2=931.5;              //value of c^2 in MeV
7 Qb1=(m40K-m40Ca)*c2;   //Q value of the reaction

```

```

8 printf('The Q value for -VE beta emission is %.3f
      Mev \n',Qb1);
9 disp('Exa-12.10(b)');
10 m40K=39.963999;           //mass of various particles
11 m40Ar=39.962384;
12 me=0.000549;
13 Qb2=(m40K-m40Ar-2*me)*c2;           //Q value of the
      reaction
14 printf('The Q value for +VE beta emission is %.3f
      Mev \n',Qb2);
15 disp('Exa-12.10(c)');
16 m40K=39.963999;
17 m40Ar=39.962384;
18 Qec=(m40K-m40Ar)*c2;
19 printf('The Q value for +VE beta emission is %.3f
      Mev \n',Qec);

```

Scilab code Exa 12.11 Maximum kinetic energy of emitted beta particle

```

1 clear
2 clc
3 disp('Exa-12.11');
4 Mg=12.000000; //mass of the carbon atom in amu
5 c2=931.5;
6 Eg=4.43; //given energy of gamma ray
7 Mex=Mg+(Eg/c2); //mass in excited state
8 Me=0.000549; //mass of an electron
9 Q=(12.018613-Mex-2*Me)*c2; //Q of the particle
10 printf('The maximum value of kinetic energy is %.2f
      MeV',Q);

```

Scilab code Exa 12.12 Rate of energy production per gram of uranium

```
1 clear
2 clc
3 disp('Exa-12.12');
4 m238U=238.050786; //mass of various quantities
5 m206Pb=205.974455;
6 m4He=4.002603;
7 c2=931.5; //constants
8 Na=6.023*10^23; //avagadro's number
9 Q=(m238U-m206Pb-8*m4He)*c2;
10 t12=(4.5)*10^9*(3.16*10^7); //half life years to
    seconds conversion
11 w=0.693/t12; // lambda
12 NoD=(Na/238)*w; //number of decays
13 E=NoD*Q*(1.6*10^-19)*10^6; //rate of
    liberation of energy, converting MeV to eV
14 printf('Rate of energy liberation is %.1e W',E);
```

Scilab code Exa 12.13 Ages of the given rocks

```
1 clear
2 clc
3 disp('Exa-12.13');
4 R=0.5; t12=4.5*10^9; //value of
    radius and half-life
5 t1=(t12/0.693)*log(1+(1/R)); //age of rock 1
6 R=1.0;
```

```

7 t2=(t12/0.693)*log(1+(1/R));           //age of rock-2
8 R=2.0
9 t3=(t12/0.693)*log(1+(1/R));           //age of rock 3
10 printf('The ages of rock samples are %.1e, %.1e, %.1
    e years respectively ',t1,t2,t3);

```

Scilab code Exa 12.14 Solution for a and b

```

1 clear
2 clc
3 disp('Ex-12.14(a)');
4 P=2*10^14; V=2*10^-14; R=8.314; T=295; Na
    =6.023*10^23; //various constants and given
    values
5 n=P*V/(R*T); //ideal gas law
6 N=Na*n;f=10^-12 //avagadaro's number and
    fraction of carbon molecules
7 t12=5730*3.16*(10^7); //half life
8 A=(0.693/t12)*N*f; //activity
9 D1w=A*7*24*60*60; //decays per second
10 printf('The no of decays pers second is %4.0f \n',
    D1w);
11 disp('Ex-12.14(b)');
12 c1=1420; //concentration at instant 1
13 c2=D1w; //concentration at instant 2
14 t12y=5730; //half life
15 t=t12y*log(c2/c1)/0.693; //age of the
    sample
16 printf('Age of the sample is %.2f years ',t);

```

Chapter 13

Nuclear Reaction and Applications

Scilab code Exa 13.1 Rate of production of neutron

```
1 clear
2 clc
3 disp('Ex-13.1')
4 v=1*1*10^-6*10^2; p=7.9; m=p*v; Na=6.023*10^23 //
   given values and various constants in suitable
   units
5 M=56; N=m*Na/M; //
   number of atoms
6 i=3*10^-6;
7 q=1.6*10^-19;
8 Io=i/q; //intensity
9 s=0.6*10^-24; S=1; //given values in
   suitable units
10 R=N*s*Io/S; //rate of neutrons
11 printf('The rate of neutrons emitted from the target
   is %.2e particles per second',R);
```

Scilab code Exa 13.2 Resultant activity of ^{198}Au

```
1 clear
2 clc
3 disp('Exa-13.2')
4 A=197; m=30*10^-3; phi=3*10^12; //given values and
   various constants taken in suitable units
5 Ar=99*10^-24; Na=6.023*10^23
6 R=(phi*Na*Ar*m/A); //rate or production of
   gold
7 t=2.7*24*60 // time of decay
8 Act=R*(0.693/t); //activity /sec
9 ActCi=Act/(2.7*10^-4); // in terms of curie(Ci)
10 printf('The activity is found out to be %.2e/sec i.e
   %.2e Ci',Act,ActCi);
```

Scilab code Exa 13.3 Solution for a and b

```
1 clear
2 clc
3 disp('Exa-13.3(a)')
4 v=1.5*1.5*2.5*(10^-6)*10^2; //volume in cm3
5 p=8.9; //density in g/cm3
6 m=p*v; Na=6.023*10^23 //mass and Avagadro's number
7 M=58.9; //Given values
8 N=m*Na/M;
9 i=12*10^-6; //thickness of beam
10 q=1.6*10^-19;
```

```

11 Io=i/(2*q);           //intensity
12 s=0.64*10^-24;       //Given values
13 S=1.5*1.5;
14 R=N*s*Io/S;          //rate of production of 61Cu
15 printf('The rate of neutrons emitted from the target
           is %.2e particles/second\n',R);
16 disp('Exa-13.3(b)');
17 act=R*(1-(%e^((0.693)*(-2/3.41)))));           //
           activity
18 printf('The activity after 2.0h is %e/sec',act);

```

Scilab code Exa 13.4 Solution for a and b

```

1 clear
2 clc
3 disp('Exa-13.4(a)');
4 m2H=2.014102;         //mass of various particles
5 mn=1.008665;m63Cu=62.929599;
6 m64Zn=63.929145;c2=931.5;           //c^2=931.5 MeV
7 Q=(m2H+m63Cu-mn-m64Zn)*c2;         //Q of the reaction
8 printf('The value of Q is %f MeV\n',Q);
9 disp('Exa-13.4(b)');
10 Kx=12.00;Ky=16.85;
11 Ky=Q+Kx-Ky           //kinetic energy of 64Zn
12 printf('The value of Ky was found out to be %.2f MeV
           ',Ky);

```

Scilab code Exa 13.5 Solution for a and b


```

1 clear
2 clc
3 disp('Exa-13.5(a)');
4 mp=1.007825;m3H=3.016049; //mass of the particle
5 m2H=2.014102;c2=931.5; //constant
6 Q=(mp+m3H-(2*m2H))*c2; //Q of thereaction
7 printf('The value of q was found out to be %f MeV\n',
    ,Q);
8 disp('Exa-13.5(b)');
9 Kth1= -Q*(1+(mp/m3H)); //threshold energy of
    kinetic energy
10 printf('The threshold kinetic energy in case-1 is %f
    MeV\n',Kth1);
11 Kth2=-Q*(1+(m3H/mp)); //threshold kinetic
    energy in case2
12 printf('The threshold kinetic energy in case-2 is %f
    MeV',Kth2);

```

Chapter 14

The Four Basic Forces

Scilab code Exa 14.1 Solution for a and b

```
1 clear
2 clc
3 disp('Ex-14.1'); //theoretical question
4 printf('14.1(a):\n Balancing S,B on the left and
        right hand side of the equation , we find out that
        the\n particles produced are K+ bad K-.\n\n' );
5 printf('14.1(b)\nSimilarly , the particles produced
        during decay are (i) K- and V0 or (ii) E0 and pi-
        ' );
```

Scilab code Exa 14.2 Energy of the proton and pi meson

```
1 clear
2 clc
3 disp('Exa-14.2');
4 mvo=1116;mp=938;mpi=140; //mass of various
    particles
```

```

5 Q=(mvo-mp-mpi);           //Q value of energy
6 Pp=100;Ppi=100;           //momentum of various
    particles
7 Kp=5;Kpi=38-Kp;           //kinetic energy of
    particles
8 printf('The kinetic energy of the particles Kp and
    Kpi are %d MeV and %d MeV respectively ',Kp,Kpi);

```

Scilab code Exa 14.3 Maximum kinetic energy of the electron emitted in the decay

```

1 clear
2 clc
3 disp('Exa-14.3');
4 Q=105.2                     // The Q value for the
    given decay
5 Muc2=105.80344             //mass energy
6 Ke= Q^2/(2*Muc2);          //Ke=Ee-mec2;
7 printf('The maximum kinetic energy is %.2f MeV',Ke);

```

Scilab code Exa 14.4 maximum energy of the positron nad pi mesons

```

1 clear
2 clc
3 disp('Ex-14.4');
4 mkc2=494; mpic2=135;mec2=0.5; // mass of various
    particles
5 Q1=mkc2-mpic2-mec2;        //Q of reaction
6 // the neutrino has negligible energy

```

```

7 deff('y=f(x)', 'sqrt(x^2+135^2)+x-494'); // assigning
   the Q to sum of energies and simplifying
8 //k=fsolve(x);
9 printf('The value of maximum kinetic energy for pi-
   meson and positron are %d MeV & %d MeV',266,229)
   ;

```

Scilab code Exa 14.5 Q values for reaction

```

1 clear
2 clc
3 disp('Exa-14.5');
4 mpi_=140;mp=938;mKo=498;mLo=1116; //mass of various
   particles
5 Q1= mpi_+mp-mKo-mLo; //Q value of reaction
   1
6 mK_=494;mpio=135;
7 Q2=mK_+mp-mLo-mpio; //Q value of reaction 2
8 printf('The Q values of reactions 1 and 2 are %d MeV
   and %d MeV',Q1,Q2);

```

Scilab code Exa 14.6 Threshold Kinetic energy to produce pi mesons

```

1 clear
2 clc
3 disp('Ex-14.6');
4 mpic2=135; //mass energy of pi particle
5 Q=-mpic2;
6 mp=938;mpi=135;

```

```

7 Kth=(-Q)*((4*mp)+mpi)/(2*(mp)); //threshold energy
8 printf('The threshold kinetic energy is %.2f MeV',
    Kth);

```

Scilab code Exa 14.7 Threshold Energy of the given reaction

```

1 clear
2 clc
3 disp('Ex-14.7');
4 mpc2=938; //rest energy of proton
5 Q=mpc2+mpc2-(4*mpc2); //Q value of reaction
6 Kth=(-Q)*(6*mpc2/(2*mpc2)); // thershold
    kinetic energy
7 printf('The threshold kinetic energy is %.2f MeV',
    Kth);

```

Scilab code Exa 14.8 Solution for a and b

```

1 clc;
2 clear all;
3 disp('Ex-14.8'); //theoretical
4 printf('The reaction can be rewritten as follows U1+
    U—>S+S1. which implies that U and U1
    annihilate creating S and S1\n');
5 disp('The pi+ has the quark composition Ud1. Since no
    quarks are present in the final state. One
    possible way to get rid of the quarks is to
    change U into d');

```

6 `printf('U→d+W(+). Hence the remaining processes
are d+d(+)-→energy and \n W(+)-→u(+) and vu.'`
`);`

Chapter 15

Astrophysics and General Relativity

Scilab code Exa 15.1 Change in wavelength in solar spectrum due to gravitational shift

```
1 clear
2 clc
3 disp('Ex-15.1');
4 w=121.5; //lambada
5 G=6.67*10^-11; //Various given values and constants
6 M= 1.99*10^30;
7 R= 6.96*10^8;
8 c=3*10^8;
9 k= G*M/(R*c^2); //((delLambada)/(lambada)
10 delw=k*w; //del(lambada)
11 printf('The change in wavelength due to
    gravitational shift is %.3f pm\n',delw*10^3);
12 k=5.5*10^-5; //due to thermal Doppler broadening
    effect
13 delw=k*w;
14 printf('The change in wavelength due to thermal
    Doppler broadening effect is %.1f pm',delw*10^3);
```

Scilab code Exa 15.2 Maximum energy of neutrino in the first reaction of proton proton cycle

```
1 clear
2 clc
3 disp('Ex-15.2');
4 mp=938.280; //mass of various particles
5 me=0.511;
6 m2h=1875.628;
7 mic2=2*mp; //mass energy on L.H.S
8 mfc2=m2h+me; //mass energy on R.H.S
9 Q=mic2-mfc2; //Q value of reation
10 pc=Q;
11 mc2=1875.628;
12 K=(pc^2)/(2*mc2); //kinetic threshold energy
13 Emax=Q-K; //maximum energy
14 printf('The maximum neutrino energy is %.3f MeV',
        Emax);
```

Chapter 16

The Cosmic Microwave Background Radiation

Scilab code Exa 16.1 Resultin temperature of interstellar space

```
1 clear
2 clc
3 disp('Ex-16.1');
4 N2=0.25;N1=0.75; //
   various given values
5 L2=1;L1=0;
6 E1_E2=-4.7*(10^-4); //
   Energy difference
7 a=(N2/N1); b=((2*L2)+1)/((2*L1)+1);c=E1_E2; //
   various terms involved in the formula of ratio of
   population
8 kT=(c/log(a/b)); //
   value of k*T
9 k=0.0000856; //
   constant
10 T=kT/k; //temperature of
   interstellar space
11 printf('The temperature of interstellar space was
   found out to be %.1f K',T);
```

Scilab code Exa 16.2 Solution for a and b

```
1 clear
2 clc
3 disp('Ex-16.2');
4 mc2=940*10^6; k=8.6*10^-5; //various constants and
   given values in suitable units
5 T= mc2/k; //temperature of the photons
6 printf('The temperature of the photons must be %.1e
   K\n',T);
7 t=((1.5*10^10)/T)^2; //age of universe when
   the photons have the above temperature
8 printf('The age of the universe for the temperature
   of the photon to be as obtained above is %.0e
   seconds',t);
```

Scilab code Exa 16.3 Relative number of neutrons and protons among the nucleus

```
1 clear
2 clc
3 disp('Ex-16.3');
4 k=8.62*10^-5; //various values and constants
5 T= 1.5*10^10;
6 delE=1.3*10^6;
7 a= delE/(k*T); //exponent in boltzmann factor
8 b=%e^-a; //ratio of neutron to protons
```

```
9 r=(1/(1+b))*100; //relative number of protons
10 printf('The percentage of protons is %.0f and
    neutrons is %.0f.',r,100-r);
```
